

G0.0 + 0.0

Sgr A East

RA: 17^h45^m44^s**1-GHz flux/Jy:** 100?**Size/arcmin:** 3.5×2.5**Dec:** −29°00′**Spectral index:** 0.8?**Type:** S

Radio: Non-thermal shell, in complex region, interacting with molecular material to the west.

X-ray: Diffuse emission, centrally peaked.

Point sources: Compact X-ray/radio source.

References:

Ekers *et al.* 1983, A&A, 122, 143. VLA at 1.4 GHz and 5 GHz (both 5''×8''), $S_{1.4\text{ GHz}} = 77\text{ Jy}$, $S_{5.0\text{ GHz}} = 31\text{ Jy}$.

Pedlar *et al.* 1989, ApJ, 342, 769. VLA at 332 MHz (12''), 1.4 GHz (1''3×2''5) and 5 GHz (1''3×2''5).

Mezger *et al.* 1989, A&A, 209, 337. Nearby molecular material.

Ho *et al.* 1991, Nature, 350, 309. VLA of NH₃ emission from surroundings.

Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (17''×33'').

Serabyn *et al.* 1992, ApJ, 395, 166. Nearby molecular material.

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×88'').

Yusef-Zadeh & Mehringer 1995, ApJ, 452, L37. VLA of nearby H₂O masers.

Yusef-Zadeh *et al.* 1999, ApJ, 512, 230. OH maser observations.

Coil & Ho 2000, ApJ, 533, 245. NH₃ observations of surroundings.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24''×43'').

Fatuzzo *et al.* 2001, ApJ, 549, 293. Electron-positron lines from the vicinity.

Yusef-Zadeh *et al.* 2001, ApJ, 560, 749. Observations of nearby molecular hydrogen.

Maeda *et al.* 2002, ApJ, 570, 671. Chandra observations.

Sakano *et al.* 2003, AN, 324 (No S1), 197. XMM observations.

Roy & Rao 2004, MNRAS, 349, L25. GMRT at 620 MHz (6'6×11'4).

Sakano *et al.* 2004, MNRAS, 350, 129. XMM observations.

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (1''×2''2).

Park *et al.* 2005, ApJ, 631, 964. Chandra observations.

Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.

Koyama *et al.* 2007, PASJ, 59, S237. Suzaku observations.

Lee *et al.* 2008, ApJ, 674, 247. Molecular H₂ observations of surroundings.

Sjouwerman & Pihlström 2008, ApJ, 681, 1287. VLA at 1.7 GHz of OH masers.

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Tsuboi *et al.* 2012, PASJ, 64, 111. CO observations of SW.

Minh *et al.* 2013, ApJ, 773, 31. NH₃ observations of region.

Zhao *et al.* 2013, ApJ, 777, 146. VLA at 4.8 GHz (0''5×0''7), 5.5 GHz (0''6×1''6), and 8.3 GHz (1''0×2''0).

Nynka *et al.* 2013, ApJ, 778, L31. NuSTAR of compact X-ray source.

G0.3 + 0.0**RA:** 17^h46^m15^s**1-GHz flux/Jy:** 22**Size/arcmin:** 15×8**Dec:** −28°38′**Spectral index:** 0.6**Type:** S

Has been called G0.33+0.04 and G0.4+0.1.

Radio: Bilateral shell, near Galactic Centre.

References:

Kassim & Frail 1996, MNRAS, 283, L51. VLA at 333 MHz (23'' \times 42''), plus review of flux densities and other observations.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (30'').

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').

G0.9 + 0.1

RA: 17^h47^m21^s

Dec: -28°09'

1-GHz flux/Jy: 18?

Spectral index: varies

Size/arcmin: 8

Type: C

Radio: Flat spectrum core within steep spectrum shell.

X-ray: Central core, with non-thermal spectrum.

Point sources: Central pulsar.

References:

Helfand & Becker 1987, ApJ, 314, 203. VLA at 1.4 and 5 GHz, and Einstein observations.

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' \times 91'').

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Mereghetti *et al.* 1998, A&A, 331, L77. X-ray detection.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' \times 24'').

Sidoli *et al.* 2000, A&A, 361, 719. BeppoSAX observations.

Gaensler *et al.* 2001, ApJ, 556, L107. Chandra observations.

Porquet *et al.* 2003, A&A, 401, 197. XMM observations.

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''6 \times 11''3).

Aharonian *et al.* 2005, A&A, 432, L25. γ -ray detection.

Dubner *et al.* 2008, A&A, 487, 1033. ATCA and VLA at 1.4 GHz (1''2 \times 2''5: S=8.3 \pm 0.7 Jy), 5 GHz (1''6 \times 2''5) and 8.3 GHz (0''8 \times 1''5).

Camilo *et al.* 2009, ApJ, 700, L34. Pulsar detection.

Holler *et al.* 2012, A&A, 539, A24. Chandra and XMM observations.

G1.0–0.1

RA: 17^h48^m30^s

Dec: -28°09'

1-GHz flux/Jy: 15

Spectral index: 0.6?

Size/arcmin: 8

Type: S

Has been called G1.05–0.1 and G1.05–0.15.

Radio: Incomplete shell, to the S of Sgr D.

X-ray: Possibly detected.

References:

Downes *et al.* 1979, A&AS, 35, 1. Review of flux densities.

Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (64'' \times 100'': S=12.3 Jy).

Liszt 1992, ApJS, 82, 495. VLA at 1.6 GHz (13'' \times 23'').

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' \times 91'').

Mehring *et al.* 1998, ApJ, 493, 274. VLA at 1.6 GHz (15'' \times 28'') and 5 GHz, including masers observations.

Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' \times 24'').

Sidoli *et al.* 2001, A&A, 372, 651. BeppoSAX possible detection.

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Nobukawa *et al.* 2009, AdSpR, 43, 1045. Suzaku observations.

Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.

G1.4–0.1

RA: 17^h49^m39^s

Dec: –27°46′

1-GHz flux/Jy: ?

Spectral index: ?

Size/arcmin: 10

Type: S

Radio: Shell, brightest in E.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43″×92″: S=2 Jy).

Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.

Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (2′.7×2′.4: S=4.2±0.5).

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8′.2×12′.2).

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G1.9+0.3

RA: 17^h48^m45^s

Dec: –27°10′

1-GHz flux/Jy: 0.6

Spectral index: 0.6

Size/arcmin: 1.5

Type: S

Radio: Shell, brighter to the N, brightening.

X-ray: Shell, with bright limbs to E and W.

References:

Green & Gull 1984, Nature, 312, 527. VLA at 5 GHz (4′.4×2′.0).

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43″×94″).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8′.3×12′.0).

Nord *et al.* 2004, AJ, 128, 1646. VLA at 330 MHz (7″×12″).

Green 2004, BASI, 32, 335. VLA at 1.5 GHz (7′.2×9′.4).

Reynolds *et al.* 2008, ApJ, 680, L41. Chandra observations.

Green *et al.* 2008, MNRAS, 387, L54. VLA at 4.86 GHz (4″×10″), for expansion studies.

Murphy *et al.* 2008, MNRAS, 389, L23. MOST at 843 MHz for flux increase.

Gómez & Rodríguez 2009, RMxAA, 45, 91. VLA at 1.5 GHz (5′.1×10′.6).

Reynolds *et al.* 2009, ApJ, 695, L149. Chandra spectroscopy.

Borkowski *et al.* 2010, ApJ, 724, L161. Chandra observations.

Carlton *et al.* 2011, ApJ, 737, L22. Chandra expansion studies.

Borkowski *et al.* 2013, ApJ, 771, L9. Chandra observations.

G3.7–0.2

RA: 17^h55^m26^s

Dec: –25°50′

1-GHz flux/Jy: 2.3

Spectral index: 0.65

Size/arcmin: 14×11

Type: S

Has been called G003.8–00.3.

Radio: Double arc.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43″×99″: S=2.4 Jy).

Gaensler 1998, ApJ, 493, 781. VLA at 1.4 GHz (9″×15″: S=1.7±0.1 Jy).

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8′.4×11′.4).

G3.8 + 0.3

RA: 17^h52^m55^s
Dec: -25°28'

1-GHz flux/Jy: 3?
Spectral index: 0.6

Size/arcmin: 18
Type: S?

Radio: Incomplete shell.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×100'':*S*=3.5 Jy).
 Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (27''×17'':*S*=6.0±0.4).

G4.2 – 3.5

RA: 18^h08^m55^s
Dec: -27°03'

1-GHz flux/Jy: 3.2?
Spectral index: 0.6?

Size/arcmin: 28
Type: S

Radio: Elongated shell.

Optical: Detected.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G4.5 + 6.8

RA: 17^h30^m42^s
Dec: -21°29'

1-GHz flux/Jy: 19
Spectral index: 0.64

Kepler, SN1604, 3C358

Size/arcmin: 3
Type: S

This is the remnant of Kepler's SN of AD1604.

Radio: Incomplete shell, brighter to the N.

Optical: Faint filaments.

X-ray: Shell, brighter to the N.

Distance: Optical expansion and proper motion indicates about 2.9 kpc, H α observations suggest 3.4 to 6.4 kpc.

References:

van den Bergh & Kamper 1977, ApJ, 218, 617. Optical proper motions.
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
 White & Long 1983, ApJ, 264, 196. Einstein observations.
 Matsui *et al.* 1984, ApJ, 287, 295. VLA at 1.4 (2''5×3''2) and 5 GHz (3''2×4''8) and Einstein image (5'').
 Dickel *et al.* 1988, ApJ, 330, 254. VLA at 1.4 (1''2×2''3) and 5 GHz (0''6×1''0) at two epochs.
 Smith *et al.* 1989, ApJ, 347, 925. EXOSAT observations.
 Hatsukade *et al.* 1990, PASJ, 42, 279. X-ray spectrum.
 Blair *et al.* 1991, ApJ, 366, 484. Optical imaging and spectroscopy.
 Bandiera & van den Bergh 1991, ApJ, 374, 186. Optical changes.
 van den Bergh 1991, PASP, 103, 194. Optical imaging.
 Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Hughes 1999, Apj, 527, 298. ROSAT and Einstein image comparison for expansion studies.
 Reynoso & Goss 1999, AJ, 118, 926. VLA at 1.4 GHz (23'' \times 13'') for H_I studies.
 Kinugasa & Tsunemi 1999, PASJ, 51, 239. ASCA observations.
 Gerardy & Fesen 2001, AJ, 121, 2781. IR spectroscopy and imaging.
 DeLaney *et al.* 2002, Apj, 580, 914. VLA at 1.3 to 1.5 GHz and 5 GHz (7'' \times 2) for spectral index studies.
 Morgan *et al.* 2003, Apj, 597, L33. Sub-mm dust observations.
 Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
 Cassam-Chenai *et al.* 2004, A&A, 414, 545. XMM observations.
 Bamba *et al.* 2005, Apj, 621, 793. Chandra observations of rim.
 Riesgo & López 2005, RMxAA, 41, 57. Optical observations of filament (previously classified as PN, H 2-12).
 Blair *et al.* 2007, Apj, 662, 998. Spitzer observations.
 Reynolds *et al.* 2007, Apj, 668, L135. Chandra observations.
 Sankrit *et al.* 2008, AJ, 135, 538. HST observations.
 Aharonian *et al.* 2008, A&A, 488, 219. γ -ray upper limit.
 Enomoto *et al.* 2008, Apj, 683, 383. γ -ray upper limit.
 Katsuda *et al.* 2008, Apj, 689, 225. Chandra proper motion studies.
 Vink *et al.* 2008, Apj, 689, 231. Chandra proper motion studies.
 Gomez *et al.* 2012, MNRAS, 420, 3557. Herschel IR dust observations.
 Williams *et al.* 2012, Apj, 755, 3. Spitzer spectroscopy.
 Burkey *et al.* 2013, Apj, 764, 63. Chandra observations.
 Yang *et al.* 2013, Apj, 766, 44. Suzaku spectroscopy.
 Park *et al.* 2013, Apj, 767, L10. Suzaku observations.

G4.8 + 6.2

RA: 17^h33^m25^s

Dec: -21°34'

1-GHz flux/Jy: 3

Spectral index: 0.6

Size/arcmin: 18

Type: S

Has been called G4.5+6.2.

Radio: Faint shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10'4).
 Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (2'.2 \times 1'.3 : S = 5.5 \pm 1.2 Jy), and NVSS at 1.4 GHz.
 Hewitt & Yusef-Zadeh 2009, Apj, 694, L16. OH maser search.

G5.2 - 2.6

RA: 18^h07^m30^s

Dec: -25°45'

1-GHz flux/Jy: 2.6?

Spectral index: 0.6?

Size/arcmin: 18

Type: S

Radio: Poorly resolved shell.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, Apj, 694, L16. OH maser search.

G5.4–1.2

Milne 56

RA: 18^h02^m10^s**1-GHz flux/Jy:** 35?**Size/arcmin:** 35**Dec:** –24°54′**Spectral index:** 0.2?**Type:** C?

Part been called G5.3–1.0. Has been suggested that this is not a SNR.

Radio: Incomplete shell, including wide ‘v’ of emission to east with small flat-spectrum source at apex.

Optical: Detected.

X-ray: Pulsar detected, with faint extension.

Point sources: Pulsar nearby, in flat spectrum source.

Distance: H_i absorption suggests > 4.3 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′: $S=38$ Jy).

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′).

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′4) and 5 GHz (4′4).

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′8: $S=21.9\pm 2.4$ Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).

Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.

Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 and 5 GHz.

Helfand & Becker 1985, Nature, 313, 118. Suggesting it is not a SNR.

Manchester *et al.* 1985, MNRAS, 212, 975. Pulsar detection.

Caswell *et al.* 1987, MNRAS, 225, 329. MOST at 843 MHz (42″×110″).

Frail & Kulkarni 1991, Nature, 352, 785. Pulsar and remnant association.

Manchester *et al.* 1991, MNRAS, 253, 7P. Pulsar and remnant association.

Milne *et al.* 1992, MNRAS, 255, 707. Parkes 64-m at 4.75 (4′5: $S=30.8\pm 2.1$ Jy) and 8.4 GHz (3′: $S=24\pm 3$ Jy), including polarisation.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′0×4′9: $S=38$ Jy).

Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz (68″×73″), plus H_i absorption.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Gaensler & Frail 2000, Nature, 406, 158. Pulsar observations, including proper motion.

Kaspi *et al.* 2001, ApJ, 562, L163. X-ray detection of pulsar, and upper limit for remnant.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.

Blazek *et al.* 2006, ApJ, 652, 1523. Proper motion study of pulsar.

Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion study of pulsar.

Liszt *et al.* 2009, A&A, 508, 1331. CO and IR observations of region.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.

G5.5+0.3**RA:** 17^h57^m04^s**1-GHz flux/Jy:** 5.5**Size/arcmin:** 15×12**Dec:** –24°00′**Spectral index:** 0.7**Type:** S

Has been called G5.55+0.32.

Radio: Shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=14.3\pm 0.3$ Jy), plus other observations.

Liszt *et al.* 2009, A&A, 508, 1331. CO and IR observations of region.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G5.9 + 3.1

RA: 17^h47^m20^s
Dec: -22°16'

1-GHz flux/Jy: 3.3?
Spectral index: 0.4?

Size/arcmin: 20
Type: S

Radio: Asymmetric shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Effelsberg 100-m at 2.7 GHz (4'3).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.1 + 0.5

RA: 17^h57^m29^s
Dec: -23°25'

1-GHz flux/Jy: 4.5
Spectral index: 0.9

Size/arcmin: 18×12
Type: S

Has been called G6.10+0.53.

Radio: Partial shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S=13.4±0.2 Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G6.1 + 1.2

RA: 17^h54^m55^s
Dec: -23°05'

1-GHz flux/Jy: 4.0?
Spectral index: 0.3?

Size/arcmin: 30×26
Type: F

Has been called G6.1+1.15.

Radio: Faint, diffuse emission.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.4–0.1

W28

RA: 18^h00^m30^s**1-GHz flux/Jy:** 310**Size/arcmin:** 48**Dec:** –23°26′**Spectral index:** varies**Type:** C

Has been called G6.6–0.2.

Radio: Several non-thermal sources in a ring, with flat spectrum core.

Optical: Diffuse emission.

X-ray: Diffuse emission from most of the remnant.

Point sources: Young pulsar near edge of remnant, but not thought to be related.

Distance: H_i observations suggest 1.9 kpc.

References:

- Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10 GHz (3′).
 van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′4) and 5 GHz (4′4).
 Dopita *et al.* 1977, ApJ, 214, 179. Some optical line ratios.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).
 Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
 Andrews *et al.* 1983, ApJ, 266, 684. VLA at 1.4 GHz (8′′) 4.9 GHz (3′′) and 15 GHz (1′′) of central region only, plus Einstein image of central region.
 Bohigas *et al.* 1983, RMxAA, 8, 155. Optical spectra.
 Odenwald *et al.* 1984, ApJ, 279, 162. Nearby IR source.
 Andrews *et al.* 1985, AJ, 90, 310. VLA of central component.
 Long *et al.* 1991, ApJ, 373, 567. Einstein and optical observations.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.1×4′.4: S=660 Jy).
 Kaspi *et al.* 1993, ApJ, 409, L57. Pulsar association.
 Frail *et al.* 1993, Nature, 365, 136. VLA at 327 MHz (smoothed to 65′′), plus pulsar association.
 Frail *et al.* 1994, ApJ, 424, L111. VLA of associated OH masers.
 Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ -ray emission.
 Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Claussen *et al.* 1997, ApJ, 489, 143. VLA of OH masers.
 Claussen *et al.* 1999, ApJ, 522, 349. High resolution observations of OH masers.
 Arikawa *et al.* 1999, PASJ, 51, L7. Observations of shocked CO.
 Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz (2′.6×5′.5) and 1.48 GHz (40′′×65′′).
 Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz (97′′×52′′: S = 425±40 Jy) and 1415 MHz (88′′×48′′: S = 246±20 Jy), and comparison with other observations.
 Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
 Rowell *et al.* 2000, A&A, 359, 337. Upper limit on high energy γ -rays.
 Douvion *et al.* 2001, A&A, 373, 281. ISO observations.
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
 Velázquez *et al.* 2002, AJ, 124, 2145. Parkes 64-m at 1.4 GHz (15′) for H_i.
 Rho & Borkowski 2002, ApJ, 575, 201. ROSAT and ASCA observations.
 Claussen *et al.* 2002, ApJ, 580, 909. Observations of nearby source.
 Yusef-Zadeh *et al.* 2003, ApJ, 583, 267. OH observations.
 Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz of associated OH masers.
 Mavromatakis *et al.* 2004, A&A, 426, 567. Optical observations.
 Reach *et al.* 2005, ApJ, 618, 297. Molecular lines and near IR observations.
 Hoffman *et al.* 2005, ApJ, 620, 257. OH maser observations.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.

Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
 Aharonian *et al.* 2008, A&A, 481, 401. γ -ray observations.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Hewitt *et al.* 2008, ApJ, 683, 189.GBT at 1.6 and 1.7 GHz for OH masers.
 Guilani *et al.* 2010, A&A, 516, L11. γ -ray observations.
 Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.
 Abdo *et al.* 2010, ApJ, 718, 348. Fermi observations.
 Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
 Sawada *et al.* 2012, PASJ, 64, 81. Suzaku observations.
 Nichols *et al.* 2012, MNRAS, 419, 251. CO observations of selected regions.
 GUSDORF *et al.* 2012, A&A, 542, L19. CO observations of regions in NE.

G6.4 + 4.0

RA: 17^h45^m10^s
Dec: -21°22'

1-GHz flux/Jy: 1.3?
Spectral index: 0.4?

Size/arcmin: 31
Type: S

Radio: Faint asymmetric shell.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.5 - 0.4

RA: 18^h02^m11^s
Dec: -23°34'

1-GHz flux/Jy: 27
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G6.51-0.48, and part has been called G6.67-0.42.

Radio: Shell, overlapping G6.4-0.1.

Optical: Detected.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 330 MHz (2'6x5'5) and 1.4 GHz (0'7x1'1).
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': S=60.8±0.4 Jy), plus other observations.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G7.0 - 0.1

RA: 18^h01^m50^s
Dec: -22°54'

1-GHz flux/Jy: 2.5?
Spectral index: 0.5?

Size/arcmin: 15
Type: S

Has been called G7.06-0.12.

Radio: Double rim, brightest in W, confused by bright H Π region M20 in SE.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz (2'6x5'5) and 1.48 GHz (40''x65'').
 Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz (97''x52'') and 1415 MHz (88''x48'').

G7.2 + 0.2**RA:** 18^h01^m07^s**Dec:** -22°38'**1-GHz flux/Jy:** 2.8**Spectral index:** 0.6**Size/arcmin:** 12**Type:** S

Has been called G7.20+0.20.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 5.2 ± 0.2 Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G7.7 – 3.7**RA:** 18^h17^m25^s**Dec:** -24°04'**1-GHz flux/Jy:** 11**Spectral index:** 0.32**1814–24****Size/arcmin:** 22**Type:** S

Radio: Shell, with high polarisation.

References:

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).

Milne *et al.* 1986, MNRAS, 223, 487. MOST at 843 MHz (44'' × 108'') and Parkes 64-m at 8.4 GHz (3' : S = 4.6 ± 0.5 Jy), with polarisation, plus review of flux densities.

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (70'' × 35'' : S = 9.9 ± 0.1 Jy), including polarisation.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G8.3 – 0.0**RA:** 18^h04^m34^s**Dec:** -21°49'**1-GHz flux/Jy:** 1.2**Spectral index:** 0.6**Size/arcmin:** 5 × 4**Type:** S

Has been called G8.31–0.09.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 2.3 ± 0.1 Jy), plus other observations.

Higashi *et al.* 2008, ApJ, 683, 957. γ -ray detection.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G8.7 – 5.0**RA:** 18^h24^m10^s**Dec:** -23°48'**1-GHz flux/Jy:** 4.4**Spectral index:** 0.3**Size/arcmin:** 26**Type:** S

Radio: Asymmetric shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.

Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G8.7–0.1

(W30)

RA: 18^h05^m30^s**1-GHz flux/Jy:** 80**Size/arcmin:** 45**Dec:** –21°26′**Spectral index:** 0.5**Type:** S?

Has been called G8.6–0.1.

Radio: Clumpy non-thermal shell, with low-frequency turnover.

X-ray: Northern edge detected.

Point sources: Pulsar inside western edge.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7′.2×9′.7 : S = 190±50 Jy).

Kassim & Weiler 1990, Nature, 343, 146. VLA at 327 MHz (3′.0×3′.7).

Kassim & Weiler 1990, ApJ, 360, 184. VLA at 327 MHz (2′.8×4′.1 : S = 129±11 Jy), and part at 1.4 GHz (0′.9×1′.8), plus review of flux densities.

Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz (37″×55″).

Finley & Ögelman 1994, ApJ, 434, L25. ROSAT observations, including pulsar.

Aharonian *et al.* 2005, Science, 307, 1938. γ -ray detection.

Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations.

Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.

Briskin *et al.* 2006, ApJ, 652, 554. Pulsar proper motion.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.

Castro & Slane 2010, ApJ, 717, 372. Fermi observations.

Ajello *et al.* 2012, ApJ, 744, 80. Fermi observations.

G8.9+0.4**RA:** 18^h03^m58^s**1-GHz flux/Jy:** 9**Size/arcmin:** 24**Dec:** –21°03′**Spectral index:** 0.6**Type:** S

Has been called G8.90+0.40.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″ : S = 18.2±0.5 Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G9.7–0.0**RA:** 18^h07^m22^s**1-GHz flux/Jy:** 3.7**Size/arcmin:** 15×11**Dec:** –20°35′**Spectral index:** 0.6**Type:** S

Has been called G9.7–0.1 and G9.70–0.06.

Radio: Shell.

References:

Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″ : S = 6.5±0.2 Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.

G9.8 + 0.6**RA:** 18^h05^m08^s**Dec:** -20°14'**1-GHz flux/Jy:** 3.9**Spectral index:** 0.5**Size/arcmin:** 12**Type:** S**Radio:** Asymmetric shell.**References:**

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Caswell 1983, MNRAS, 204, 833. Molonglo at 408 MHz (3': S=5.8±0.6 Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41''×63'': S=3.5±0.4 Jy).
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G9.9 – 0.8**RA:** 18^h10^m41^s**Dec:** -20°43'**1-GHz flux/Jy:** 6.7**Spectral index:** 0.4**Size/arcmin:** 12**Type:** S

Has been called G9.95–0.81.

Radio: Shell.**Optical:** Detected.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': S=11.0±0.3 Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G10.5 – 0.0**RA:** 18^h09^m08^s**Dec:** -19°47'**1-GHz flux/Jy:** 0.9**Spectral index:** 0.6**Size/arcmin:** 6**Type:** S

Has been called G10.59–0.04.

Radio: Partial shell.**X-ray:** Detected.**References:**

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': S=1.4±0.1 Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G11.0–0.0**RA:** 18^h10^m04^s**Dec:** –19°25′**1-GHz flux/Jy:** 1.3**Spectral index:** 0.6**Size/arcmin:** 11×9**Type:** S

Has been called G11.0+0.0 and G11.03–0.05.

Radio: Partial shell.

X-ray: Diffuse emission.

References:

Bamba *et al.* 2003, ApJ, 589, 253. ASCA observations.

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 3.1 \pm 0.2$ Jy), plus other observations.

G11.1–1.0**RA:** 18^h14^m03^s**Dec:** –19°46′**1-GHz flux/Jy:** 5.8**Spectral index:** 0.5**Size/arcmin:** 18×12**Type:** S

Has been called G11.2–1.1 and G11.17–1.04.

Radio: Shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 11.0 \pm 0.3$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5'': $S = 3.40 \pm 0.25$ Jy) including polarisation and review of flux densities.

G11.1–0.7**RA:** 18^h12^m46^s**Dec:** –19°38′**1-GHz flux/Jy:** 1.0**Spectral index:** 0.7**Size/arcmin:** 11×7**Type:** S

Has been called G11.15–0.71.

Radio: Partial shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 2.3 \pm 0.1$ Jy), plus other observations.

G11.1+0.1**RA:** 18^h09^m47^s**1-GHz flux/Jy:** 2.3**Size/arcmin:** 12×10**Dec:** −19°12′**Spectral index:** 0.4**Type:** S

Has been called G11.18+0.11.

Radio: Shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=3.5\pm0.2$ Jy), plus other observations.

G11.2−0.3**RA:** 18^h11^m27^s**1-GHz flux/Jy:** 22**Size/arcmin:** 4**Dec:** −19°25′**Spectral index:** 0.5**Type:** C

Probably associated with the SN of AD386.

Radio: Symmetrical clumpy shell, with flatter spectrum core.

X-ray: Shell, with hard spectrum centrally brightened region around pulsar.

Point sources: Central pulsar.

Distance: H_i absorption indicates 4.4 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Radhakrishnan *et al.* 1972, ApJS, 24, 49. H_i absorption.

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20''×25'') and Einstein observations, with review of flux densities.

Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz, plus H_i absorption, Einstein observations.

Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26'': $S=4.04\pm0.24$ Jy).

Green *et al.* 1988, MNRAS, 231, 735. VLA at 1.4 and 5 GHz.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'2×4'1': $S=39$ Jy).

Reynolds *et al.* 1994, MNRAS, 271, L1. ROSAT image and spectra.

Vasisht *et al.* 1996, ApJ, 456, L59. ASCA observations.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Torii *et al.* 1997, ApJ, 489, L145. ASCA detection of pulsar.

Torii *et al.* 1999, ApJ, 523, L69. X-ray timing observations of pulsar.

Kothes & Reich 2001, A&A, 372, 627. Effelsberg 100-m at 4.25, 10.45, 14.9 and 32 GHz (2'5, 1'1, 0'86 and 0'45: $S=9.6\pm0.5, 6.3\pm0.4, 5.7\pm0.4$ and 3.8 ± 0.4).

Kaspi *et al.* 2001, ApJ, 560, 371. Chandra observations.

Tam *et al.* 2002, ApJ, 572, 202. VLA at 1.4/1.5 GHz (2''6×1''8: $S=16.6\pm0.9$ Jy) and 5 GHz (2''1×1''5: $S=8.4\pm0.9$ Jy) for spectral studies.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 14.7 GHz.

Tam & Roberts 2003, ApJ, 598, L27. Multi-epoch VLA observations at 1.4/1.5 GHz and 5 GHz, for expansion studies.

Roberts *et al.* 2003, ApJ, 588, 992. Chandra observations.

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (18'').

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.

Koo *et al.* 2007, ApJ, 657, 308. IR observations.

Dean *et al.* 2008, MNRAS, 384, L29. Integral observations of pulsar and nebula.
 Moon *et al.* 2009, ApJ, 703, L81. IR spectroscopy.
 Mizuno *et al.* 2010, AJ, 139, 1542. Spitzer observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5'' : S = 9.0 \pm 0.5$ Jy) including polarisation and review of flux densities.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Lee *et al.* 2013, ApJ, 770, 143. IR observations.

G11.4–0.1

RA: $18^{\text{h}}10^{\text{m}}47^{\text{s}}$
Dec: $-19^{\circ}05'$

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 8
Type: S?

Radio: Incomplete shell, possibly with central core.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3' : S = 9.4$ Jy) contaminated by sidelobes of a nearby source, and Parkes 64-m at 5 GHz ($4' : S = 2.8$ Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'8'' : S = 2.0 \pm 0.4$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'6''$).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3'2'' \times 4'1'' : S = 18$ Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($44'' \times 63'' : S = 5.1 \pm 0.6$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz ($25''$), 1.5 GHz ($25''$), and 74 MHz.

G11.8–0.2

RA: $18^{\text{h}}12^{\text{m}}25^{\text{s}}$
Dec: $-18^{\circ}44'$

1-GHz flux/Jy: 0.7
Spectral index: 0.3

Size/arcmin: 4
Type: S

Has been called G11.89–0.21.

Radio: Shell.

X-ray: Detected.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.9 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.0–0.1

RA: $18^{\text{h}}12^{\text{m}}11^{\text{s}}$
Dec: $-18^{\circ}37'$

1-GHz flux/Jy: 3.5
Spectral index: 0.7

Size/arcmin: 7?
Type: ?

Radio: Incomplete shell, defined in E only.

X-ray: Detected.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3' : S = 6.6$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 1.1$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'6''$).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3'2'' \times 4'1''$).

Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($41'' \times 61''$; $S=0.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

G12.2 + 0.3

RA: $18^{\text{h}}11^{\text{m}}17^{\text{s}}$
Dec: $-18^{\circ}10'$

1-GHz flux/Jy: 0.8
Spectral index: 0.7

Size/arcmin: 6×5
Type: S

Has been called G12.26+0.30.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42''$; $S=1.5 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.5 + 0.2

RA: $18^{\text{h}}12^{\text{m}}14^{\text{s}}$
Dec: $-17^{\circ}55'$

1-GHz flux/Jy: 0.6
Spectral index: 0.4

Size/arcmin: 6×5
Type: C?

Has been called G12.58+0.22.

Radio: Diffuse, central brightened.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42''$; $S=0.8 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.7 – 0.0

RA: $18^{\text{h}}13^{\text{m}}19^{\text{s}}$
Dec: $-17^{\circ}54'$

1-GHz flux/Jy: 0.8
Spectral index: 0.8

Size/arcmin: 6
Type: S

Has been called G12.72–0.00.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42''$; $S=2.0 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.8–0.0**RA:** 18^h13^m37^s**Dec:** –17°49′**1-GHz flux/Jy:** 0.8**Spectral index:** 0.5**Size/arcmin:** 3**Type:** C?

Has been called G12.82–0.02 and G12.83–0.02.

Radio: Shell.**X-ray:** Diffuse.**Point sources:** Central X-ray pulsar.**References:**

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2005, ApJ, 629, L105. VLA at 330 MHz (19''×32''), plus other observations.
 Ubertini *et al.* 2005, ApJ, 629, L109. X-ray, γ -ray and other observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 1.2 \pm 0.1$ Jy), plus other observations.
 Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.
 Albert *et al.* 2006, ApJ, 637, L41. γ -ray observations.
 Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
 Funk *et al.* 2007, A&A, 470, 249. XMM observations, CO observations of surroundings.
 Helfand *et al.* 2007, ApJ, 665, 1297. Chandra observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Gotthelf & Halpern 2009, ApJ, 700, L158. Pulsar detection.
 Halpern *et al.* 2012, ApJ, 753, L14. Pulsar observations.

G13.3–1.3**RA:** 18^h19^m20^s**Dec:** –18°00′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 70×40**Type:** S?**Radio:** Amorphous emission.**Optical:** Filaments in S.**X-ray:** Elongated emission.**Distance:** Absorption indicates 2--4 kpc.**References:**

Seward *et al.* 1995, ApJ, 449, 681. ROSAT detection, optical studies and observations of CO.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G13.5+0.2**RA:** 18^h14^m14^s**Dec:** –17°12′**1-GHz flux/Jy:** 3.5?**Spectral index:** 1.0?**Size/arcmin:** 5×4**Type:** S

Has been called G13.46+0.16.

Radio: Elongated, incomplete shell.**References:**

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ($S = 0.65 \pm 0.05$ Jy) and 1.4 GHz (15'' : $S = 2.67 \pm 0.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G14.1–0.1**RA:** 18^h16^m40^s**Dec:** –16°41′**1-GHz flux/Jy:** 0.5**Spectral index:** 0.6**Size/arcmin:** 6×5**Type:** S

Has been called G14.18–0.12.

Radio: Shell.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S = 0.9 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G14.3+0.1**RA:** 18^h15^m58^s**Dec:** –16°27′**1-GHz flux/Jy:** 0.6**Spectral index:** 0.4**Size/arcmin:** 5×4**Type:** S

Has been called G14.30+0.14. Has been suggested this is not an SNR.

Radio: Partial shell.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S = 1.2 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Pinheiro Gonçalves *et al.* 2011, AJ, 142, 47. IR observations.

G15.1–1.6**RA:** 18^h24^m00^s**Dec:** –16°34′**1-GHz flux/Jy:** 5.5?**Spectral index:** 0.0?**Size/arcmin:** 30×24**Type:** S?**Radio:** Elongated, incomplete shell.**Optical:** Diffuse shell.**References:**

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).
 Boumis *et al.* 2008, A&A, 481, 705. Optical detection.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5: $S = 4.8 \pm 0.3$ Jy) including polarisation and review of flux densities.

G15.4+0.1**RA:** 18^h18^m02^s**Dec:** −15°27′**1-GHz flux/Jy:** 5.6**Spectral index:** 0.62**Size/arcmin:** 15×14**Type:** S

Has been called G15.42+0.18.

Radio: Shell.

Distance: H_I observations suggest 4.8 kpc.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=10.9\pm 0.3$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5: $S=2.1\pm 0.2$ Jy) including polarisation and review of flux densities.

Castelletti *et al.* 2013, A&A, 557, L15. GMRT at 1.4 GHz (15″) plus SGPS H_I and CO observations of region.

G15.9+0.2**RA:** 18^h18^m52^s**Dec:** −15°02′**1-GHz flux/Jy:** 5.0**Spectral index:** 0.63**Size/arcmin:** 7×5**Type:** S?

Radio: Incomplete shell, with bright concentration to the E.

X-ray: Shell, brighter to S and E.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=7.7$ Jy) and Parkes 64-m at 5 GHz (4′: $S=1.9$ Jy).

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′8: $S=1.1\pm 0.2$ Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).

Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz (58″×44″).

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (77″×61″: $S=11.2\pm 1.0$ Jy), 1.4 GHz (23″×14″: $S=3.9\pm 0.1$ Jy) and 4.9 GHz (13″×16″).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Reynolds *et al.* 2006, ApJ, 652, L45. Chandra observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5: $S=2.0\pm 0.3$ Jy) including polarisation and review of flux densities.

G16.0−0.5**RA:** 18^h21^m56^s**Dec:** −15°14′**1-GHz flux/Jy:** 2.7**Spectral index:** 0.6**Size/arcmin:** 15×10**Type:** S

Has been called G16.05−0.57.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=4.9\pm 0.2$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Beaumont *et al.* 2011, ApJ, 741, 14. CO of region.

G16.2–2.7**RA:** 18^h29^m40^s**Dec:** –16°08′**1-GHz flux/Jy:** 2.5**Spectral index:** 0.4**Size/arcmin:** 17**Type:** S**Radio:** Double rim.**References:**

Trushkin 1999, A&A, 352, L103. Review of radio observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 1.28±0.10 Jy) including polarisation and review of flux densities.**G16.4–0.5****RA:** 18^h22^m38^s**Dec:** –14°55′**1-GHz flux/Jy:** 4.6**Spectral index:** 0.3?**Size/arcmin:** 13**Type:** S

Has been called G16.41–0.55.

Radio: Partial shell.**References:**Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″ : S = 10.0±0.3 Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 3.0±0.3 Jy) including polarisation and review of flux densities.**G16.7 + 0.1****RA:** 18^h20^m56^s**Dec:** –14°20′**1-GHz flux/Jy:** 3.0**Spectral index:** 0.6**Size/arcmin:** 4**Type:** C

Has been called G16.73+0.08.

Radio: Asymmetric shell with flat-spectrum core.**X-ray:** Non-thermal core.**References:**Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz (5 : S = 0.95 Jy) and 1.4 GHz (15″ : S = 2.43 Jy) and Ooty at 327 MHz (S = 5.13 Jy).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.

Reynoso & Mangum 2000, ApJ, 545, 874. CO observations of surroundings.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.Helfand *et al.* 2003, ApJ, 592, 941. XMM observations.

Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (19″ × 25″).

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 6 GHz.Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 1.23±0.11 Jy) including polarisation and review of flux densities.

G17.0–0.0**RA:** 18^h21^m57^s**Dec:** –14°08′**1-GHz flux/Jy:** 0.5**Spectral index:** 0.5**Size/arcmin:** 5**Type:** S

Has been called G17.02–0.04.

Radio: Shell.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=0.7\pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G17.4–2.3**RA:** 18^h30^m55^s**Dec:** –14°52′**1-GHz flux/Jy:** 5**Spectral index:** 0.5?**Size/arcmin:** 24?**Type:** S**Radio:** Incomplete, poorly defined shell.**Optical:** Filaments to SE, and diffuse emission.**References:**

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2002, A&A, 385, 1042. Optical observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5: $S=2.3\pm 0.2$ Jy) including polarisation and review of flux densities.

G17.4–0.1**RA:** 18^h23^m08^s**Dec:** –13°46′**1-GHz flux/Jy:** 0.4**Spectral index:** 0.7**Size/arcmin:** 6**Type:** S

Has been called G17.48–0.12.

Radio: Partial shell.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=0.9\pm 0.1$ Jy), plus other observations.

G17.8–2.6**RA:** 18^h32^m50^s**Dec:** –14°39′**1-GHz flux/Jy:** 5**Spectral index:** 0.5**Size/arcmin:** 24**Type:** S**Radio:** Well defined shell.**References:**

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9^{\circ}5':S=2.23\pm 0.13$ Jy) including polarisation and review of flux densities.

G18.1–0.1

RA: $18^{\text{h}}24^{\text{m}}34^{\text{s}}$
Dec: $-13^{\circ}11'$

1-GHz flux/Jy: 4.6
Spectral index: 0.5

Size/arcmin: 8
Type: S

Has been called G18.1–0.2 and G18.16–0.16.

Radio: Shell.

X-ray: Detected.

Distance: Association with other sources suggests 4 kpc.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz ($8'$).

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S=7.6\pm 0.1$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Paron *et al.* 2013, MNRAS, 433, 1619. CO, optical and other observations of region.

G18.6–0.2

RA: $18^{\text{h}}25^{\text{m}}55^{\text{s}}$
Dec: $-12^{\circ}50'$

1-GHz flux/Jy: 1.4
Spectral index: 0.4

Size/arcmin: 6
Type: S

Has been called G18.62–0.28.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S=1.9\pm 0.1$ Jy), plus other observations.

G18.8+0.3

Kes 67

RA: $18^{\text{h}}23^{\text{m}}58^{\text{s}}$
Dec: $-12^{\circ}23'$

1-GHz flux/Jy: 33
Spectral index: 0.46

Size/arcmin: 17×11
Type: S

Has been called G18.9+0.3.

Radio: Incomplete shell, in complex region near the H_{II} region W39.

Distance: Association with molecular cloud and H_I absorption suggests 12 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S=17\pm 7$ Jy).

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz ($3' : S=38$ Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'4$) and 5 GHz ($4'4$).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'6$).

Milne *et al.* 1989, PASA, 8, 187. Parkes 64-m at 8.4 GHz ($3'0 : S=12.9\pm 1.0$ Jy), including polarisation.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($2'.9 \times 3'.5 : S = 55$ Jy).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ($75'' \times 55'' : S = 29.9 \pm 0.3$ Jy).
 Dubner *et al.* 1999, AJ, 118, 930. Parkes 64-m at 1.6 GHz ($15''$) for H α , VLA at 1.6 GHz ($17'' \times 12''$) for OH, plus CO observations.
 Dubner *et al.* 2004, A&A, 426, 201. CO observations of environment.
 Tian *et al.* 2007, A&A, 474, 541. VGPS at 1.4 GHz ($1'$) including H α , plus CO observations of region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 15.3 \pm 0.9$ Jy) including polarisation and review of flux densities.
 Vasquez *et al.* 2012, A&A, 545, A89. CO observations of region.
 Paron *et al.* 2012, A&A, 547, A60. CO and other molecular observations of region.

G18.9–1.1

RA: $18^{\text{h}}29^{\text{m}}50^{\text{s}}$

Dec: $-12^{\circ}58'$

1-GHz flux/Jy: 37

Spectral index: 0.39

Size/arcmin: 33

Type: C?

Has been called G18.95–1.1 and G18.94–1.04.

Radio: Non-thermal, diffuse partially limb-brightened, with central ridge.

Optical: Detected.

X-ray: Partial shell.

Point sources: Compact X-ray source, with diffuse nebula.

Distance: Various observations suggest 2 kpc.

References:

Fürst *et al.* 1985, Nature, 314, 720. Effelsberg 100-m at 4.75 GHz ($2'.4 : S = 23.8$ Jy), plus other flux densities.
 Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz ($7'.2 \times 8' : S = 82 \pm 15$ Jy), plus review of flux densities.
 Barnes & Turtle 1988, IAUCom, 101, 347. Molonglo at 408 MHz ($2'.9 \times 3'.1 : S = 58 \pm 9$ Jy) and Parkes 64-m at 5 GHz ($4'.4 \times 4'.1 : S = 23 \pm 6$ Jy).
 Patnaik *et al.* 1988, Nature, 332, 136. Ooty at 327 MHz ($0'.6 \times 1'.6$).
 Fürst *et al.* 1989, A&A, 209, 361. Effelsberg 100-m at 4.75 GHz ($2'.45 : S = 23.8$ Jy) and VLA at 1.5 and 4.9 GHz ($19'' \times 14''$), and Effelsberg 100-m at 1.4 GHz ($9'$) for H α .
 Aschenbach *et al.* 1991, A&A, 246, L32. ROSAT observations.
 Fürst *et al.* 1997, A&A, 319, 655. ROSAT observations, and Effelsberg 100-m at 10.6 GHz ($1'.1$).
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.
 Harrus *et al.* 2004, ApJ, 603, 152. ROSAT and ASCA observations.
 Tüllmann *et al.* 2010, ApJ, 720, 848. Chandra detection of compact source.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 19.6 \pm 1.0$ Jy) including polarisation and review of flux densities.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G19.1+0.2

RA: $18^{\text{h}}24^{\text{m}}56^{\text{s}}$

Dec: $-12^{\circ}07'$

1-GHz flux/Jy: 10

Spectral index: 0.5

Size/arcmin: 27

Type: S

Has been called G19.15+0.27.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 17.4 \pm 0.4$ Jy), plus other observations.

G20.0–0.2**RA:** 18^h28^m07^s**Dec:** –11°35′**1-GHz flux/Jy:** 10**Spectral index:** 0.1**Size/arcmin:** 10**Type:** F**Radio:** Faint, filled-centre, polarised.**X-ray:** Centrally brightened.**Point sources:** OH source 20.1–0.1 is nearby.**References:**

Becker & Helfand 1985, ApJ, 297, L25. VLA at 1.4 and 5 GHz (12'').

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7'2×8':S=8.5±2 Jy), plus review of flux densities.

Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'.3), including polarisation.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5:S=9.2±0.5 Jy) including polarisation and review of flux densities.Petriella *et al.* 2013, A&A, 554, A73. Chandra observations, plus CO and H_i observations of region.

G20.4+0.1**RA:** 18^h27^m51^s**Dec:** –11°00′**1-GHz flux/Jy:** 9?**Spectral index:** 0.1?**Size/arcmin:** 8**Type:** S?

Has been called G20.47+0.16.

Radio: Shell.**References:**Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'':S=4.2±0.1 Jy), plus other observations.Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5:S=7.5±0.5 Jy) including polarisation and review of flux densities.

G21.0–0.4**RA:** 18^h31^m12^s**Dec:** –10°47′**1-GHz flux/Jy:** 1.1**Spectral index:** 0.6**Size/arcmin:** 9×7**Type:** S

Has been called G21.04–0.47.

Radio: Shell.**References:**Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'':S=2.3±0.2 Jy), plus other observations.

G21.5–0.9**RA:** 18^h33^m33^s**Dec:** –10°35′**1-GHz flux/Jy:** 7**Spectral index:** varies**Size/arcmin:** 5**Type:** C

Early observations relate to the central core only.

Radio: Filled-centre, with high frequency turnover.

X-ray: Central core, with extended, faint halo.

Point sources: Central pulsar.

Distance: H_i absorption indicates 4.6 kpc.

References:

Wilson & Weiler 1976, A&A, 53, 89. WSRT at 5 GHz (6''×35'').

Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20''×5'') and 8 GHz (7''×2''), plus review of flux densities.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Becker & Szymkowiak 1981, ApJ, 248, L23. VLA at 5 GHz (8''), and Einstein observations.

Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT X-ray spectrum, and VLA H_i observations.

Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (26''5 : S = 5.64±0.29 Jy).

Fürst *et al.* 1988, PASJ, 40, 347. NRO array at 22.3 GHz (4''4×7''3).

Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.

Salter *et al.* 1989, A&A, 225, 167. Observations at 90.7 (29''5 : S = 3.8±0.4 Jy) and 141.9 GHz (S = 2.5±1.2 Jy).

Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz (S = 3.94±0.70 Jy), plus review of flux densities.

Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'3×3'4 : S = 9 Jy).

Wallace *et al.* 1994, A&A, 286, 565. H_i of surroundings.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Slane *et al.* 2000, ApJ, 533, L29. Chandra observations identifying X-ray halo.

Warwick *et al.* 2001, A&A, 365, L248. XMM observations of X-ray halo.

Bock *et al.* 2001, ApJ, 561, L203. BIMA at 94 GHz (8''6×4''6).

Safi-Harb *et al.* 2001, ApJ, 561, 308. Chandra and other X-ray observations.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 22 GHz (8'') and 32 GHz, including polarisation.

La Palombara & Mereghetti 2002, A&A, 383, 916. XMM upper limit on pulsations.

Bocchino *et al.* 2005, A&A, 442, 539. XMM and Chandra observations.

Gupta *et al.* 2005, Current Science, 89, 853. Pulsar discovery.

Camilo *et al.* 2006, ApJ, 637, 456. Pulsar discovery.

Bietenholz & Bartel 2008, MNRAS, 386, 1411. VLA at 4.75 GHz (0''53×0''82) for expansion studies.

Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1') including H_i.

Matheson & Safi-Harb 2010, ApJ, 724, 572. Chandra observations.

Bietenholz *et al.* 2011, MNRAS, 412, 1221. VLA at 1.4 GHz (14''×18'').

Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 6 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 6.5±0.4 Jy) including polarisation and review of flux densities.

Zajczyk *et al.* 2012, A&A, 542, A12. IR observations, including polarisation.

G21.5–0.1**RA:** 18^h30^m50^s**Dec:** –10°09′**1-GHz flux/Jy:** 0.4**Spectral index:** 0.5**Size/arcmin:** 5**Type:** S

Has been called G21.56–0.10.

Radio: Partial shell.

X-ray: Detected.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=0.5\pm 0.1$ Jy), plus other observations.

G21.6–0.8**RA:** 18^h33^m40^s**Dec:** –10°25′**1-GHz flux/Jy:** 1.4**Spectral index:** 0.5?**Size/arcmin:** 13**Type:** S

Has been called G21.64–0.84.

Radio: Faint, irregular shell.

References:

Bietenholz *et al.* 2011, MNRAS, 412, 1221. VLA at 327 MHz (85″: $S=2.8$ Jy) and 1.4 GHz (14″×18″).

G21.8–0.6

Kes 69

RA: 18^h32^m45^s**Dec:** –10°08′**1-GHz flux/Jy:** 65**Spectral index:** 0.56**Size/arcmin:** 20**Type:** S

Radio: Incomplete shell.

X-ray: Detected.

Distance: Association with CO indicates 5.2 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).

Wilson 1972, A&A, 19, 354. H₂CO absorption.

Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 5 GHz (6′) and 10 GHz (3′).

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′: $S=42.3\pm 4.6$ Jy).

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).

Seward 1990, ApJS, 73, 781. Einstein observations.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.3×3′.5: $S=132$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.

Yusef-Zadeh *et al.* 2003, ApJ, 585, 319. X-ray observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Wood *et al.* 2008, AJ, 135, 2358. VLA at 4.8 GHz, including polarisation.

Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1′) including Hi.

Zhou *et al.* 2009, ApJ, 691, 516. CO and HCO⁺ of region.

Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9^{\circ}5'S=24.0\pm 1.3$ Jy) including polarisation and review of flux densities.

Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.

G22.7–0.2

RA: $18^{\text{h}}33^{\text{m}}15^{\text{s}}$

Dec: $-09^{\circ}13'$

1-GHz flux/Jy: 33

Spectral index: 0.6

Size/arcmin: 26

Type: S?

Radio: Non-thermal ring in complex region, overlapping G23.3–0.3.

Point sources: Variable radio source near centre.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3'.3\times 3'.4:S=82$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Becker *et al.* 2010, AJ, 140, 157. Variable radio source detection.

Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.

G23.3–0.3

W41

RA: $18^{\text{h}}34^{\text{m}}45^{\text{s}}$

Dec: $-08^{\circ}48'$

1-GHz flux/Jy: 70

Spectral index: 0.5

Size/arcmin: 27

Type: S

Radio: Distorted ring, in complex region, overlapping G22.7–0.2.

X-ray: Possible extended emission, with compact sources.

Point sources: Pulsar association suggested.

Distance: H_I and CO observations indicate 4.2 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3'.3\times 3'.4:S=138$ Jy).

Gaensler & Johnston 1995, MNRAS, 275, L73. Possible pulsar association.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Aharonian *et al.* 2005, Science, 307, 1938. γ -ray detection.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations.

Albert *et al.* 2006, ApJ, 643, L53. γ -ray observations.

Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.

Tian *et al.* 2007, ApJ, 657, L25. VGPS at 1.4 GHz (1') including H_I, plus XMM observations.

Leahy & Tian 2008, AJ, 135, 167. VGPS at 1.4 GHz (1') including H_I, plus CO observations.

Mukherjee *et al.* 2009, ApJ, 691, 1707. XMM and γ -ray observations.

Frail *et al.* 2013, ApJ, 773, L19. OH observations.

Castro *et al.* 2013, ApJ, 774, 36. Fermi observations.

G23.6 + 0.3**RA:** 18^h33^m03^s**Dec:** −08°13′**1-GHz flux/Jy:** 8?**Spectral index:** 0.3**Size/arcmin:** 10?**Type:** ?

Has been suggested this is not an SNR.

Radio: Not well resolved, in complex region.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Pinheiro Gonçalves *et al.* 2011, AJ, 142, 47. IR observations.

G24.7 − 0.6**RA:** 18^h38^m43^s**Dec:** −07°32′**1-GHz flux/Jy:** 8**Spectral index:** 0.5**Size/arcmin:** 15?**Type:** S?

Radio: Incomplete shell, defined in SW.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : S = 12.3 Jy) and Parkes 64-m at 5 GHz (4′ : S = 3.6 Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′.8 : S = 2.2 ± 0.5 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (44″ × 56″ : S = 1.9 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

G24.7 + 0.6**RA:** 18^h34^m10^s**Dec:** −07°05′**1-GHz flux/Jy:** 20?**Spectral index:** 0.2?**Size/arcmin:** 30 × 15**Type:** C?

Radio: Filled-centre, with faint shell, and a compact H_{II} region to the S.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4′.3 : S = 19 ± 3 Jy) and 4.75 GHz (2′.4 : S = 17 ± 4 Jy) and NRO 45-m at 10.2 GHz (2′.7 : S = 15 ± 3 Jy).
 Becker & Helfand 1987, ApJ, 316, 660. VLA at 1.4 GHz (12″), and X-ray upper limit.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

G25.1 − 2.3**RA:** 18^h45^m10^s**Dec:** −08°00′**1-GHz flux/Jy:** 8**Spectral index:** 0.5?**Size/arcmin:** 80 × 30?**Type:** S

Radio: Incomplete shell, extent not well defined.

References:

Gao *et al.* 2011, A&A, 532, A144. Urumqi 25-m at 5 GHz ($9'5'' : S = 3.7 \pm 0.4$ Jy), plus other observations.

G27.4+0.0

4C–04.71

RA: $18^{\text{h}}41^{\text{m}}19^{\text{s}}$ **1-GHz flux/Jy:** 6**Size/arcmin:** 4**Dec:** $-04^{\circ}56'$ **Spectral index:** 0.68**Type:** S

Early references refer to G27.3–0.1 (Kes 73), a supposed larger remnant.

Radio: Incomplete shell.

X-ray: Diffuse emission, with central low period pulsar.

Point sources: Central AXP.

Distance: H α absorption suggests 7.5 to 9.8 kpc.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5'$).

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz ($3' : S = 4.4$ Jy).

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'4''$) and 5 GHz ($4'4''$).

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'8'' : S = 2.0 \pm 0.5$ Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'6''$).

Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz ($60'' \times 45'' : S = 3.5$ Jy). Revise $S_{408 \text{ MHz}} = 10.4$ Jy, and $S_{5 \text{ GHz}} = 1.9 \pm 0.2$ Jy.

Kriss *et al.* 1985, ApJ, 288, 703. Einstein observations, plus VLA at 1.4 and 5 GHz ($12''$).

Sanbonmatsu & Helfand 1992, AJ, 104, 2189. VLA at 1.4 GHz for H α absorption.

Helfand *et al.* 1994, ApJ, 434, 627. ROSAT observations, particularly of central source.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including nearby masers.

Vasisht & Gotthelf 1997, ApJ, 486, L129. ASCA detection of pulsar.

Gotthelf & Vasisht 1997, ApJ, 486, L133. ASCA observations.

Gotthelf *et al.* 1999, ApJ, 522, L49. X-ray timing observations of pulsar.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Mereghetti *et al.* 2001, MNRAS, 321, 143. Search for optical/IR counterpart to pulsar.

Kuiper *et al.* 2004, ApJ, 613, 1173. X-ray observations of pulsar.

Tian & Leahy 2008, ApJ, 677, 292. VGPS at 1.4 GHz ($1'$) including H α .

Mizuno *et al.* 2010, AJ, 139, 1542. Spitzer observations.

An *et al.* 2013, ApJ, 779, 163. NuSTAR and γ -ray observations.

G27.8+0.6

RA: $18^{\text{h}}39^{\text{m}}50^{\text{s}}$ **1-GHz flux/Jy:** 30**Size/arcmin:** 50×30 **Dec:** $-04^{\circ}24'$ **Spectral index:** varies**Type:** F

Radio: Filled-centre, with spectral turnover.

X-ray: Possible pulsar wind nebula.

References:

Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz ($4'3'' : S = 23 \pm 2$ Jy), and 4.75 GHz ($2'4'' : S = 18 \pm 2$ Jy) and NRO 45-m at 10.2 GHz (smoothed to $4'3'' : S = 8.5 \pm 2$ Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

Misanovic *et al.* 2010, ApJ, 725, 931. XMM pulsar/wind nebula search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5''S=21.0\pm 1.1$ Jy) including polarisation and review of flux densities.

G28.6–0.1

RA: $18^{\text{h}}43^{\text{m}}55^{\text{s}}$

Dec: $-03^{\circ}53'$

1-GHz flux/Jy: ?

Spectral index: ?

Size/arcmin: 13×9

Type: S

Radio: Poorly defined regions of non-thermal emission.

X-ray: Diffuse shell, with thermal and non-thermal emission.

References:

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 1.4 GHz ($15''$) and 5 GHz ($15''$).

Bamba *et al.* 2001, PASJ, 53, L21. ASCA observations.

Ueno *et al.* 2003, ApJ, 588, 338. Chandra observations.

Ebisawa *et al.* 2005, ApJ, 635, 214. Chandra observations.

G28.8+1.5

RA: $18^{\text{h}}39^{\text{m}}00^{\text{s}}$

Dec: $-02^{\circ}55'$

1-GHz flux/Jy: ?

Spectral index: 0.4?

Size/arcmin: 100?

Type: S?

Radio: Part of rim detected.

X-ray: Diffuse, Centrally brightened.

References:

Schwentker 1994, A&A, 286, L47. ROSAT observations.

Song *et al.* 2000, PASJ, 52, 181. ASCA observations.

Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

Misanovic *et al.* 2010, ApJ, 725, 931. XMM pulsar/wind nebula search.

G29.6+0.1

RA: $18^{\text{h}}44^{\text{m}}52^{\text{s}}$

Dec: $-02^{\circ}57'$

1-GHz flux/Jy: 1.5?

Spectral index: 0.5?

Size/arcmin: 5

Type: S

Radio: Diffuse shell.

Point sources: AXP associated.

References:

Gaensler *et al.* 1999, ApJ, 526, L37. VLA at 5 GHz ($13''$) and 8 GHz ($8''$)

Vasisht *et al.* 2000, ApJ, 542, L49. X-ray observations of AXP.

G29.7–0.3

Kes 75

RA: 18^h46^m25^s**1-GHz flux/Jy:** 10**Size/arcmin:** 3**Dec:** –02°59′**Spectral index:** 0.63**Type:** C

Has erroneously been called G29.6+0.1.

Radio: Shell with flatter spectrum emission from centre.

X-ray: Thermal shell and non-thermal core, and central pulsar.

Point sources: X-ray pulsar.

Distance: Association with CO implies 11 kpc.

References:

- Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Milne & Dickel 1974, *AujPh*, 27, 549. Parkes 64-m at 2.7 GHz (8′4″: $S=5\pm 20\%$ Jy).
 Becker & Kundu 1975, *AJ*, 80, 679. NRAO 140-ft at 10.6 GHz (3′).
 Becker & Kundu 1976, *ApJ*, 204, 427. NRAO interferometer at 2.7 GHz (20″×7″) and 8 GHz (25″×8″), plus review of flux densities.
 Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6″).
 Becker *et al.* 1983, *ApJ*, 268, L93. VLA at 1.4 GHz (3″), plus Einstein observations.
 Becker & Helfand 1984, *ApJ*, 283, 154. VLA at 5 GHz (2′6″), plus HI.
 Morsi & Reich 1987, *A&AS*, 71, 189. Effelsberg 100-m at 32 GHz (26′5″: $S=1.02\pm 0.07$ Jy).
 Salter *et al.* 1989, *ApJ*, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
 Kassim 1992, *AJ*, 103, 943. VLA at 327 MHz (3′4″×3′7″: $S=27.4$ Jy).
 Biggs & Lyne 1996, *MNRAS*, 282, 691. Pulsar search.
 Blanton & Helfand 1996, *ApJ*, 470, 961. ASCA observations.
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Gotthelf *et al.* 2000, *ApJ*, 542, L37. X-ray pulsar detection.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
 Mereghetti *et al.* 2002, *ApJ*, 574, 873. BeppoSAX observations of pulsar.
 Helfand *et al.* 2003, *ApJ*, 582, 783. Chandra observations.
 Bock & Gaensler 2005, *ApJ*, 626, 343. BIMA at 88.6 GHz (10″×13″).
 Morton *et al.* 2007, *ApJ*, 667, 219. Spitzer and Chandra observations.
 McBride *et al.* 2008, *A&A*, 477, 249. γ -ray and Chandra observations.
 Leahy & Tian 2008, *A&A*, 480, L25. VGPS at 1.4 GHz (1′) including HI.
 Kumar & Safi-Harb 2008, *ApJ*, 678, L43. Chandra observations.
 Ng *et al.* 2008, *ApJ*, 686, 508. Chandra observations.
 Gavriil *et al.* 2008, *Science*, 319, 1802. X-ray observations of pulsar.
 Su *et al.* 2009, *ApJ*, 694, 376. CO observations of region, plus Chandra observations.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9′5″: $S=3.6\pm 0.6$ Jy) including polarisation and review of flux densities.

G30.7–2.0**RA:** 18^h54^m25^s**1-GHz flux/Jy:** 0.5?**Size/arcmin:** 16**Dec:** –02°54′**Spectral index:** 0.7?**Type:** ?

Radio: Poorly defined.

References:

- Reich *et al.* 1988, *IAUCo*, 101, 293. Summary of parameters.
 Reich *et al.* 1990, *A&AS*, 85, 633. Effelsberg 100-m at 2.7 GHz (4′3″).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

G30.7 + 1.0**RA:** 18^h44^m00^s**Dec:** -01°32′**1-GHz flux/Jy:** 6**Spectral index:** 0.4**Size/arcmin:** 24×18**Type:** S?**Radio:** Non-thermal, highly polarised part shell?**Point sources:** Compact source near centre.**References:**Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2′.4 : S = 3.4±0.4 Jy), plus other flux densities.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.6×3′.9 : S = 8.6 Jy).

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 2.93±0.19 Jy) including polarisation and review of flux densities.**G31.5 – 0.6****RA:** 18^h51^m10^s**Dec:** -01°31′**1-GHz flux/Jy:** 2?**Spectral index:** ?**Size/arcmin:** 18?**Type:** S?

Has been called G31.55–0.65.

Radio: Distorted shell? near H_{II} region.**Optical:** Diffuse, incomplete shell.**References:**Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2′.4), plus other flux densities.Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.**G31.9 + 0.0**

3C391

RA: 18^h49^m25^s**Dec:** -00°55′**1-GHz flux/Jy:** 25**Spectral index:** varies**Size/arcmin:** 7×5**Type:** S**Radio:** Shell, brightest in NW, with low frequency turnover.**X-ray:** Diffuse with central core.**Distance:** H_I absorption is seen to the tangent point (8.5 kpc).**References:**Radhakrishnan *et al.* 1972, ApJS, 24, 49. H_I absorption.

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′).

Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15 GHz (58″ : S > 1.6).

Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20″×23″) and 8 GHz (24″×9″), plus review of flux densities.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).Goss *et al.* 1979, A&A, 78, 75. FIRST at 1.4 GHz (54″×66″ : S = 20±2 Jy) and Effelsberg 100-m at 10.7 GHz (77″ : S = 7.5±0.8 Jy).

Wang & Seward 1984, ApJ, 279, 705. Einstein observations.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.4×3′.7 : S = 44.8 Jy).

Reynolds & Moffett 1993, AJ, 105, 2226. VLA at 1.4 GHz (6″), including possible associated CO.

Moffett & Reynolds 1994, ApJ, 425, 668. VLA at 330 MHz (smoothed to 30″ : S = 38.5±0.5 Jy) 1.46 GHz (6′.7) and 4.85 GHz (6′.2×6′.4), including spectral index and polarisation studies.

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.
 Rho & Petre 1996, ApJ, 467, 698. ROSAT observations.
 Wilner *et al.* 1998, AJ, 115, 247. CO observations of surroundings.
 Reach & Rho 1998, ApJ, 507, L93. ISO observations.
 Reach & Rho 1999, ApJ, 511, 836. CO, HCO⁺ and CS observations of surroundings.
 Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Chen & Slane 2001, ApJ, 563, 202. ASCA observations.
 Reach *et al.* 2002, ApJ, 564, 302. Observations of shocked molecular species.
 Chen *et al.* 2004, ApJ, 616, 885. Chandra observations.
 Brogan *et al.* 2005, AJ, 130, 148. VLA at 74 MHz, 330 MHz and 1.5 GHz (70'').
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Castro & Slane 2010, ApJ, 717, 372. Fermi observations.
 Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 8.9±0.6 Jy) including polarisation and review of flux densities.

G32.0—4.9

3C396.1

RA: 19^h06^m00^s**1-GHz flux/Jy:** 22?**Size/arcmin:** 60?**Dec:** -03°00'**Spectral index:** 0.5?**Type:** S?**Radio:** Possible large shell?**References:**

Milne & Hill 1969, AujPh, 22, 211. Parkes 64-m at 635 MHz (31' : S = 25±30% Jy), 1410 MHz (15' : S = 19±15% Jy) and 2650 MHz (8'.4 : S = 8.6±30% Jy). Fluxes if size is 60', plus review of flux densities.
 Caswell 1970, AujPh, 23, 105. Revision of low frequency flux densities.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 111 MHz (S = 105±30 Jy).

G32.1—0.9**RA:** 18^h53^m10^s**1-GHz flux/Jy:** ?**Size/arcmin:** 40?**Dec:** -01°08'**Spectral index:** ?**Type:** C?**Radio:** Possible faint shell, not well defined.**X-ray:** Diffuse, with clumps.**References:**

Folgheraiter *et al.* 1997, MNRAS, 292, 365. ROSAT and ASCA observations.

G32.4 + 0.1**RA:** 18^h50^m05^s**Dec:** -00°25'**1-GHz flux/Jy:** 0.25?**Spectral index:** ?**Size/arcmin:** 6**Type:** S

Has been called G32.45+0.1.

Radio: Shell.

X-ray: Shell.

Distance: X-ray absorption suggests 17 kpc.

References:

Yamaguchi *et al.* 2004, PASJ, 56, 1059. XMM and other observations.

Ueno *et al.* 2005, in XRRC, E4.18. XMM observations.

G32.8 - 0.1**RA:** 18^h51^m25^s**Dec:** -00°08'**1-GHz flux/Jy:** 11?**Spectral index:** 0.2?**Size/arcmin:** 17**Type:** S?

Kes 78

Part has been called G33.1-0.1.

Radio: Elongated shell?

Optical: Detected.

X-ray: NE rim detected.

Distance: Association with CO indicates 4.8 kpc.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5': S=7.2±0.5 Jy).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 430 MHz (S=19.0±15.5 Jy).

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3': S=12.8 Jy) and Parkes 64-m at 5 GHz (4': S=7.7 Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'6×3'8: S=31.3 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

Zhou & Chen 2011, ApJ, 743, 4. XMM observations, plus CO of region.

G33.2 - 0.6**RA:** 18^h53^m50^s**Dec:** -00°02'**1-GHz flux/Jy:** 3.5**Spectral index:** varies**Size/arcmin:** 18**Type:** S

Radio: Incomplete shell.

Optical: Filaments and diffuse emission.

References:

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Reich 1982, A&A, 106, 314. Effelsberg 100-m at 2.7 GHz (4'4: S=2.6±0.3 Jy) and 4.75 GHz (2'5: S=1.75±0.2 Jy).

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (68''×52'' : S=2.7±0.3 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2009, A&A, 499, 789. Optical observations.

G33.6 + 0.1

Kes 79, 4C00.70, HC13

RA: 18^h52^m48^s**1-GHz flux/Jy:** 20**Size/arcmin:** 10**Dec:** +00°41′**Spectral index:** 0.51**Type:** S

Has been called G33.7+0.0.

Radio: Shell, with bright central region, in complex region.

X-ray: Multiple shells and filaments.

Point sources: Central X-ray pulsar.

Distance: H_I absorption gives about 7.8 kpc.

References:

- Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′:S=35.5 Jy) and Parkes 64-m at 5 GHz (4′:S=7.8 Jy).
 Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S₄₃₀ MHz=69±33 Jy.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′:S=6.8±1.5 Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′8″:S=11.4±1.1 Jy).
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1′), plus observations of the nearby point source.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H_I absorption to nearby point source, possibly extragalactic.
 Sequist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source, plus Einstein observations.
 Green 1989, MNRAS, 238, 737. OH absorption.
 Frail & Clifton 1989, ApJ, 336, 854. VLA at 1.4 GHz (1′×2′9″), including H_I absorption.
 Velusamy *et al.* 1991, AJ, 102, 676. VLA at 327 MHz (1′), 1.5 (7″×14″) and 5 GHz (7″), including spectral comparison.
 Green & Dewdney 1992, MNRAS, 254, 686. Observations of adjacent molecular material.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′6″×3′8″:S=34.8 Jy).
 Seward & Velusamy 1995, ApJ, 439, 715. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Tsunemi & Enoguchi 2002, PASJ, 54, 735. ASCA observations.
 Seward *et al.* 2003, ApJ, 584, 414. Chandra observations.
 Stanimirović 2003, ApJ, 592, 953. Arecibo OH absorption.
 Sun *et al.* 2004, ApJ, 605, 742. Chandra observations.
 Gotthelf *et al.* 2005, ApJ, 627, 390. XMM pulsar detection.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Halpern *et al.* 2007, ApJ, 665, 1304. X-ray pulsar timing observations.
 Giacani *et al.* 2009, A&A, 507, 841. VLA at 74 MHz (36″×39″:S=76±10 Jy), 324 MHz (13″:S=39±8 Jy) and 1.5 GHz (17″×19″:S=11.5±1.5 Jy), plus review of flux densities and XMM observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5″:S=9.4±0.5 Jy) including polarisation and review of flux densities.

G34.7–0.4

W44, 3C392

RA: 18^h56^m00^s**1-GHz flux/Jy:** 250**Size/arcmin:** 35×27**Dec:** +01°22′**Spectral index:** 0.37**Type:** C

Has been called G34.6–0.5.

Radio: Distorted shell, brighter to the E, with pulsar and associated nebula.

Optical: Diffuse emission.

X-ray: Centrally concentrated, thermal spectrum, plus pulsar wind nebula.

Point sources: Pulsar within the boundary of the remnant.

Distance: H_I absorption indicates 2.8 kpc.

References:

- Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10.7 GHz (3′:S=105±7 Jy).
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′:S=177±10 Jy).
 Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.
 Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′:S=299 Jy).
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430 MHz}=540±187 Jy.
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Smith *et al.* 1985, MNRAS, 217, 99. Einstein observations.
 Wolszczan *et al.* 1991, ApJ, 372, L99. Pulsar detection.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.6×3′.8:S=469 Jy).
 Jones *et al.* 1993, MNRAS, 265, 631. VLA at 1.4 GHz (15″), plus X-ray spectra.
 Rho *et al.* 1994, ApJ, 430, 757. Optical and ROSAT observations.
 Koo & Heiles 1995, ApJ, 442, 679. H_I of surrounding shell.
 Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ -ray emission.
 Harrus *et al.* 1996, ApJ, 464, L161. ASCA observations.
 Frail *et al.* 1996, ApJ, 464, L165. VLA at 1.5 and 8.4 GHz (8″.9×7″.8) of pulsar nebula.
 Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.
 Harrus *et al.* 1997, ApJ, 488, 781. X-ray observations.
 Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.
 Giacani *et al.* 1997, AJ, 113, 1379. VLA at 1.4 GHz (15″), plus optical images.
 Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.
 Cox *et al.* 1999, ApJ, 524, 179. Revision of distance.
 Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
 Petre *et al.* 2002, ApJ, 579, 404. Chandra observations of pulsar and wind nebula.
 Mavromatakis *et al.* 2003, A&A, 405, 591. Optical observations.
 Shelton *et al.* 2004, ApJ, 611, 906. Chandra observations of part.
 Reach *et al.* 2005, ApJ, 618, 297. Molecular line and near-IR observations.
 Hoffman *et al.* 2005, ApJ, 627, 803. Observations of OH masers.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
 Castelletti *et al.* 2007, A&A, 471, 537. VLA at 74 MHz (36″×39″:S=634±70 Jy) and 324 MHz (13″:S=411±50 Jy).
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Abdo *et al.* 2010, Science, 327, 1103. Fermi observations.
 Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 118±6 Jy) including polarisation and review of flux densities.
 Giuliani *et al.* 2011, ApJ, 742, L30. γ -ray observations.
 Uchiyama *et al.* 2012, ApJ, 749, L35. Fermi observations.
 Uchida *et al.* 2012, PASJ, 64, 141. Suzaku observations.
 Ackermann *et al.* 2013, Science, 339, 807. Fermi observations.
 Yoshiike *et al.* 2013, ApJ, 768, 179. CO and H α observations of region.
 Sashida *et al.* 2013, ApJ, 774, 10. HCO $^+$ and CO observations of region.
 Park *et al.* 2013, ApJ, 777, 14. Arecibo H α observations of region.

G35.6–0.4

RA: 18^h57^m55^s
Dec: +02°13'

1-GHz flux/Jy: 9
Spectral index: 0.5

Size/arcmin: 15×11
Type: S?

Re-identified as SNR in 2009.

Radio: Diffuse, with some limb brightening.

References:

Green 2009, MNRAS, 399, 177. Identification in the radio as a SNR.
 Paron & Giacani 2010, A&A, 509, L4. CO and IR observations of region.
 Zhu *et al.* 2013, ApJ, 775, 95. H α and other observations.

G36.6–0.7

RA: 19^h00^m35^s
Dec: +02°56'

1-GHz flux/Jy: 1.0
Spectral index: 0.7?

Size/arcmin: 25?
Type: S?

Radio: polarised arc, possibly part of a larger shell?

References:

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4), plus other flux densities.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.4×3'.7 : S = 6.7 Jy).
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 0.39±0.04 Jy) including polarisation and review of flux densities.

G36.6+2.6

RA: 18^h48^m49^s
Dec: +04°26'

1-GHz flux/Jy: 0.7?
Spectral index: 0.5?

Size/arcmin: 17×13?
Type: S

Radio: Poorly resolved shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

G38.7–1.3**RA:** 19^h06^m40^s**Dec:** +04°28′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 32×19?**Type:** S

G38.7–1.4 refers to the E portion.

Radio: Incomplete shell.**Optical:** Arc of filaments, brighter to E.**X-ray:** Detected in E.**References:**Schauadel *et al.* 2002, ASPC, 271, 391. ROSAT observations of E.Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.**G39.2–0.3**

3C396, HC24, NRAO 593

RA: 19^h04^m08^s**Dec:** +05°28′**1-GHz flux/Jy:** 18**Spectral index:** 0.34**Size/arcmin:** 8×6**Type:** C**Radio:** Shell, brighter to W, with faint ‘tail’ to E.**X-ray:** Diffuse, brighter to W, with central core.**Point sources:** Central X-ray source.**Distance:** H α absorption suggests > 7.7 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S₄₃₀ MHz = 54±38 Jy.

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′:S=4.1±1.0 Jy).

Caswell *et al.* 1975, A&A, 45, 239. H α absorption.Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (65″×48″).

Becker & Helfand 1987, AJ, 94, 1629. VLA at 1.4 GHz (12″:S=14 Jy) and 5 GHz, plus Einstein observations.

Patnaik *et al.* 1990, A&A, 232, 467. VLA at 1.5 GHz (25″) and 1.4 GHz (7″8×7″5) and 5 GHz (25″) including polarisation, plus Ooty at 327 MHz (100″×31″), including review of flux densities.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′5×3′6:S=42.5 Jy).

Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz for spectral index studies.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

Harrus & Slane 1999, ApJ, 516, 811. ASCA observations.

Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.Olbert *et al.* 2003, ApJ, 592, L45. Chandra observations.Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.Scaife *et al.* 2007, MNRAS, 377, L69. 33 GHz observations.Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.Lee *et al.* 2009, ApJ, 691, 1042. IR observations.Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5:S=8.8±0.5 Jy) including polarisation and review of flux densities.Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.Su *et al.* 2011, ApJ, 727, 43. Chandra and CO observations of region.

G39.7–2.0

W50, SS433

RA: 19^h12^m20^s**1-GHz flux/Jy:** 85?**Size/arcmin:** 120×60**Dec:** +04°55′**Spectral index:** 0.7?**Type:** ?

Eastern part has been called G40.0–3.1. Is this a SNR?

Radio: Elongated shell, containing SS433, adjacent to the H_{II} region S74.

Optical: Faint filaments at the edge of the radio emission.

X-ray: Emission from SS433 and two lobes.

Point sources: SS433 is the compact source in the centre of the W50.

Distance: H_I absorption indicates 6.0±0.5 kpc.

References:

van den Bergh 1980, ApJ, 236, L23. Optical in H α and [S_{II}].

Zealey *et al.* 1980, MNRAS, 192, 731. Optical spectra.

van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H_I absorption to nearby point source (not SS433).

Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.

Watson *et al.* 1983, ApJ, 273, 688. X-ray observations.

Downes *et al.* 1986, MNRAS, 218, 393. Effelsberg 100-m at 4.75 GHz (2′.4 : S = 34±4 Jy), plus previous 1.7 and 2.7 GHz data.

Romney *et al.* 1987, ApJ, 321, 822. VLBI of SS433, including distance.

Elston & Baum 1987, AJ, 94, 1633. Mosaic with VLA at 1.4 GHz (30′′) of fine structure only.

Kawai *et al.* 1989, PASJ, 41, 491. X-ray observations of SS433.

Band 1989, ApJ, 336, 937. Einstein and EXOSAT observations.

Yamauchi *et al.* 1994, PASJ, 46, L109. X-ray spectral observations.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Brinkmann *et al.* 1996, A&A, 312, 306. ROSAT observations.

Kotani *et al.* 1996, PASJ, 48, 619. X-ray line emission from SS433.

Safi-Harb & Ögelman 1997, ApJ, 483, 868. ROSAT observations.

Dubner *et al.* 1998, AJ, 116, 1842. VLA at 328 MHz (64′′×60′′ : S = 160±20 Jy), and 1.4 GHz (56′′×54′′), plus NRAO 140-ft at 1.4 GHz (21′) for H_I observations.

Safi-Harb & Petre 1999, ApJ, 512, 784. X-ray observations.

Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.

Brinkmann *et al.* 2007, A&A, 463, 611. XMM observations of E lobe.

Boumis *et al.* 2007, MNRAS, 381, 308. Optical observations.

Lockman *et al.* 2007, MNRAS, 381, 881. H_I observations.

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 37±4 Jy), including polarisation and review of flux densities.

G40.5–0.5**RA:** 19^h07^m10^s**1-GHz flux/Jy:** 11**Size/arcmin:** 22**Dec:** +06°31′**Spectral index:** 0.4**Type:** S

Radio: Shell, brightest to the NE.

References:

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).

Downes *et al.* 1980, A&A, 92, 47. Effelsberg 100-m at 1.7 GHz (7′.6 : S = 9.3±1.3 Jy), and 2.7 GHz (4′.4 : S = 7.2±0.5 Jy), plus review of flux densities.

Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.

Yang *et al.* 2006, ChJAA, 6, 210. CO observations of surroundings.

Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 6.4 \pm 0.3$ Jy) including polarisation and review of flux densities.

G41.1–0.3

3C397

RA: 19^h07^m34^s**1-GHz flux/Jy:** 25**Size/arcmin:** 4.5×2.5**Dec:** +07°08'**Spectral index:** 0.50**Type:** S

Radio: 3C397 is two sources: the E is the SNR, the W is a H_{II} region.

X-ray: Brighter to the E and W, with central component.

Distance: Possible limit of > 7.5 kpc for non-thermal component from H_I absorption.

References:

Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 10 GHz (3') and 5 GHz (6').

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5' : S = 21.3±1.2 Jy for both components).

Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430 MHz} = 82±51 Jy, also Algonquin 46-m at 10.6 GHz (3' : S = 12±2 Jy), and Haystack 36-m at 15.5 GHz (2'3 : S = 8.5±3.0 Jy).

Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : S = 29.8 Jy) and Parkes 64-m at 5 GHz (4' : S = 8.7 Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (58''×52'').

Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz (8''), plus Einstein observations.

Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30'' : S = 4.10±0.19 Jy).

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.5×3'.6 : S = 46.3 Jy).

Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz, for spectral index studies.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Chen *et al.* 1999, ApJ, 520, 737. ASCA and ROSAT observations.

Dyer & Reynolds 1999, ApJ, 526, 365. VLA at 1.5 GHz (6''.9×6''.6) and 4.8 GHz (46''.4×5''.6), including polarisation and comparison with ROSAT image.

Safi-Harb *et al.* 2000, ApJ, 545, 922. ROSAT, ASCA and other X-ray observations.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.

Safi-Harb *et al.* 2005, ApJ, 618, 321. Chandra observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Jiang *et al.* 2010, ApJ, 712, 1147. CO observations of region.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 18.5±1.1 Jy) including polarisation and review of flux densities.

Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.

G41.5 + 0.4**RA:** 19^h05^m50^s**Dec:** +07°46′**1-GHz flux/Jy:** 1?**Spectral index:** ?**Size/arcmin:** 10**Type:** S?**Radio:** Partial clumpy shell, brighter to NE.**References:**Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 332 MHz (20″: $S = 1.8 \pm 0.4$ Jy).Alves *et al.* 2012, MNRAS, 422, 2429. Radio observations.**G42.0 – 0.1****RA:** 19^h08^m10^s**Dec:** +08°00′**1-GHz flux/Jy:** 0.5?**Spectral index:** ?**Size/arcmin:** 8**Type:** S?**Radio:** Irregular shell.**References:**Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 332 MHz (20″: $S = 1.8 \pm 0.4$ Jy).Alves *et al.* 2012, MNRAS, 422, 2429. Radio observations.**G42.8 + 0.6****RA:** 19^h07^m20^s**Dec:** +09°05′**1-GHz flux/Jy:** 3?**Spectral index:** 0.5?**Size/arcmin:** 24**Type:** S

Has been called G42.8+0.65.

Radio: Faint shell.**Point sources:** Near soft gamma repeater, and young pulsar.**References:**Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2′.4: $S = 1.5 \pm 0.2$ Jy), plus other flux densities.Vasisht *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3′.2×3′.4).Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.

Lorimer & Xilouris 2000, ApJ, 545, 385. Pulsar detection.

Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50″), and other observations of the region.**G43.3 – 0.2****RA:** 19^h11^m08^s**Dec:** +09°06′**1-GHz flux/Jy:** 38**Spectral index:** 0.46**Size/arcmin:** 4×3**Type:** S

W49B

Radio: Shell, brightest to the SE and W, near the H_{II} region W49A.**X-ray:** Centrally brightened, elongated E--W.**Point sources:** Compact X-ray source.**Distance:** H_I absorption indicates 10 kpc.**References:**

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 2.7 GHz (4').
 Downes & Wilson 1974, *A&A*, 34, 133. Effelsberg 100-m at 10.7 GHz (1'3).
 Green *et al.* 1975, *A&A*, 44, 187. Effelsberg 100-m at 15.0 GHz (58'' : $S=9.0\pm 0.7$ Jy).
 Lockhart & Goss 1978, *A&A*, 67, 355. H α absorption.
 Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Pye *et al.* 1984, *MNRAS*, 207, 649. Einstein observations, and VLA at 1.4 and 5 GHz (both 12'').
 Smith *et al.* 1985, *ApJ*, 296, 469. EXOSAT spectrum.
 Morsi & Reich 1987, *A&AS*, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30'' : $S=6.90\pm 0.38$ Jy).
 Moffett & Reynolds 1994, *ApJ*, 437, 705. VLA at 330 MHz (6''7 \times 7''7 : $S=64.4$ Jy), 1.48 GHz (4''8 \times 5''2 : $S=31.8$ Jy) and 4.85 GHz (4''0 \times 4''1), including polarisation.
 Fujimoto *et al.* 1995, *PASJ*, 47, L31. ASCA observations.
 Gorham *et al.* 1996, *ApJ*, 458, 257. Pulsar search.
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Hwang *et al.* 2000, *ApJ*, 532, 970. ROSAT image and ASCA spectroscopy.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
 Aharonian *et al.* 2001, *A&A*, 375, 1008. Limit on high energy γ -rays.
 Brogan & Troland 2001, *ApJ*, 550, 799. VLA at 1.4 GHz (27'' \times 24'' and 5'') for H α Zeeman splitting.
 Lacey *et al.* 2001, *ApJ*, 559, 954. VLA at 74 MHz (26'' \times 23'' : $S=55.6$ Jy) and 326 MHz (6''6 \times 6''2 : $S=56.0$ Jy).
 Kaplan *et al.* 2002, *ApJ*, 566, 378. VLA at 333 MHz (50''), and other observations of the region.
 Kawasaki *et al.* 2005, *ApJ*, 631, 935. ASCA observations.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Miceli *et al.* 2006, *A&A*, 453, 567. XMM observations.
 Keohane *et al.* 2007, *ApJ*, 654, 938. IR and Chandra observations.
 Ozawa *et al.* 2009, *ApJ*, 706, L71. Suzaku observations.
 Abdo *et al.* 2010, *ApJ*, 722, 1303. Fermi observations.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9'5 : $S=19.1\pm 1.0$ Jy) including polarisation and review of flux densities.
 Rodes-Roca 2013, *A&A*, 555, A115. IR of compact X-ray source.
 Yang *et al.* 2013, *ApJ*, 766, 44. Suzaku spectroscopy.
 Lopez *et al.* 2013, *ApJ*, 777, 145. Chandra spectroscopy.

G43.9 + 1.6

RA: 19^h05^m50^s

Dec: +10°30'

1-GHz flux/Jy: 9.0

Spectral index: 0.5

Size/arcmin: 60?

Type: S?

Radio: Large, poorly defined faint shell.

Point sources: Soft gamma repeater nearby.

References:

Reich *et al.* 1988, *IAUCo*, 101, 293. Summary of parameters.
 Reich *et al.* 1990, *A&AS*, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Vasisht *et al.* 1994, *ApJ*, 431, L35. VLA at 327 MHz (3'2 \times 3'4).
 Hurley *et al.* 1996, *ApJ*, 463, L13. Observations of soft gamma repeater field.
 Kaplan *et al.* 2002, *ApJ*, 566, 378. VLA at 333 MHz (50''), and other observations of the region.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9'5 : $S=4.55\pm 0.24$ Jy) including polarisation and review of flux densities.

G45.7–0.4**RA:** 19^h16^m25^s**Dec:** +11°09′**1-GHz flux/Jy:** 4.2?**Spectral index:** 0.4?**Size/arcmin:** 22**Type:** S**Radio:** Shell, brightest to the SE, poorly defined to NW.**References:**

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2′4″:S=2.6±0.3 Jy), plus other flux densities.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G46.8–0.3

(HC30)

RA: 19^h18^m10^s**Dec:** +12°09′**1-GHz flux/Jy:** 17**Spectral index:** 0.54**Size/arcmin:** 17×13**Type:** S

Has been called G46.6–0.2.

Radio: Shell, two bright arcs to NNW and SSE.**Distance:** H_I absorption suggests 6.8--8.8 kpc.**References:**

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′:S=9.8±0.9 Jy), and 37-m at 1.7 GHz (S=14.5±5.5 Jy).

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′:S=20.3 Jy) and Parkes 64-m at 5 GHz (4′:S=7.1 Jy).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430 MHz}=46±21 Jy.

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′8″:S=5.9±0.6 Jy).

Sato 1979, ApL, 20, 43. H_I observations.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6″).

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (58″×53″:S=13.3±0.1 Jy).

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5″:S=7.02±0.18 Jy) including polarisation and review of flux densities.

G49.2–0.7

(W51)

RA: 19^h23^m50^s**Dec:** +14°06′**1-GHz flux/Jy:** 160?**Spectral index:** 0.3?**Size/arcmin:** 30**Type:** S?**Radio:** In complex region, parameters uncertain.**Optical:** Some diffuse emission possibly associated.**X-ray:** Elongated east--west.**Distance:** Association with CO gives 6 kpc. H_I suggest 4.3 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Parkes 64-m at 5 GHz (4′).

Sato 1973, PASJ, 25, 135. H_I absorption.

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′), S=51.5±3.2 Jy, for the non-thermal component, but probably confused.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6″).

Seward 1990, ApJS, 73, 781. Einstein observations.

Copetti & Schmidt 1991, MNRAS, 250, 127. 151 MHz observations.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'1×3'5).
 Subrahmanyan & Goss 1995, MNRAS, 275, 755. VLA at 330 MHz (1'1).
 Koo *et al.* 1995, ApJ, 447, 211. ROSAT observations.
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. OH masers.
 Koo & Moon 1997, ApJ, 475, 194. Arecibo (3') and VLA (40''×42'') at 1.4 GHz for H_i.
 Koo & Moon 1997, ApJ, 485, 263. NRAO 12-m CO and HCO⁺ observations.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.
 Koo *et al.* 2002, AJ, 123, 1629. ASCA observations.
 Koo *et al.* 2005, ApJ, 633, 946. Chandra observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_i.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Abdo *et al.* 2009, ApJ, 706, L1. Fermi observations.
 Koo *et al.* 2010, AJ, 140, 262. H_i Zeeman splitting observations.
 Ceccarelli *et al.* 2011, ApJ, 740, L4. Molecular line observations of region.
 Aleksić *et al.* 2012, A&A, 541, A13. γ -ray observations.
 Hanabata *et al.* 2013, PASJ, 65, 42. Suzaku observations.
 Tian & Leahy 2013, ApJ, 769, L17. H_i observations of region.
 Brogan *et al.* 2013, ApJ, 771, 91. VLA at 74 MHz (84''×92'') and 320 MHz (33''×35''), plus OH, molecular line and other observations.
 Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_i in region.

G53.6–2.2

3C400.2, NRAO 611

RA: 19^h38^m50^s**1-GHz flux/Jy:** 8**Size/arcmin:** 33×28**Dec:** +17°14'**Spectral index:** 0.50**Type:** S

Has been called G53.7–2.2.

Radio: Ring of emission, with extension to NW.

Optical: Filaments and diffuse emission.

X-ray: Centrally brightened, offset to NW.

Distance: Association with H_i gives 2.8 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5':S=5.3±0.6 Jy).
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5':S=4.8±0.3 Jy).
 Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3':S=11.7 Jy).
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430 MHz}=20±10 Jy, S_{318 MHz}=20±3.6 Jy.
 Goss *et al.* 1975, A&A, 43, 459. WSRT at 610 MHz (1'×3':S=13.2±1.6 Jy).
 Sabbadin & d'Odorico 1976, A&A, 49, 119. Optical spectra.
 van den Bergh 1978, ApJS, 38, 119. Optical observations.
 Rosado 1983, RMxAA, 8, 59. Optical spectra.
 Blair & Long 1988, PASP, 100, 461. Optical imaging and spectroscopy.
 see also: Blair & Long 1988, PASP, 100, 651. Erratum.
 Long *et al.* 1991, ApJ, 373, 567. Einstein and optical observations.
 Winkler *et al.* 1993, ApJ, 405, 608. Optical imaging.
 Dubner *et al.* 1994, AJ, 108, 207. VLA at 327 MHz (59'') and 1.49 GHz (52''), plus X-rays.
 Saken *et al.* 1995, ApJ, 443, 231. ROSAT observations.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Giacani *et al.* 1998, A&AS, 133, 61. DRAO at 1.4 GHz for H_i studies, including distance.
 Yoshita *et al.* 2001, PASJ, 53, 93. ASCA observations, and spectral comparison with ROSAT.

Ambrocio-Cruz *et al.* 2006, RMxAA, 42, 241. Optical imaging and spectroscopy.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 4.00 \pm 0.22$ Jy) including polarisation and review of flux densities.

G54.1 + 0.3

RA: $19^{\text{h}}30^{\text{m}}31^{\text{s}}$
Dec: $+18^{\circ}52'$

1-GHz flux/Jy: 0.5
Spectral index: 0.1

Size/arcmin: 12?
Type: C?

Radio: Filled-centre core, with faint diffuse emission.

X-ray: Centrally concentrated, with more extended diffuse emission.

Point sources: Central pulsar.

Distance: H α absorption suggests 4.5--9 kpc, association with CO suggest 8.2 kpc.

References:

Green 1985, MNRAS, 216, 691. Radio at 2.7 GHz ($7'' \times 20''$).
 Reich *et al.* 1985, A&A, 151, L10. Effelsberg 100-m at 4.75 GHz ($2'.4 : S = 0.37 \pm 0.04$ Jy)
 Velusamy & Becker 1988, AJ, 95, 1162. VLA at 1.4 ($14'' : S = 0.48 \pm 0.03$ Jy), 1.6 ($14'' : S = 0.42 \pm 0.03$ Jy) and 5 GHz ($5'' : S = 0.33 \pm 0.02$ Jy), Ooty at 327 MHz ($S = 0.50 \pm 0.08$ Jy), plus review of flux densities.
 Seward 1989, AJ, 97, 481. Einstein observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Lu *et al.* 2001, A&A, 370, 570. ROSAT and ASCA observations.
 Lu *et al.* 2002, ApJ, 568, L49. Chandra observations.
 Camilo *et al.* 2002, ApJ, 574, L71. Pulsar detection.
 Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.
 Leahy *et al.* 2008, AJ, 136, 1477. VGPS at 1.4 GHz ($1'$) including H α .
 Koo *et al.* 2008, ApJ, 673, L147. AKARI observations of surroundings.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Bocchino *et al.* 2010, A&A, 520, A71. XMM and Suzaku observations.
 Lang *et al.* 2010, ApJ, 709, 1125. VLA at 1.4 GHz ($6'.6 \times 6'.8$), 4.7 GHz ($3''.2 \times 3''.3$), and 8.2 GHz ($3''.0 \times 3''.2$) and Spitzer observations.
 Acciari *et al.* 2010, ApJ, 719, L69. γ -ray observations.
 Lee *et al.* 2012, JKAS, 45, 117. CO observations of region.

G54.4–0.3

RA: $19^{\text{h}}33^{\text{m}}20^{\text{s}}$
Dec: $+18^{\circ}56'$

1-GHz flux/Jy: 28
Spectral index: 0.5

(HC40)
Size/arcmin: 40
Type: S

Has been called G54.5–0.3.

Radio: Shell, in complex region.

Optical: Faint filaments.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 34.4 \pm 5.0$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'.6$).
 Caswell 1985, AJ, 90, 1224. DRAO at 1.4 GHz ($1'.3 \times 2'.6 : S = 18 \pm 4$ Jy).
 Velusamy *et al.* 1986, JApA, 7, 105. WSRT at 609 MHz ($50'' \times 191''$ smoothed to $100'' \times 200''$).

Junkes *et al.* 1992, A&AS, 96, 1. Surrounding CO.
 Junkes *et al.* 1992, A&A, 261, 289. Nearby IRAS sources.
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_I.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_I in region.

G55.0 + 0.3

RA: 19^h32^m00^s
Dec: +19°50′

1-GHz flux/Jy: 0.5?
Spectral index: 0.5?

Size/arcmin: 20×15?
Type: S

Has been called G55.2+0.5.

Radio: Faint, partial shell.

Point sources: Old pulsar nearby.

Distance: Association with H_I features implies 14 kpc.

References:

Matthews *et al.* 1998, ApJ, 493, 312. WSRT at 327 MHz (1′.0×2′.9 : S = 0.98±0.15 Jy), DRAO at 1.4 GHz (1′.0×2′.9 : S = 0.25±0.12 Jy), plus H_I observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G55.7 + 3.4

RA: 19^h21^m20^s
Dec: +21°44′

1-GHz flux/Jy: 1?
Spectral index: 0.3?

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Point sources: Old pulsar within the boundary of the remnant.

References:

Goss *et al.* 1977, A&A, 61, 93. WSRT observations at 610 MHz (57″×156″ : S = 1.9±0.2 Jy) and 1415 MHz (27″×72″ : S = 1.0±0.1 Jy).
 Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 1.3 to 1.9 GHz (30″).
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 0.52±0.03 Jy) including polarisation and review of flux densities.

G57.2 + 0.8

(4C21.53)

RA: 19^h34^m59^s
Dec: +21°57′

1-GHz flux/Jy: 1.8
Spectral index: 0.62

Size/arcmin: 12?
Type: S?

Radio: Extended non-thermal arc.

Point sources: Near the millisecond pulsar, but not thought to be related.

References:

Sieber & Seiradakis 1984, A&A, 130, 257. Effelsberg 100-m at 1.4 GHz (8′.8 : 1.34±0.1), 2.7 GHz (4′.3 : 0.86±0.1), plus other surveys of the area.
 Caswell *et al.* 1985, AJ, 90, 488. DRAO at 1.4 GHz (1′×3′).
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5:S=0.74±0.04 Jy) including polarisation and review of flux densities.

G59.5 + 0.1

RA: 19^h42^m33^s
Dec: +23°35'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 15
Type: S

Has been called G59.6+0.1.

Radio: Incomplete shell.

Optical: Diffuse shell.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1'0×2'5 : S = 5.1±0.2 Jy), and northern sky survey at 4.9 GHz.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Xu & Wang 2012, A&A, 543, A24. CO observations of SE.

G59.8 + 1.2

RA: 19^h38^m55^s
Dec: +24°19'

1-GHz flux/Jy: 1.5
Spectral index: 0.0

Size/arcmin: 20×16?
Type: ?

Has been called G59.7+1.2.

Radio: Poorly defined source.

Optical: Faint diffuse emission and filaments.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.

Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.

Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).

Boumis *et al.* 2005, A&A, 443, 175. Optical observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5:S=1.43±0.08 Jy) including polarisation and review of flux densities.

G63.7 + 1.1

RA: 19^h47^m52^s
Dec: +27°45'

1-GHz flux/Jy: 1.8
Spectral index: 0.24

Size/arcmin: 8
Type: F

Radio: Centrally brightened, with core.

References:

Wallace *et al.* 1997, AJ, 114, 2068. WSRT at 1.4 GHz (14''×26'' : S = 1.63 Jy), DRAO at 1.4 GHz (smoothed to 2''), plus review of flux densities and other observations.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5:S=1.12±0.06 Jy) including polarisation and review of flux densities.

G64.5 + 0.9**RA:** 19^h50^m25^s**Dec:** +28°16′**1-GHz flux/Jy:** 0.15?**Spectral index:** 0.5**Size/arcmin:** 8**Type:** S?**Radio:** Shell with central source.**References:**Hurley-Walker *et al.* 2009, MNRAS, 398, 249. Radio identification.**G65.1 + 0.6****RA:** 19^h54^m40^s**Dec:** +28°35′**1-GHz flux/Jy:** 5.5**Spectral index:** 0.61**Size/arcmin:** 90×50**Type:** S**Radio:** Large, faint shell.**Point sources:** Old pulsar nearby.**Distance:** Possible association with H_I suggests 9 kpc.**References:**Landecker *et al.* 1990, A&A, 232, 207. DRAO at 408 MHz (3′.5×7′.0: S=9.5±0.1 Jy), and 1.4 GHz (1′.0×2′.0: S=5.4±1.0 Jy).Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.Tian & Leahy 2006, A&A, 455, 1053. CGPS at 408 MHz (2′.8×5′.9: S=8.6±0.8 Jy) and 1.4 GHz (0′.8×1′.7: S=4.9±0.5 Jy) including H_I.Kotthes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′: S=9.1±1.0 Jy) and 1420 MHz (~1′: S=3.9±0.5 Jy), including review of flux densities.Aleksić *et al.* 2010, ApJ, 725, 1629. Fermi observations.Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5: S=3.2±0.3 Jy), including polarisation and review of flux densities.**G65.3 + 5.7****RA:** 19^h33^m00^s**Dec:** +31°10′**1-GHz flux/Jy:** 42**Spectral index:** 0.6**Size/arcmin:** 310×240**Type:** S?

Has been called G65.2+5.7.

Radio: Large, faint ring, near S91 and S94.**Optical:** Filamentary ring.**X-ray:** Diffuse, centrally brightened.**Distance:** Optical proper motions and velocities indicates 0.8 kpc.**References:**Gull *et al.* 1977, ApJ, 215, L69. Optical plates.Reich *et al.* 1979, A&A, 72, 270. Effelsberg 100-m observations at 1.42 GHz (smoothed to 11′: S=42.4±1.6 Jy), estimate $S_{408\text{ MHz}}=91\pm 5$ Jy from previous sky survey.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Rosado 1981, ApJ, 250, 222. Optical interferometry.

Fesen *et al.* 1983, ApJS, 51, 337. Deep [O III] imagery.Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.

Seward 1990, ApJS, 73, 781. Einstein observations.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Mavromatakis *et al.* 2002, A&A, 388, 355. Optical observations.
 Boumis *et al.* 2004, A&A, 424, 583. Optical observations.
 Shelton *et al.* 2004, ApJ, 615, 275. ROSAT observations.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Xiao *et al.* 2009, A&A, 503, 827. Effelsberg 100-m at 2.7 GHz ($4'4'' : S = 22 \pm 3$ Jy), and Urumqi 21-m at 4.8 GHz ($9'5'' : S = 16.8 \pm 1.8$ Jy) including polarisation and review of flux densities.
 Gosachinskii 2010, AstL, 36, 260. H α observations.
 Kim *et al.* 2010, ApJ, 722, 388. Far UV observations.

G65.7 + 1.2

DA 495

RA: 19^h52^m10^s**1-GHz flux/Jy:** 5.1**Size/arcmin:** 22**Dec:** +29°26'**Spectral index:** varies**Type:** F

Has mistakenly been called G55.7+1.2.

Radio: Centrally brightened with thick shell?

X-ray: Detected.

Point sources: Compact X-ray source near centre.

Distance: H α polarisation observations suggest 1.5 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 2.8 \pm 0.4$ Jy), and 37-m at 1.7 GHz ($S = 4.4 \pm 0.5$ Jy), plus review of flux densities.
see also: Willis 1973, A&A, 27, 483. Erratum.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430 \text{ MHz}} = 8.7 \pm 4.9$ Jy, $S_{318 \text{ MHz}} = 9.7 \pm 2.2$ Jy.
 Landecker & Caswell 1983, AJ, 88, 1810. DRAO at 1.4 GHz ($0'9'' \times 1'5'' : S = 4.4 \pm 0.2$ Jy).
 Velusamy *et al.* 1989, JApA, 10, 161. Ooty at 327 MHz ($36'' \times 64''$), WSRT at 610 MHz ($62''$) and VLA at 1.4 GHz ($36''$ and $12''$), including IRAS imaging.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2004, ApJ, 607, 855. H α polarisation absorption.
 Arzoumanian *et al.* 2004, ApJ, 610, L101. ROSAT and ASCA observations of compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 6.5 \pm 0.6$ Jy) and 1420 MHz ($\sim 1' : S = 4.0 \pm 0.2$ Jy), including polarisation and review of flux densities.
 Arzoumanian *et al.* 2008, ApJ, 687, 505. Chandra observations.
 Kothes *et al.* 2008, ApJ, 687, 516. CGPS at 408 MHz ($2'9'' \times 6'0'' : S = 6.5 \pm 0.5$ Jy) and 1.4 GHz ($0'82'' \times 1'75'' : S = 4.0 \pm 0.2$ Jy), Effelsberg 100-m at 4.85 GHz ($2'45'' : S = 1.6 \pm 0.1$ Jy) and 10.55 GHz (smoothed to $2'45'' : S = 1.1 \pm 0.1$ Jy), plus review of flux densities.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5'' : S = 1.95 \pm 0.10$ Jy) including polarisation and review of flux densities.

G65.8 – 0.5**RA:** 19^h59^m20^s**1-GHz flux/Jy:** ?**Size/arcmin:** 10 × 6?**Dec:** +28°38'**Spectral index:** ?**Type:** S

Radio: Arc in W.

Optical: Diffuse shell, brighter in W.

References:

Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G66.0–0.0

RA: 19^h57^m50^s
Dec: +29°03′

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 31×25?
Type: S

Radio: Some emission in N.

Optical: Incomplete shell.

References:

Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G67.6+0.9

RA: 19^h57^m45^s
Dec: +30°53′

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 50×45?
Type: S

Radio: Arc in S.

Optical: Filamentary shell.

References:

Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G67.7+1.8

RA: 19^h54^m32^s
Dec: +31°29′

1-GHz flux/Jy: 1.0
Spectral index: 0.61

Size/arcmin: 15×12
Type: S

Radio: Double arc shell.

Optical: Filaments in N.

X-ray: Detected.

Point sources: Compact X-ray source.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1′9×1′0: $S = 1.9 \pm 0.1$ Jy), and northern sky survey at 4.9 GHz ($S = 0.42 \pm 0.05$ Jy).

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S = 1.1 \pm 0.1$ Jy) and 1420 MHz ($\sim 1'$: $S = 0.68 \pm 0.04$ Jy), including polarisation and review of flux densities.

Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Hui & Becker 2009, A&A, 494, 1005. Chandra observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5: $S = 0.30 \pm 0.03$ Jy) including polarisation and review of flux densities.

G67.8+0.5**RA:** 20^h00^m00^s**Dec:** +30°51'**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 7×5**Type:** ?**Radio:** Poorly resolved arc.**Optical:** Diffuse shell, brighter to W.**References:**Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.**G68.6–1.2****RA:** 20^h08^m40^s**Dec:** +30°37'**1-GHz flux/Jy:** 1.1**Spectral index:** 0.2**Size/arcmin:** 23**Type:** ?**Radio:** Faint, poorly defined source.**References:**Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz ($\sim 1'$: S=0.57 \pm 0.08 Jy), including review of flux densitiesSun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5: S=0.80 \pm 0.04 Jy) including polarisation and review of flux densities.**G69.0+2.7****RA:** 19^h53^m20^s**Dec:** +32°55'**1-GHz flux/Jy:** 120?**Spectral index:** varies**Size/arcmin:** 80?**Type:** ?

CTB 80

An association with a SN in AD1408 has been suggested. Has been called G68.8+2.8. Is it a SNR?

Radio: Compact core, flat spectrum plateau, and steeper spectrum extensions, with spectral break?**Optical:** Expanding nebulosity near centre, with filaments to the SW and far NE.**X-ray:** Diffuse emission with compact source.**Point sources:** Pulsar at western edge of core.**Distance:** H α observations suggest 1.5 kpc.**References:**Angerhofer *et al.* 1981, A&A, 94, 313. WSRT at 610 MHz (56'' \times 103'') 1.4 GHz (24'' \times 44'') and 5 GHz (7'' \times 13''), plus optical.Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.Sofue *et al.* 1983, PASJ, 35, 437. NRO 45-m at 10.2 GHz (2'7).

Velusamy & Kundu 1983, JApA, 4, 253. VLA of compact sources.

Blair *et al.* 1984, ApJ, 282, 161. Optical images and spectra.

Wang & Seward 1984, ApJ, 285, 607. Einstein observations.

- Strom *et al.* 1984, A&A, 139, 43. Radio observations of flat spectrum component, VLA 5 GHz (1''7) and 1.4 GHz.
- Mantovani *et al.* 1985, A&A, 145, 50. Bologna at 408 MHz (2'6×4'9: $S=67.5\pm 10.5$ Jy), Effelsberg 100-m at 1.41 (9': $S=62\pm 9$ Jy), 1.72 (7'6: $S=66\pm 5$ Jy), 2.7 (4'5: $S=52\pm 4$ Jy) and 4.75 GHz (2'4: $S=44\pm 3.3$ Jy), plus review of flux densities.
- Kulkarni *et al.* 1988, Nature, 331, 50. Pulsar detection.
- Angelini *et al.* 1988, ApJ, 330, L43. EXOSAT spectra.
- Fesen *et al.* 1988, Nature, 334, 229. IRAS of surrounding shell.
- Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.
- Whitehead *et al.* 1989, MNRAS, 237, 1109. Optical of core.
- Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
- Hester & Kulkarni 1989, ApJ, 340, 362. Optical imaging and spectroscopy.
- Koo *et al.* 1990, ApJ, 364, 178. Large, expanding H_i shell.
- Greidanus & Strom 1990, A&A, 240, 376. Optical kinematics of core.
- Koo *et al.* 1993, ApJ, 417, 196. VLA at 1.4 GHz of surrounding H_i (1'), plus IRAS.
- Safi-Harbi *et al.* 1995, ApJ, 439, 722. ROSAT observations.
- Srinivasan 1997, ApJ, 489, 170. γ -ray observations of pulsar.
- Mavromatakis *et al.* 2001, A&A, 371, 300. Optical observations.
- Butler *et al.* 2002, A&A, 395, 845. HST detection of pulsar.
- Migliazzo *et al.* 2002, ApJ, 567, L141. Pulsar proper motion study.
- Castelletti *et al.* 2003, AJ, 126, 2114. GMRT at 240 and 618 MHz (26''×17'' and 10''×6'') and VLA at 324 and 1380 MHz (73''×63'' and 93''×78'').
- Moon *et al.* 2004, ApJ, 610, L33. Chandra and HST observations of core.
- Li *et al.* 2005, ApJ, 628, 931. Chandra observations of pulsar and surroundings.
- Golden *et al.* 2005, ApJ, 635, L153. High resolution radio observations of pulsar and surroundings.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=72\pm 7$ Jy) and 1420 MHz ($\sim 1'$: $S=56\pm 5$ Jy), including polarisation and review of flux densities.
- Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_i.
- Albert *et al.* 2007, ApJ, 669, 1143. γ -ray observations.
- Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion of pulsar.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9'5: $S=36\pm 4$ Jy), including polarisation and review of flux densities.
- Leahy & Ranasinghe 2012, MNRAS, 423, 718. CGPS at 1.4 GHz, including H_i, plus ROSAT observations.
- Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_i in region.

G69.7 + 1.0

RA: 20^h02^m40^s
Dec: +32°43'

1-GHz flux/Jy: 2.0
Spectral index: 0.7

Size/arcmin: 16×14
Type: S

Radio: Poorly resolved source.

X-ray: Detected.

References:

- Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
- Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.
- Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Yoshita *et al.* 2000, PASJ, 52, 867. ROSAT and ASCA observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=3.2\pm 0.4$ Jy) and 1420 MHz ($\sim 1'$: $S=1.5\pm 0.1$ Jy), including review of flux densities.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5: $S=0.78\pm 0.07$ Jy) including polarisation and review of flux densities.

G73.9+0.9

RA: 20^h14^m15^s
Dec: +36°12'

1-GHz flux/Jy: 9
Spectral index: 0.23

Size/arcmin: 27
Type: S?

Radio: Diffuse, centrally brightened to SW.

Optical: Faint shell.

References:

- Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'4 : S = 6.7±0.5 Jy), plus other flux densities.
- Chastenay & Pineault 1988, IAUCom, 101, 297. DRAO at 408 MHz (3'5×5'9) and 1.4 GHz (1'0×1'7).
- Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz (3'4×5'8 : S = 12.7±1.2 Jy) and 1.4 GHz (1'0×1'7 : S = 7.4±1.0 Jy).
- Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
- Pineault *et al.* 1996, AJ, 112, 201. DRAO at 1.4 GHz (smoothed to 2') for H_i.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Mavromatakis 2003, A&A, 398, 153. Optical observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : S = 10.0±1.7 Jy) and 1420 MHz (~1' : S = 7.6±0.6 Jy), including polarisation and review of flux densities.
- Sitnik 2010, ARep, 54, 317. H α and CO observations of region.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 6.2±0.3 Jy) including polarisation and review of flux densities.
- Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G74.0–8.5

Cygnus Loop

RA: 20^h51^m00^s
Dec: +30°40'

1-GHz flux/Jy: 210
Spectral index: varies

Size/arcmin: 230×160
Type: S

Has been suggested that this is two overlapping remnants.

Radio: Shell, brightest to the NE, with fainter breakout region to S, with spectral variations.

Optical: Large filamentary loop, brightest to the NE, not well defined to the S or W.

X-ray: Shell in soft X-rays.

Point sources: Several compact radio sources within the boundary of the remnant, including CL4, plus X-ray sources in S.

Distance: Optical proper motion and shock velocity gives 0.44 kpc.

References:

- Green 1990, AJ, 100, 1927. DRAO at 408 MHz (3'3×6'7) for spectral index study, plus X-ray and optical.
- Graham *et al.* 1991, AJ, 101, 175. Shocked molecular H outside rim in NE.
- Shull & Clarke 1991, PASP, 103, 811. Optical spectroscopy of nearside filaments.
- Fesen *et al.* 1992, AJ, 104, 719. H α imagery.
- Long *et al.* 1992, ApJ, 400, 214. Optical of Balmer dominated filament.
- Arendt *et al.* 1992, ApJ, 400, 562. IRAS observations.
- Hester *et al.* 1994, ApJ, 420, 721. H α , [OIII] and other optical observations of Balmer dominated filaments in NE.
- Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

- Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
- Levenson *et al.* 1997, ApJ, 484, 304. ROSAT observations.
- Leahy *et al.* 1997, AJ, 114, 2081. DRAO at 1.4 GHz ($1' \times 2'$), including polarisation.
- Leahy & Roger 1998, ApJ, 505, 784. DRAO at 1.4 GHz ($1'0 \times 1'9$) and 408 MHz ($3'4 \times 6'9$), for spectral index studies in comparison with other radio observations.
- Levenson *et al.* 1998, ApJS, 118, 541. Optical images.
- Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density ($S = 1400 \pm 400$ Jy).
- Bohigas *et al.* 1999, ApJ, 518, 324. Optical spectroscopy of surroundings.
- Levenson *et al.* 1999, ApJ, 526, 874. ROSAT images.
- Blair *et al.* 1999, AJ, 118, 942. HST observations, for distance.
- Aschenbach & Leahy 1999, A&A, 341, 602. ROSAT image, and comparison with radio.
- Danforth *et al.* 2000, AJ, 119, 2319. UV, optical and X-ray comparison of selected regions.
- Miyata *et al.* 2001, ApJ, 550, 1023. ASCA observations of compact X-ray sources.
- Miyata & Tsunemi 2001, ApJ, 552, 624. ASCA spectroscopy of regions in N and E.
- Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
- Danforth *et al.* 2001, AJ, 122, 938. Far-UV spectroscopy, H α and other optical observations of NE region.
- Levenson & Graham 2001, ApJ, 559, 948. HST of SE region.
- Uyaniker *et al.* 2002, A&A, 389, L61. Effelsberg 100-m at 2.7 GHz ($4'3$) including polarisation, and comparison with ROSAT data.
- Leahy 2002, AJ, 123, 2689. DRAO at 1.4 GHz ($4' \times 2'$) for H α .
- Blair *et al.* 2002, ApJS, 140, 367. UV spectroscopy.
- Welsh *et al.* 2002, A&A, 391, 705. Optical absorption to background stars.
- Sankrit & Blair 2002, ApJ, 565, 297. UV observations of NE filament.
- Levenson *et al.* 2002, ApJ, 576, 798. Chandra observations of W edge.
- Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'5 : S = 184 \pm 18$ Jy).
- Leahy 2004, MNRAS, 351, 385. Chandra observations of SW.
- Uyaniker *et al.* 2004, A&A, 426, 909. Effelsberg 100-m at 2.7 GHz ($4'3$), with comparison with other data for spectral index studies.
- Blair *et al.* 2005, AJ, 129, 2268. HST of outer filaments.
- Leahy 2005, AJ, 130, 165. DRAO at 1.4 GHz of SE.
- Levenson & Graham 2005, ApJ, 622, 366. Chandra observations of knot in SE.
- Sun *et al.* 2006, A&A, 447, 937. Urumqi 25-m at 4.8 GHz ($9'5 : S = 90 \pm 9$ Jy), with comparisons with other data for spectral index studies.
- Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
- Seon *et al.* 2006, ApJ, 644, L175. Far UV observations.
- Sankrit *et al.* 2007, AJ, 133, 1383. UV observations of part.
- Tsunemi *et al.* 2007, ApJ, 671, 1717. XMM observations of NE to SW.
- Miyata *et al.* 2007, PASJ, 59, S163. Suzaku observations of NE.
- Nemes *et al.* 2008, ApJ, 675, 1293. XMM observations of NE.
- Katsuda *et al.* 2008, ApJ, 680, 1198. Chandra observations of NE.
- Uchida *et al.* 2008, ApJ, 688, 1102. XMM observations.
- Uchida *et al.* 2009, PASJ, 61, 503. Suzaku observations of N.
- Kimura *et al.* 2009, PASJ, 61, S137. Suzaku observations from NE to SW.
- Tsunemi *et al.* 2009, PASJ, 61, S147. Suzaku observations of SE.
- Kosugi *et al.* 2010, PASJ, 62, 1035. Suzaku observations of SE.
- Sankrit *et al.* 2010, ApJ, 712, 1092. Spitzer observations.
- McEntaffer & Brantseg 2011, ApJ, 730, 99. Chandra observations of E.
- Katagiri *et al.* 2011, ApJ, 741, 44. Fermi observations.
- Katsuda *et al.* 2012, ApJ, 754, L7. Suzaku and XMM observations.
- Leahy & Hassan 2013, ApJ, 764, 55. Suzaku and XMM observations of SW.
- Oakley *et al.* 2013, ApJ, 766, 51. X-ray spectroscopy.

G74.9 + 1.2

CTB 87

RA: 20^h16^m02^s**1-GHz flux/Jy:** 9**Size/arcmin:** 8×6**Dec:** +37°12′**Spectral index:** varies**Type:** F**Radio:** Filled-centre, with high polarisation and high frequency turnover.**X-ray:** Centrally brightened.**Point sources:** Compact X-ray source in SE.**Distance:** H α absorption indicates 12 kpc, optical extinction gives 6.1 kpc.**References:**

- Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 12.2 \pm 9.7$ Jy, $S_{318\text{ MHz}} = 17.7 \pm 5.0$ Jy.
 Weiler & Shaver 1978, A&A, 70, 389. WSRT at 610 MHz ($57'' \times 94'' : S = 9.1 \pm 1.2$ Jy), 1.4 ($24'' \times 40'' : S = 8.7 \pm 1.2$ Jy) and 5 GHz ($24'' \times 40'' : S = 5.6 \pm 1.3$ Jy).
 Geldzahler *et al.* 1980, A&A, 84, 237. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 7.6 \pm 0.5$ Jy).
 Wilson 1980, ApJ, 241, L19. Einstein observations.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H α absorption of nearby compact source.
 Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (smoothed to $40'' : S = 1.47 \pm 0.19$ Jy).
 Green & Gull 1989, MNRAS, 237, 555. VLA at 1.4 GHz ($1'.2 \times 1'.4$) including H α .
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz, plus review of flux densities.
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ($3'.4 \times 5'.8 : S = 11.6 \pm 0.4$ Jy) and 1.4 GHz ($1'.0 \times 1'.7 : S = 7.2 \pm 0.3$ Jy).
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz ($3'.5 \times 5'.2 : S = 13.3 \pm 0.8$ Jy) and Effelsberg 100-m at 4.8 GHz ($S = 7.5 \pm 0.7$ Jy).
 Wallace *et al.* 1994, A&A, 286, 565. H α of surroundings.
 Cho *et al.* 1994, AJ, 108, 634. CO of adjacent molecular clouds.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Wallace *et al.* 1997, A&A, 317, 212. DRAO at 408 MHz ($3'.4 \times 5'.5$) and 1.4 GHz ($1'.0 \times 1'.6$) including H α (smoothed to $2'$).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2003, ApJ, 588, 852. CGPS at 1.4 GHz ($1'.6 \times 1'$) including H α , plus CO observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 11.9 \pm 0.9$ Jy) and 1420 MHz ($\sim 1' : S = 7.1 \pm 1.1$ Jy), including polarisation and review of flux densities.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Sitnik 2010, ARep, 54, 317. H α and CO observations of region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 6.4 \pm 0.4$ Jy) including polarisation and review of flux densities.
 Matheson *et al.* 2013, ApJ, 774, 33. Chandra observations.

G76.9 + 1.0**RA:** 20^h22^m20^s**1-GHz flux/Jy:** 2?**Size/arcmin:** 9**Dec:** +38°43′**Spectral index:** ?**Type:** C**Radio:** Bipolar shell.**Point sources:** Central pulsar.**References:**

Landecker *et al.* 1993, A&A, 276, 522. VLA at 1.49 GHz (14''), 4.86 GHz (13'' \times 16'') and 8.55 GHz (11'' \times 12''), including polarisation and review of flux densities.
 Landecker *et al.* 1997, A&AS, 123, 199. Miyun at 232-MHz (3'8 \times 5'4).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (\sim 3':S=2.3 \pm 0.2 Jy) and 1420 MHz (\sim 1':S=1.35 \pm 0.07 Jy), including polarisation and review of flux densities.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5':S=0.79 \pm 0.07 Jy) including polarisation and review of flux densities.
 Marthi *et al.* 2011, MNRAS, 416, 2560. GMRT at 618 MHz (51'' \times 54''), 1160 MHz (2''2 \times 3''4), and Chandra observations of central source.
 Arzoumanian *et al.* 2011, ApJ, 739, 39. Pulsar detection.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G78.2 + 2.1
DR4, γ Cygni SNR**RA:** 20^h20^m50^s**1-GHz flux/Jy:** 320**Size/arcmin:** 60**Dec:** +40 $^{\circ}$ 26'**Spectral index:** 0.51**Type:** S

Has been called G78.1+1.8.

Radio: In complex region (early catalogues refer to other proposed remnants in this region).

Optical: Faint filaments, spectra indicate a SNR superposed on a H α region.

X-ray: Weak emission from the SE of the remnant.

Point sources: X-ray pulsar at edge of remnant.

Distance: Associations with other objects suggests 1.7 to 2.6 kpc.

References:

Higgs *et al.* 1977, AJ, 82, 718. DRAO at 1.4 GHz (2' \times 3':S=270 \pm 40 Jy) plus some 10 GHz (4') survey data, reveals true extent of remnant.
 d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.
 van den Bergh 1978, ApJS, 38, 119. Optical observations.
 Landecker *et al.* 1980, A&AS, 39, 133. DRAO H α observations (2' \times 3'1).
 Higgs *et al.* 1983, AJ, 88, 97. CO of surroundings.
 Bohigas *et al.* 1983, RMxAA, 8, 155. Optical spectra, find thermal only.
 Braun & Strom 1986, A&AS, 63, 345. WSRT H α observations.
 Fukui & Tatematsu 1988, IAUCom, 101, 261. CO observations of the vicinity (2'7).
 Green 1989, MNRAS, 238, 737. OH observations.
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz (3'4 \times 5'8':S=480 \pm 60 Jy) and 1.4 GHz (1'0 \times 1'7':S=270 \pm 40 Jy).
 Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz (3'5 \times 5'2':S=540 \pm 40 Jy) and Effelsberg 100-m at 4.8 GHz (S=150 \pm 15 Jy).
 Esposito *et al.* 1996, ApJ, 461, 820. Associated γ -ray emission.
 Brazier *et al.* 1996, MNRAS, 281, 1033. γ -ray and X-ray point source.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Zhang *et al.* 1997, A&A, 324, 641. Multi-frequency radio comparison.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
 Uchiyama *et al.* 2002, ApJ, 571, 866. ASCA observations.
 Mavromatakis 2003, A&A, 408, 237. Optical observations.
 Bykov 2004, A&A, 427, L21. Hard X-ray observations.
 Becker *et al.* 2004, ApJ, 615, 897. Chandra and other observations of compact sources.
 Weisskopf *et al.* 2006, ApJ, 652, 387. Chandra and other observations of compact sources.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=500\pm 35$ Jy) and 1420 MHz ($\sim 1'$: $S=226\pm 19$ Jy), including review of flux densities.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_i.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Ladouceur & Pineault 2008, A&A, 490, 197. CGPS at 408 MHz ($2'.9\times 4'.5$) and 1.4 GHz ($0'.8\times 1'.5$).
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'.5$: $S=170\pm 18$ Jy), including polarisation and review of flux densities.
 Leahy *et al.* 2013, MNRAS, 436, 968. ROSAT and Chandra observations, and CGPS for H_i.
 Aliu *et al.* 2013, ApJ, 770, 93. γ -ray observations.
 Lin *et al.* 2013, ApJ, 770, L9. Pulsar detection.

G82.2 + 5.3

W63

RA: 20^h19^m00^s**1-GHz flux/Jy:** 120?**Size/arcmin:** 95×65**Dec:** +45°30'**Spectral index:** 0.5?**Type:** S

Has been called G82.5+5.3.

Radio: Shell in the Cygnus X complex.

Optical: In complex region, but spectra indicate SNR filaments.

X-ray: Detected.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5'$: $S=59.0\pm 3.5$ Jy).
 Sabbadin 1976, A&A, 51, 159. Optical spectra.
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'.8$). Incomplete mapping.
 Rosado & González 1981, RMxAA, 5, 93. Optical spectra.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'.5$: $S=82.5\pm 5.5$ Jy).
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation, of part.
 Mavromatakis *et al.* 2004, A&A, 415, 1051. ROSAT, ASCA and optical observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=144\pm 12$ Jy) and 1420 MHz ($\sim 1'$: $S=93\pm 5$ Jy), including review of flux densities.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'.5$: $S=49\pm 5$ Jy), including polarisation and review of flux densities.

G83.0 – 0.3

RA: 20^h46^m55^s**1-GHz flux/Jy:** 1**Size/arcmin:** 9×7**Dec:** +42°52'**Spectral index:** 0.4**Type:** S

Radio: Incomplete shell.

References:

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=1.2\pm 0.3$ Jy) and 1420 MHz ($\sim 1'$: $S=0.8\pm 0.1$ Jy), including polarisation and review of flux densities.

G84.2 – 0.8**RA:** 20^h53^m20^s**Dec:** +43°27′**1-GHz flux/Jy:** 11**Spectral index:** 0.5**Size/arcmin:** 20×16**Type:** S**Radio:** Elongated shell, with a filament aligned with the major axis.**X-ray:** Detected.**Distance:** H_I absorption suggests 6 kpc.**References:**

Matthews *et al.* 1977, A&A, 55, 1. WSRT at 610 MHz (56″×81″: $S = 12.4 \pm 1.5$ Jy) and Effelsberg 100-m at 2.7 GHz (4′4″: $S = 6.8 \pm 1.3$ Jy).

Matthews & Shaver 1980, A&A, 87, 255. WSRT at 1415 MHz (23″×32″), and Effelsberg 100-m at 2.7 GHz (4′4″: $S = 5.6 \pm 0.5$ Jy).

Feldt & Green 1993, A&A, 274, 421. DRAO at 1.4 GHz (1′×1′5), including H_I, plus CO observations.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1′) including polarisation.

Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3′$: $S = 14.5 \pm 0.5$ Jy) and 1420 MHz ($\sim 1′$: $S = 7.2 \pm 0.8$ Jy), including review of flux densities.

Leahy & Green 2012, ApJ, 760, 25. CGPS, including H_I, plus Chandra observations.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G85.4 + 0.7**RA:** 20^h50^m40^s**Dec:** +45°22′**1-GHz flux/Jy:** ?**Spectral index:** 0.2**Size/arcmin:** 24?**Type:** S**Radio:** Faint, incomplete shell, within larger thermal shell.**X-ray:** Centrally brightened.**Distance:** H_I observations suggest 3.5 kpc.**References:**

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz (4′4″×2′8″: $S < 0.45$ Jy) and 1.4 GHz (1′1″×0′8″), plus H_I, X-ray and optical data.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3′$: $S = 2.9 \pm 0.5$ Jy) and 1420 MHz ($\sim 1′$: $S = 2.3 \pm 0.2$ Jy), including review of flux densities.

Jackson *et al.* 2008, ApJ, 674, 936. XMM and H_I observations.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G85.9 – 0.6**RA:** 20^h58^m40^s**Dec:** +44°53′**1-GHz flux/Jy:** ?**Spectral index:** 0.2**Size/arcmin:** 24**Type:** S**Radio:** Faint, incomplete shell.**Optical:** Diffuse shell.**X-ray:** Centrally brightened.**Distance:** H_I observations suggest 4.8 kpc.**References:**

- Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($4'.4 \times 2'.8 : S < 0.9$ Jy) and 1.4 GHz ($1'.1 \times 0'.8$), plus H α , X-ray and optical data.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.0 \pm 1.3$ Jy) and 1420 MHz ($\sim 1' : S = 2.2 \pm 0.8$ Jy), including review of flux densities.
- Jackson *et al.* 2008, ApJ, 674, 936. XMM and H α observations.
- Gök *et al.* 2009, Ap&SS, 324, 17. Optical observations.
- Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G89.0 + 4.7

HB21

RA: 20^h45^m00^s**1-GHz flux/Jy:** 220**Size/arcmin:** 120×90**Dec:** +50°35'**Spectral index:** 0.38**Type:** S

Radio: Distorted shell (4C50.52, an extragalactic double, is within the boundary of the remnant).

Optical: Filaments and patches.

X-ray: Centrally brightened.

Distance: Various associations imply 0.8 kpc.

References:

- Hirabayashi & Takahashi 1972, PASJ, 24, 231. 30-m dish at 4.2 GHz ($11' : S = 160 \pm 40$ Jy).
- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 148 \pm 16$ Jy), plus optical filaments.
- Hill 1974, MNRAS, 169, 59. Half-Mile Telescope at 1.4 GHz ($3' \times 3'.9$).
- Haslam *et al.* 1975, A&A, 39, 453. Effelsberg 100-m at 2.7 GHz ($4'.4$).
- Fukui & Tatematsu 1988, IAUCo, 101, 261. CO observations of the vicinity ($2'.7$).
- Tatematsu *et al.* 1990, A&A, 237, 189. DRAO at 408 MHz ($3'.5 \times 4'.5$) and 1.4 GHz ($1'.0 \times 1'.3$), including H α , plus CO observations of adjacent molecular cloud.
- Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
- Leahy & Aschenbach 1996, A&A, 315, 260. ROSAT observations.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Koo *et al.* 2001, ApJ, 552, 175. NRAO 12-m and other CO observations ($27''$ and $45''$) of eastern part.
- Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'.5 : S = 228 \pm 5$ Jy).
- Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation.
- Byun *et al.* 2006, ApJ, 637, 283. CO observations of surroundings.
- Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
- Leahy 2006, ApJ, 647, 1125. CGPS at 408 MHz ($2'.8 \times 3'.7$) and 1.4 GHz ($0'.8 \times 1'.1$).
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 259 \pm 19$ Jy) and 1420 MHz ($\sim 1' : S = 183 \pm 9$ Jy), including polarisation and review of flux densities.
- Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H α .
- Mavromatakis *et al.* 2007, A&A, 461, 991. Optical observations.
- Shinn *et al.* 2009, ApJ, 693, 1883. IR observations.
- Pannuti *et al.* 2010, AJ, 140, 1787. ASCA and Chandra observations.
- Shinn *et al.* 2010, AdSpR, 45, 445. IR observations in S.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'.5 : S = 107 \pm 11$ Jy), including polarisation and review of flux densities.
- Reichardt *et al.* 2012, A&A, 546, A21. γ -ray detection.
- Shinn *et al.* 2012, ApJ, 759, 34. AKARI observations of S.
- Pivato *et al.* 2013, ApJ, 779, 179. Fermi observations.

G93.3 + 6.9

DA 530, 4C(T)55.38.1

RA: 20^h52^m25^s**1-GHz flux/Jy:** 9**Size/arcmin:** 27×20**Dec:** +55°21′**Spectral index:** 0.45**Type:** C?

Has been called G93.2+6.7.

Radio: Shell, with two bright limbs, highly polarised.

X-ray: Compact central source.

Distance: H_I observations suggest 2.2 kpc.

References:

Roger & Costain 1976, A&A, 51, 151. DRAO at 1.42 GHz (2′×2′.4 : S=6.9 Jy).

Haslam *et al.* 1980, A&A, 92, 57. Effelsberg 100-m at 1.72 GHz (7′.6 : S=6.47±0.52 Jy) and 2.7 GHz (4′.4 : S=5.64±0.64 Jy), plus review of flux densities.

Lalitha *et al.* 1984, A&A, 131, 196. Effelsberg 100-m at 4.75 GHz (smoothed to 3′ : S=4.01±0.57 Jy).

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Landecker *et al.* 1999, ApJ, 527, 866. DRAO at 408 MHz (3′.5×4′.3) and 1.4 GHz (1′.0×1′.2), including polarisation and H_I.

Foster & Routledge 2003, ApJ, 598, 1005. H_I for distance.

Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′ : S=10.5±0.7 Jy) and 1420 MHz, including review of flux densities.

Jiang *et al.* 2007, ApJ, 670, 1142. Chandra observations.

Bocchino *et al.* 2008, AdSpR, 41, 407. XMM observations.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G93.7 – 0.2

CTB 104A, DA 551

RA: 21^h29^m20^s**1-GHz flux/Jy:** 65**Size/arcmin:** 80**Dec:** +50°50′**Spectral index:** 0.65**Type:** S

Has been called G93.6–0.2 and G93.7–0.3.

Radio: Distorted, faint shell.

Distance: Association with H_I features suggests 1.5 kpc.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′ : S=18.4±1.0 Jy).

Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz (7′.6 : S=53.5±5.0 Jy), plus review of flux densities.

Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to 2′ : S=58±6 Jy).

Mantovani *et al.* 1991, A&A, 247, 545. Effelsberg 100-m at 4.75 GHz (smoothed to 3′ : S=33.5±4.0 Jy), including polarisation, plus review of flux densities.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

Uyaniker *et al.* 2002, ApJ, 565, 1022. CGPS 1.4 GHz (54″×49″), including H_I, and 408 MHz (3′.7×2′.8).

Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1′) including polarisation.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′ : S=67±6 Jy) and 1420 MHz (~1′ : S=35±4 Jy), including polarisation and review of flux densities.

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S=25.0±2.5 Jy), including polarisation and review of flux densities.

G94.0 + 1.0

3C434.1

RA: 21^h24^m50^s
Dec: +51°53'**1-GHz flux/Jy:** 13
Spectral index: 0.45**Size/arcmin:** 30×25
Type: S**Radio:** Incomplete shell, containing H_I shell.**Distance:** Association with stellar wind bubble implies 5.2 kpc.**References:**

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 6.1 \pm 0.8$ Jy), and 37-m at 1.7 GHz ($S = 11 \pm 3$ Jy).
- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 5.8 \pm 0.4$ Jy). Also NRAO 140-ft at 5 GHz (6').
- Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ($7'.6 : S = 12.0 \pm 1.3$ Jy), plus review of flux densities.
- Goss *et al.* 1984, A&A, 138, 469. WSRT at 610 MHz (smoothed to $100'' : S = 16 \pm 1.7$ Jy) and Effelsberg 100-m at 4.75 GHz ($2'.4 : S = 7.2 \pm 0.5$ Jy).
- Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 16 \pm 3$ Jy).
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1') including polarisation.
- Foster *et al.* 2004, A&A, 417, 79. DRAO at 1.4 GHz, including H_I.
- Foster 2005, A&A, 441, 1043. CGPS at 408 MHz ($2'.8 \times 3'.6$) and 1.4 GHz ($0'.8 \times 1'.0$) for spectral index studies, plus other observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 20 \pm 2$ Jy) and 1420 MHz ($\sim 1' : S = 11.3 \pm 1.0$ Jy), including review of flux densities.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 6.2 \pm 0.4$ Jy) including polarisation and review of flux densities.
- Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
- Jeong *et al.* 2013, ApJ, 770, 105. CO observations of region.

G96.0 + 2.0**RA:** 21^h30^m30^s
Dec: +53°59'**1-GHz flux/Jy:** 0.35
Spectral index: 0.6**Size/arcmin:** 26
Type: S**Radio:** Faint, arc in S, poorly defined in N.**Distance:** Association for H_I indicates 4 kpc.**References:**

- Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz ($2'.8 \times 3'.5$) and 1.4 GHz ($50'' \times 61''$) including H_I.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 0.42 \pm 0.06$ Jy) and 1420 MHz ($\sim 1' : S = 0.24 \pm 0.02$ Jy), including review of flux densities.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 0.14 \pm 0.02$ Jy) including polarisation and review of flux densities.

G106.3+2.7**RA:** 22^h27^m30^s**Dec:** +60°50′**1-GHz flux/Jy:** 6**Spectral index:** 0.6**Size/arcmin:** 60×24**Type:** C?

Incorporates the pulsar wind nebula G106.6+2.9 (the ‘Boomerang’).

Radio: Faint extended source, which brighter ‘head’ to NE.

X-ray: Pulsar and wind nebula.

Point sources: Pulsar.

References:

Pineault & Joncas 2000, *AJ*, 120, 3218. DRAO at 408 MHz (3′.9×3′.5: $S=10.5\pm0.3$ Jy) and 1.4 GHz (1′.2×1′.0: $S=4.9\pm0.6$ Jy), plus H_i.

Halpern *et al.* 2001, *ApJ*, 547, 323. X-ray and radio observations of the ‘head’.

Halpern *et al.* 2001, *ApJ*, 552, L125. Pulsar detection.

Kothes *et al.* 2001, *ApJ*, 560, 236. CGPS at 1.4 GHz, including H_i, plus CO and other observations.

Kothes *et al.* 2004, *ApJ*, 607, 855. H_i polarisation absorption.

Ng & Romani 2004, *ApJ*, 601, 479. Chandra detection of pulsar wind nebula.

Kothes *et al.* 2006, *ApJ*, 638, 225. Effelsberg 100-m at 4.85 (2′.4), 8.35 (1′.4), 10.5 (1′.2) and 32 GHz (0′.45) of pulsar wind nebula, including polarisation.

Kothes *et al.* 2006, *A&A*, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=8.6\pm1.0$ Jy) and 1420 MHz ($\sim 1'$: $S=4.8\pm0.5$ Jy), including polarisation and review of flux densities.

Abdo *et al.* 2007, *ApJ*, 664, L91. γ -ray observations.

Casandjian & Grenier 2008, *A&A*, 489, 849. γ -ray observations.

Acciari *et al.* 2009, *ApJ*, 703, L6. γ -ray observations.

Gao *et al.* 2011, *A&A*, 529, A159. Urumqi 25-m at 5 GHz (9′.5: $S=2.0\pm0.3$ Jy), including polarisation and review of flux densities.

G108.2–0.6**RA:** 22^h53^m40^s**Dec:** +58°50′**1-GHz flux/Jy:** 8**Spectral index:** 0.5**Size/arcmin:** 70×54**Type:** S

Radio: Faint shell.

Distance: Possible associated H_i structures suggest 3.2 kpc.

References:

Tian *et al.* 2007, *A&A*, 465, 907. DRAO at 408 MHz (2′.8×3′.3: $S=11.5\pm1.2$ Jy) and 1.4 GHz (1′.0×1′.2: $S=6.6\pm0.7$ Jy) including H_i.

G109.1–1.0

CTB 109

RA: 23^h01^m35^s**1-GHz flux/Jy:** 22**Size/arcmin:** 28**Dec:** +58°53′**Spectral index:** 0.45**Type:** S

Radio: Semicircular shell, with the Molecular cloud S152 is to the immediate W.

X-ray: Semicircular shell, with pulsar at W edge.

Point sources: Long period X-ray pulsar.

Distance: Various observations imply 3.2 kpc.

References:

- Hughes *et al.* 1981, ApJ, 246, L127. WSRT at 610 MHz ($1': S=40\pm 5$ Jy) shows bad CLEAN artefacts.
- Blair & Kirshner 1981, Nature, 291, 132. Optical spectra.
- Downes 1983, MNRAS, 203, 695. Effelsberg 100-m at 2.7 GHz ($4'.4: S=13.0\pm 1.5$ Jy).
- Sofue *et al.* 1983, PASJ, 35, 447. NRO 45-m at 10.2 GHz ($2'.7$).
- Hughes *et al.* 1984, ApJ, 283, 147. WSRT at 610 MHz ($20'': S=26\pm 3$ Jy) 1.4 GHz ($10''$), DRAO at 1.4 GHz ($1'\times 1'.2: S=16.8\pm 2$ Jy) and Algonquin 46-m at 6.5 GHz ($4'.5: S=6.7\pm 1$ Jy).
- Braun & Strom 1986, A&AS, 63, 345. WSRT H_I observations.
- Tatematsu *et al.* 1987, PASJ, 39, 755. NRO 45-m at 10 GHz ($2'.7$), plus polarisation.
- Tatematsu *et al.* 1987, A&A, 184, 279. CO observations of the surroundings ($2'.7$).
- Hanson *et al.* 1988, A&A, 195, 114. EXOSAT of pulsar.
see Hanson *et al.* 1988, A&A, 207, 204. Erratum.
- Morini *et al.* 1988, ApJ, 333, 777. EXOSAT observations.
- Koyama *et al.* 1989, PASJ, 41, 461. X-ray observations of pulsar.
- Coe *et al.* 1989, MNRAS, 238, 649. IRAS observations of surroundings.
- Green 1989, MNRAS, 238, 737. OH observations.
- Tatematsu *et al.* 1990, ApJ, 351, 157. CO of surroundings, plus X-ray observations.
- Davies & Coe 1991, MNRAS, 249, 313. Optical and IR observations near pulsar.
- Hurford & Fesen 1995, MNRAS, 277, 549. ROSAT imaging.
- Rho & Petre 1997, ApJ, 484, 828. ROSAT observations.
- Parmar *et al.* 1998, A&A, 330, 175. X-ray observations.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
- Patel *et al.* 2001, ApJ, 563, L45. Chandra observations of pulsar.
- Kothes *et al.* 2002, ApJ, 576, 169. CGPS at 1.4 GHz ($68''\times 59''$), including H_I, plus CO observations.
- Sasaki *et al.* 2004, ApJ, 617, 322. XMM observations.
- Sasaki *et al.* 2006, ApJ, 642, L149. CO observations of surroundings, plus Chandra observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3': S=26\pm 3$ Jy) and 1420 MHz ($\sim 1': S=17.4\pm 1.2$ Jy), including polarisation and review of flux densities.
- Tian *et al.* 2010, MNRAS, 404, L1. CGPS for H_I absorption.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5: S=9.8\pm 0.5$ Jy) including polarisation and review of flux densities.
- Castro *et al.* 2012, ApJ, 756, 88. Fermi observations.
- Kothes & Foster 2012, ApJ, 746, L4. H_I and CO observations of region.
- Sasaki *et al.* 2013, A&A, 552, A45. Chandra observations of NE.
- Tendulakar *et al.* 2013, ApJ, 772, 31. Pulsar proper motion study.

G111.7–2.1

Cassiopeia A, 3C461

RA: 23^h23^m26^s**1-GHz flux/Jy:** 2720**Size/arcmin:** 5**Dec:** +58°48′**Spectral index:** 0.77**Type:** S

Presumably the remnant of a late 17th century SN.

Radio: Bright shell with compact knots and extended plateau of emission.

Optical: Fast knots and quasi-stationary flocculi, with many filaments at large radii, and NE ‘jet’.

X-ray: Incomplete shell, with hard spectral component.

Point sources: Central compact X-ray source.

Distance: Optical expansion, plus proper motions indicate 3.4 kpc.

References:

- Anderson *et al.* 1991, ApJ, 373, 146. VLA at 1.4 and 5 GHz (1'') for spectral variations.
- Anderson *et al.* 1995, ApJ, 441, 300. VLA at 1.4 GHz and 4.8 GHz (both 1''), for polarisation studies.
- Anderson & Rudnick 1995, ApJ, 441, 307. VLA at 1.4 GHz and 4.8 GHz, for proper motion studies.
- Kassim *et al.* 1995, ApJ, 455, L59. VLA at 74 (25'') and 332 MHz (6''), including spectral index comparisons.
- Anderson & Rudnick 1996, ApJ, 456, 234. VLA for spectral index comparisons.
- Schwarz *et al.* 1997, A&AS, 123, 43. WSRT at 1.4 GHz (30'') for H_I absorption.
- Reynoso *et al.* 1997, A&A, 317, 203. VLA at 1.4 GHz (11''×14'') for H_I absorption.
- O'Sullivan & Green 1999, MNRAS, 303, 575. Flux density changes at 13.5, 15.5 and 16.5 GHz.
- Agüeros & Green 1999, MNRAS, 305, 957. 151 MHz observations for bulk expansion studies.
- Mason *et al.* 1999, AJ, 118, 2908. Flux density at 32 GHz ($S=194\pm 5$ Jy).
- Reichart & Stephens 2000, ApJ, 537, 904. Flux density changes at 1405 MHz.
- Gotthelf *et al.* 2001, ApJ, 552, L39. Chandra observations, showing outer shock.
- Bleeker *et al.* 2001, A&A, 365, L225. XMM observations.
- Maccarone *et al.* 2001, A&A, 368, 267. BeppoSAX observations.
- Douvion *et al.* 2001, A&A, 369, 589. ISO observations.
- Thorstensen *et al.* 2001, AJ, 122, 297. Optical expansion studies.
- Hwang *et al.* 2001, ApJ, 560, L175. Chandra observations of Doppler shifted lines.
- Fesen *et al.* 2001, AJ, 122, 2644. HST observations.
- Willingale *et al.* 2002, A&A, 381, 1039. XMM observations.
- Mereghetti *et al.* 2002, ApJ, 569, 275. XMM observations of central source.
- Reynoso & Goss 2002, ApJ, 575, 871. VLA at 5 GHz (6''4×6''0) for H₂CO absorption studies.
- Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz, including polarisation.
- Rho *et al.* 2003, ApJ, 592, 299. IR observations.
- Laming & Hwang 2003, ApJ, 597, 347. Chandra spectroscopy.
- Hines *et al.* 2004, ApJS, 154, 290. Spitzer observations.
- Morse *et al.* 2004, ApJ, 614, 727. HST proper motion studies.
- Hwang *et al.* 2004, ApJ, 615, L117. Deep Chandra imaging.
- Krause *et al.* 2004, Nature, 432, 596. IR and molecular line observations of surroundings.
- Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
- Wilson & Batrla 2005, A&A, 430, 561. Observational constraints on dust.
- Krause *et al.* 2005, Science, 308, 1604. Spitzer light echoes from surroundings.
- Fesen *et al.* 2006, ApJ, 636, 848. Optical and IR limits for central sources.
- Vinyūkin 2006, ARep, 50, 143. Secular decrease at 36, 81.5 and 151.5 MHz.
- Laming *et al.* 2006, ApJ, 644, 260. Chandra observations, particularly of NE ‘jet’.
- Fesen *et al.* 2006, ApJ, 645, 283. HST observations for expansion studies.
- Renaud *et al.* 2006, ApJ, 647, L41. γ -ray line observations.
- Lazendic *et al.* 2006, ApJ, 651, 250. Chandra spectroscopy of bright knots.
- Ennis *et al.* 2006, ApJ, 652, 376. Spitzer observations.

Mookerjee *et al.* 2006, MNRAS, 371, 761. Sub-mm [C_I] observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_i.
 Patnaude & Fesen 2007, AJ, 133, 147. X-ray variability of compact features.
 Albert *et al.* 2007, A&A, 474, 937. γ -ray observations.
 Rho *et al.* 2008, ApJ, 673, 271. Spitzer observations.
 Uchiyama & Aharonian 2008, ApJ, 677, L105. X-ray variability of compact features.
 Rest *et al.* 2008, ApJ, 681, L81. SN light echo.
 Hammell & Fesen 2008, ApJS, 179, 195. HST of outer knots.
 Krause *et al.* 2008, Science, 320, 1195. SN light echo spectrum.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Helmboldt & Kassim 2009, AJ, 138, 838. Low radio frequency temporal variations.
 Maeda *et al.* 2009, PASJ, 61, 1217. Suzaku observations.
 Smith *et al.* 2009, ApJ, 693, 713. Spitzer observations.
 Patnaude & Fesen 2009, ApJ, 697, 535. X-ray proper motions.
 Pavlov & Luna 2009, ApJ, 703, 910. Chandra observations of central source.
 Barlow *et al.* 2010, A&A, 518, L138. Herschel observations.
 Abdo *et al.* 2010, ApJ, 710, L92. Fermi observations.
 Acciari *et al.* 2010, ApJ, 714, 163. γ -ray observations.
 Sibthorpe *et al.* 2010, ApJ, 719, 1553. AKARI and sub-mm observations of region.
 DeLaney *et al.* 2010, ApJ, 725, 2038. Spitzer and Chandra observations for 3-D structure.
 Patnaude *et al.* 2011, ApJ, 729, L28. Chandra observations of fading.
 Rest *et al.* 2011, ApJ, 732, 3. SN light echo observations.
 Fesen *et al.* 2011, ApJ, 736, 109. HST variability studies.
 Besel & Krause 2012, A&A, 541, L3. IR light echoes.
 Hwang & Laming 2012, ApJ, 746, 130. Chandra observations.
 Rho *et al.* 2012, ApJ, 747, L6. AKARI CO observations.
 Vogt *et al.* 2012, ApJ, 750, 155. Spitzer light echoes.
 Isenne *et al.* 2012, ApJ, 757, 126. Spitzer spectroscopy.
 Asgekar *et al.* 2013, A&A, 551, L11. LOFAR of carbon recombination lines.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
 Rutherford *et al.* 2013, ApJ, 769, 64. Chandra spectroscopy.
 Milisavljevic & Fesen 2013, ApJ, 772, 134. Optical spectroscopy for 3-D structure.
 Koo *et al.* 2013, Science, 342, 1346. IR spectroscopy.
 Willström *et al.* 2013, A&A, 558, L2. CO observations.

G113.0+0.2

RA: 23^h36^m35^s

Dec: +61°22'

1-GHz flux/Jy: 4

Spectral index: 0.5?

Size/arcmin: 40×17?

Type: ?

Radio: Elongated, extent not well defined.

Point sources: Contains old pulsar.

Distance: Association for H_i indicates 3.1 kpc.

References:

Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz (2'8×3'1) and 1.4 GHz (49''×55'') including H_i.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3') and 1420 MHz (~1').
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 1.9±0.5 Jy) including polarisation and review of flux densities.

G114.3+0.3**RA:** 23^h37^m00^s**Dec:** +61°55′**1-GHz flux/Jy:** 5.5**Spectral index:** 0.5**Size/arcmin:** 90×55**Type:** S**Radio:** Shell, with H_{II} region S165 within the boundary of the remnant.**Optical:** Faint emission in centre and to S.**Point sources:** Pulsar near centre of remnant.**Distance:** Association with H_I and other features implies 0.7 kpc.**References:**

- Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4′.4 : S = 3.6 Jy) and S_{1.4 GHz} = 4.4 Jy from 1.4 GHz survey data, plus H_I from Maryland--Green Bank survey.
- Kulkarni *et al.* 1993, Nature, 362, 135. Pulsar association.
- Fürst *et al.* 1993, A&A, 276, 470. Pulsar association.
- Becker *et al.* 1996, A&A, 306, 464. ROSAT of pulsar.
- Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
- Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
- Mavromatakis *et al.* 2002, A&A, 383, 1011. Optical observations.
- Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz (49″×55″), including H_I (1′.0×1′.1).
- Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz (3′.4×3′.9 : S = 12.0±6.0 Jy) and 1.4 GHz (1′.0×1′.1 : S = 9.8±0.8 Jy).
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz (~1′ : S = 5.4±0.8 Jy), including review of flux densities.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 6.9±0.7 Jy), including polarisation and review of flux densities.

G116.5+1.1**RA:** 23^h53^m40^s**Dec:** +63°15′**1-GHz flux/Jy:** 10**Spectral index:** 0.5**Size/arcmin:** 80×60**Type:** S**Radio:** Distinct shell, with high polarisation.**Optical:** Detected.**Distance:** Association with H_I features implies 1.6 kpc.**References:**

- Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4′.4 : S = 4.7±0.4 Jy) and S_{1.4 GHz} = 8.0±0.8 Jy from 1.4 GHz survey data, plus H_I from Maryland--Green Bank survey.
- Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz (49″×55″), including H_I (1′.0×1′.1).
- Mavromatakis *et al.* 2005, A&A, 435, 141. Optical observations.
- Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz (3′.4×3′.8 : S = 15.0±1.5 Jy) and 1.4 GHz (1′.0×1′.1 : S = 10.6±0. Jy).
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′ : S = 12.5±1.6 Jy) and 1420 MHz (~1′ : S = 10.3±0.70 Jy), including polarisation and review of flux densities.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 5.7±0.6 Jy), including polarisation and review of flux densities.

G116.9+0.2

CTB 1

RA: 23^h59^m10^s**1-GHz flux/Jy:** 8**Size/arcmin:** 34**Dec:** +62°26′**Spectral index:** 0.57**Type:** S

Has been called G117.3+0.1 and G116.9+0.1.

Radio: Incomplete shell.

Optical: Filaments on sky survey.

X-ray: Centrally brightened, with NE ‘breakout’.

Point sources: Pulsar to NE.

Distance: Association with H_I features implies 1.6 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 3.9 \pm 1.0$ Jy), and 37-m at 1.7 GHz ($S = 5.5 \pm 2.0$ Jy), plus review of flux densities.

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 4.2 \pm 0.2$ Jy).

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'8 : S = 3.0 \pm 0.3$ Jy).

Dickel & Willis 1980, A&A, 85, 55. WSRT at 610 MHz ($56'' \times 64''$) and 1.4 GHz ($22'' \times 25''$).

Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz ($4'4 : S = 4.8 \pm 0.4$ Jy) and $S_{1.4 \text{ GHz}} = 7.8 \pm 0.8$ Jy from 1.4 GHz survey data, plus H_I from Maryland--Green Bank survey.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Landecker *et al.* 1982, AJ, 87, 1379. DRAO at 1.42 GHz and H_I ($2' \times 2'3 : S = 8.3 \pm 0.5$ Jy), plus review of flux densities.

see also: Landecker *et al.* 1983, AJ, 88, 877. Erratum.

Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.

Hailey & Craig 1994, ApJ, 434, 635. Optical spectroscopy.

Hailey & Craig 1995, ApJ, 455, L151. ROSAT of nearby pulsar.

Fesen *et al.* 1997, AJ, 113, 767. Optical observations.

Craig *et al.* 1997, ApJ, 488, 307. ROSAT observations.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz ($69''$), including polarisation.

Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz ($49'' \times 55''$), including H_I ($1'0 \times 1'1$).

Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.

Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz ($3'4 \times 3'8 : S = 15.0 \pm 1.5$ Jy) and 1.4 GHz ($1'0 \times 1'1 : S = 8.1 \pm 0.4$ Jy).

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.5 \pm 0.8$ Jy) and 1420 MHz ($\sim 1' : S = 7.0 \pm 0.8$ Jy), including review of flux densities.

Pannuti *et al.* 2010, AJ, 140, 1787. ASCA and Chandra observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5 : S = 3.6 \pm 0.4$ Jy) including polarisation and review of flux densities.

G119.5+10.2

CTA 1

RA: 00^h06^m40^s**1-GHz flux/Jy:** 36**Size/arcmin:** 90?**Dec:** +72°45'**Spectral index:** 0.6**Type:** S

Has been called G119.5+10.3.

Radio: Incomplete shell, with 'breakout' to NW.

Optical: Faint diffuse nebulosities.

X-ray: Centrally brightened.

Point sources: Central pulsar.

Distance: Associated H_i shell indicates 1.4 kpc.

References:

- Sieber *et al.* 1979, A&A, 74, 361. Effelsberg 100-m at 2.7 GHz (4'4": $S=23.6\pm 2.8$ Jy).
 Sieber *et al.* 1981, A&A, 103, 393. Effelsberg 100-m at 2.7 GHz and 1.7 GHz (7'6": $S=31.6\pm 2.5$ Jy), and 151 MHz (4': $S=62.6\pm 6$ Jy).
 Fesen *et al.* 1981, ApJ, 247, 148. Optical, including spectra.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Simonetti 1992, ApJ, 386, 170. VLA observations of background sources for rotation measure studies.
 Pineault *et al.* 1993, AJ, 105, 1060. DRAO at 1.4 GHz (1': $S=34\pm 4$ Jy), plus H_i and IRAS.
 Seward *et al.* 1995, ApJ, 453, 284. ROSAT observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Slane *et al.* 1997, ApJ, 485, 221. ASCA and ROSAT observations.
 Pineault *et al.* 1997, A&A, 324, 1152. DRAO at 408 MHz (3'5") and 1.4 GHz (1'0").
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Brazier *et al.* 1998, MNRAS, 295, 819. Studies of central, compact γ -ray source.
 Mavromatakis *et al.* 2000, A&A, 353, 371. Optical imagery.
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 1.4 GHz (9'4").
 Slane *et al.* 2004, ApJ, 601, 1045. XMM observations of central source.
 Halpern *et al.* 2004, ApJ, 612, 398. Chandra observations of central nebula, plus optical and radio limits for compact source.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Abdo *et al.* 2008, Science, 322, 1218. Fermi detection of pulsar.
 Lin *et al.* 2010, ApJ, 725, L1. XMM observations of pulsar.
 Caraveo *et al.* 2010, ApJ, 725, L6. XMM observations of pulsar.
 Sun *et al.* 2011, A&A, 535, A64. Urumqi 25-m at 4.8 GHz (9'5": $S=11.6\pm 1.2$ Jy) and Effelsberg 100-m at 2.6 GHz (4'4": $S=20.3\pm 2.0$ Jy) including polarisation.
 Lin *et al.* 2012, MNRAS, 426, 2283. Suzaku observations.
 Mignani *et al.* 2013, MNRAS, 430, 1354. Optical limits for pulsar.
 Aliu *et al.* 2013, ApJ, 764, 38. γ -ray observations.

G120.1 + 1.4

Tycho, 3C10, SN1572

RA: 00^h25^m18^s**1-GHz flux/Jy:** 56**Size/arcmin:** 8**Dec:** +64°09′**Spectral index:** 0.58**Type:** S

This is the remnant of the Tycho's SN of AD1572.

Radio: Shell, brightest to the NE.

Optical: Faint filaments/knots to the NNW, NE and E.

X-ray: Shell, brighter to the NE.

Point sources: Faint radio source near centre of the remnant, thought to be extragalactic.

Distance: H_I observations suggest 2.3--3 kpc, optical proper motion and shock velocity gives 2.4 kpc.

References:

- Duin & Strom 1975, A&A, 39, 33. WSRT at 610 MHz (57''×64'') and 5 GHz (7''×8'').
- Klein *et al.* 1979, A&A, 76, 120. Effelsberg 100-m at 10.7 GHz (1'2 : S = 13.1±0.8 Jy), plus review of flux densities.
- Strom *et al.* 1982, MNRAS, 200, 473. WSRT at 1415 MHz (27''×31'') from 1971 and 1979, for expansion.
- Dickel *et al.* 1982, ApJ, 257, 145. Comparison of radio, X-ray and optical observations.
- Reid *et al.* 1982, ApJ, 261, 485. Einstein observations.
- Seward *et al.* 1983, ApJ, 266, 287. Einstein observations.
- Tan & Gull 1985, MNRAS, 216, 949. Radio at 2.7 GHz (4'') from 1980 and 1983, and One-Mile Telescope at 1.4 GHz (23'') from 1965 and 1980, for expansion.
- Albinson *et al.* 1986, MNRAS, 219, 427. H_I observations.
- Green & Gull 1987, MNRAS, 224, 1055. VLA H_I absorption observations towards central radio source.
- Smith *et al.* 1988, ApJ, 325, 288. EXOSAT spectrum and image (smoothed to 13'').
- Dickel *et al.* 1991, AJ, 101, 2151. VLA at 1.4 and 5 GHz (1''5).
- Wood *et al.* 1992, AJ, 103, 1338. VLA at 5 GHz (1''5) polarisation studies.
- Fink *et al.* 1994, A&A, 283, 635. X-ray spectra.
- Vancura *et al.* 1995, ApJ, 441, 680. X-ray spectra and ROSAT image.
- Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
- Schwarz *et al.* 1995, A&A, 299, 193. WSRT and Effelsberg 100-m at 1.4 GHz (50''), and VLA at 1.4 GHz (13''), for neutral hydrogen studies.
- Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
- Hwang & Gotthelf 1997, ApJ, 475, 665. ASCA observations.
- Reynoso *et al.* 1997, ApJ, 491, 816. VLA at 1.4 GHz (1''4×1''5) from 1984 and 1994 for expansion studies.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Reynoso *et al.* 1999, AJ, 117, 1827. VLA at 1.4 GHz (13''), for H_I studies.
- Katz-Stone *et al.* 2000, ApJ, 529, 453. VLA at 333 MHz and 1.4 GHz (9''5×7''7), for spectral index studies.
- Ghavamian *et al.* 2000, ApJ, 535, 266. Optical observations of shock precursors.
- Hughes 2000, ApJ, 545, L53. ROSAT X-ray expansion.
- Decourchelle *et al.* 2001, A&A, 365, L218. XMM observations.
- Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
- Douvion *et al.* 2001, A&A, 373, 281. ISO observations.
- Aharonian *et al.* 2001, A&A, 373, 292. Upper limit on high energy γ -rays.
- Hwang *et al.* 2002, ApJ, 581, 1101. Chandra observations.
- Lee *et al.* 2004, ApJ, 605, L113. Observations of molecular clouds in vicinity.
- Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
- Warren *et al.* 2005, ApJ, 634, 376. Chandra observations.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=86\pm 5$ Jy) and 1420 MHz ($\sim 1'$: $S=40.5\pm 1.5$ Jy), including polarisation and review of flux densities.

Lee *et al.* 2007, ApJ, 659, L133. H α observations.

Cassam-Chenai *et al.* 2007, ApJ, 665, 315. Chandra observations.

Rest *et al.* 2008, ApJ, 681, L81. SN light echo.

Krause *et al.* 2008, Nature, 456, 617. SN light echo spectrum.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Tamagawa *et al.* 2009, PASJ, 61, S167. Suzaku observations.

Furuzawa *et al.* 2009, ApJ, 693, L61. Suzaku observations.

Lee *et al.* 2010, ApJ, 715, L146. H α observations.

Katsuda *et al.* 2010, ApJ, 709, 1387. Chandra proper motions study.

Raymond *et al.* 2010, ApJ, 712, 901. H α spectroscopy.

Hayato *et al.* 2010, ApJ, 725, 894. Suzaku observations of expansion.

Ishihara *et al.* 2010, A&A, 521, L61. AKARI observations.

Tian & Leahy 2011, ApJ, 729, L15. H α and CO observations.

Acciari *et al.* 2011, ApJ, 730, L20. γ -ray observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5$: $S=20.0\pm 2.0$ Jy) including polarisation and review of flux densities.

Giordano *et al.* 2012, ApJ, 744, L2. Fermi observations.

Gomez *et al.* 2012, MNRAS, 420, 3557. Herschel IR dust observations.

Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.

Williams *et al.* 2013, ApJ, 770, 129. Spitzer observations.

G126.2 + 1.6

RA: 01^h22^m00^s

Dec: +64°15'

1-GHz flux/Jy: 6

Spectral index: 0.5

Size/arcmin: 70

Type: S?

Radio: Poorly defined shell.

Optical: Filaments, mostly in W.

References:

Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz ($9'$: $S=6.8\pm 0.7$ Jy) and 2.7 GHz ($4'4$: $S=3.9\pm 0.4$ Jy).

Blair *et al.* 1980, ApJ, 242, 592. Optical detection and spectra.

Rosado 1982, RMxAA, 5, 127. Optical spectra.

Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.

Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz ($4'4$) and 4.8 GHz ($2'6$).

Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz ($3'5\times 3'9$: $S=12\pm 2.5$ Jy) and part at 1.4 GHz ($1'1\times 1'0$), plus review of flux densities.

Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'5$: $S=6.1\pm 1.6$ Jy).

Boumis *et al.* 2005, A&A, 443, 175. Optical observations.

Tian & Leahy 2006, A&A, 447, 205. CGPS at 408 MHz ($3'4\times 3'8$: $S=9.7\pm 3.9$ Jy) and 1.4 GHz ($1'0\times 1'1$: $S=6.7\pm 2.1$ Jy), plus other observations for spectral index studies.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=5.7\pm 0.7$ Jy) and 1420 MHz ($\sim 1'$: $S=6.4\pm 1.1$ Jy), including review of flux densities.

Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz ($9'5$: $S=2.6\pm 0.6$ Jy), including polarisation. *see also:* Sun *et al.* 2007, A&A, 469, 1003. Erratum.

G127.1+0.5

R5

RA: 01^h28^m20^s**1-GHz flux/Jy:** 12**Size/arcmin:** 45**Dec:** +63°10′**Spectral index:** 0.45**Type:** S

Has been called G127.3+0.7.

Radio: Distinct shell, with bright central source.

Optical: Detected.

Point sources: Flat radio spectrum (extragalactic) source at centre of remnant.

Distance: 1.2--1.3 kpc if associated with NGC 559.

References:

Caswell 1977, MNRAS, 181, 789. Half-Mile Telescope at 1.42 GHz (3′5×3′9), plus other observations of central source.

Pauls 1977, A&A, 59, L13. Effelsberg 100-m at 1.4 GHz (9′:S=8±1 Jy).

Salter *et al.* 1978, A&A, 66, 77. Effelsberg 100-m at 2.7 GHz (4′4), plus 5 and 8.7 GHz of the central source.

Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz (9′:S=10.8±1.3 Jy).

Pauls *et al.* 1982, A&A, 112, 120. WSRT at 610 MHz (56″×62″) and H_I absorption to the point source with the VLA.

Geldzahler & Shaffer 1982, ApJ, 260, L69. Observations of central source.

Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz (4′4) and 4.8 GHz (2′6).

Goss & van Gorkom 1984, JApA, 5, 425. WSRT H_I absorption of central source.

Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz (3′5×3′9 : S = 17.9±2.0 Jy) and 1.4 GHz (1′13×1′0 : S = 10.1±0.8 Jy), plus review of flux densities.

Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′5 : S = 14.6±0.8 Jy).

Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact sources.

Leahy & Tian 2006, A&A, 451, 251. CGPS at 408 MHz (3′4×3′8 : S = 17.1±/1.7 Jy) and 1.4 GHz (1′0×1′2 : S = 10.0±/0.8 Jy).

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′ : S = 15.9±1.0 Jy) and 1420 MHz (~1′ : S = 9.7±0.6 Jy), including polarisation and review of flux densities.

Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz (9′5 : S = 6.3±0.7 Jy), including polarisation. *see also:* Sun *et al.* 2007, A&A, 469, 1003. Erratum.

G130.7+3.1

3C58, SN1181

RA: 02^h05^m41^s**1-GHz flux/Jy:** 33**Size/arcmin:** 9×5**Dec:** +64°49′**Spectral index:** 0.07**Type:** F

This is the remnant of the SN of AD1181.

Radio: Filled-centre, highly polarised, with high frequency turnover.

Optical: Faint filaments.

X-ray: Centrally brightened, with faint jet.

Point sources: Central pulsar.

Distance: H_I absorption indicates 2 kpc.

References:

Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz (58″ : S = 26.7±0.5 Jy).

- Wilson & Weiler 1976, A&A, 49, 357. WSRT at 610 MHz ($58'' \times 64''$), 1.4 GHz ($24'' \times 27''$) and 5 GHz ($7'' \times 8''$).
- van den Bergh 1978, ApJ, 220, L9. Optical observations.
- Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.
- Green & Gull 1982, Nature, 299, 606. H α absorption distance.
- Fesen 1983, ApJ, 270, L53. Optical spectra.
- Reynolds & Aller 1985, AJ, 90, 2312. VLA at 1.4 GHz ($2'$), for limits of shell.
- Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT spectrum.
- Green 1986, MNRAS, 218, 533. 151 MHz observations ($1'.2 \times 1'.3 : S = 36 \pm 4$ Jy), plus 2.7 GHz ($4''$), plus Einstein observations for limit on shell.
- Green 1987, MNRAS, 225, 11P. Flux density increase at 408 MHz.
- Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz ($26''.5 : S = 24.2 \pm 1.4$ Jy).
- Reynolds & Aller 1988, ApJ, 327, 845. VLA at 1.4 ($2''.4$) and 4.9 GHz ($2''.5$).
- Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ($90'' : S = 15.0 \pm 2.0$ Jy), plus review of flux densities.
- Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
- Green & Scheuer 1992, MNRAS, 258, 833. IRAS upper limits.
- Roberts *et al.* 1993, A&A, 274, 427. H α absorption.
- Wallace *et al.* 1994, A&A, 286, 565. H α of surroundings.
- Helfand *et al.* 1995, ApJ, 453, 741. ROSAT observations.
- Torii *et al.* 2000, PASJ, 52, 875. ASCA observations.
- Bietenholz *et al.* 2001, ApJ, 560, 772. VLA at 74 MHz ($26'' : S = 33.6$ Jy) and 327 MHz ($8''.2 : S = 33.9$ Jy), for spectral index and expansion studies.
- Bocchino *et al.* 2001, A&A, 369, 1078. XMM observations.
- Murray *et al.* 2002, ApJ, 568, 226. Chandra pulsar detection.
- Camilo *et al.* 2002, ApJ, 571, L41. Pulsar detection in radio.
- Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz ($26''$) for polarised intensity.
- Slane *et al.* 2002, ApJ, 571, L45. Chandra observations.
- Slane *et al.* 2004, ApJ, 616, 403. Deep Chandra imaging.
- Bietenholz 2006, ApJ, 645, 1180. VLA at 1.4 GHz ($1''.36$) for expansion studies.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 32.2 \pm 2.0$ Jy) and 1420 MHz ($\sim 1' : S = 31.9 \pm 1.0$ Jy), including polarisation and review of flux densities.
- Gotthelf *et al.* 2007, ApJ, 654, 267. XMM observations.
- Slane *et al.* 2008, ApJ, 676, L33. Spitzer and other IR observations.
- Fesen *et al.* 2008, ApJS, 174, 379. Optical observations for proper motion studies.
- Shearer & Neustroev 2008, MNRAS, 390, 235. Optical observations of pulsar nebula.
- Shibanov *et al.* 2008, A&A, 486, 273. Optical observations of pulsar nebula.
- Abdo *et al.* 2009, ApJ, 699, L102. Fermi observations of pulsar.
- Livingstone *et al.* 2009, ApJ, 706, 1163. Pulsar observations.
- Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 31.7 \pm 3.0$ Jy) including polarisation and review of flux densities.
- Bietenholz *et al.* 2013, MNRAS, 431, 2590. Proper motion study of pulsar.
- Kothes *et al.* 2013, A&A, 560, A18. CGPS H α observations for distance.

G132.7+1.3

HB3

RA: 02^h17^m40^s**1-GHz flux/Jy:** 45**Size/arcmin:** 80**Dec:** +62°45′**Spectral index:** 0.6**Type:** S

Has been called G132.4+2.2.

Radio: Faint shell, adjacent to W3/4/5 complex.

Optical: Complete, filamentary shell, shock excited spectra.

X-ray: Partial shell.

Point sources: Pulsar nearby.

Distance: Interaction with surroundings suggests 2.2 kpc.

References:

van den Bergh *et al.* 1973, *ApJS*, 26, 19. Optical observations.

Velusamy & Kundu 1974, *A&A*, 32, 375. NRAO 300-ft at 2.7 GHz (5′:S=33.8±7.0 Jy).

d’Odorico & Sabbadin 1977, *A&AS*, 28, 439. Optical spectra.

Read 1981, *MNRAS*, 194, 863. Radio at 151 MHz (4′4) and 1.4 GHz (2′) showing H_I shell.

Lozinskaya 1981, *SvAL*, 7, 17. Mean optical velocity.

Fesen & Gull 1983, *PASP*, 95, 196. Optical image.

Leahy *et al.* 1985, *ApJ*, 294, 183. Einstein observations.

Landecker *et al.* 1987, *AJ*, 94, 111. DRAO at 408 MHz (3′5×4′:S=75±15 Jy), plus review of flux densities.

Routledge *et al.* 1991, *A&A*, 247, 529. DRAO at 1.4 GHz (1′0×1′1) for H_I, plus CO observations.

Fesen *et al.* 1995, *AJ*, 110, 2876. Optical imaging and spectroscopy, DRAO at 408 MHz (3′5×4′) and 1.4 GHz (1′0×1′1).

Lorimer *et al.* 1998, *A&A*, 331, 1002. Pulsar search.

Koralesky *et al.* 1998, *AJ*, 116, 1323. VLA detection of compact OH emission.

Reich *et al.* 2003, *A&A*, 408, 961. Effelsberg 100-m at 863 MHz (14′5:S=51.5±3.5 Jy).

Tian & Leahy 2005, *A&A*, 436, 187. CGPS at 408 MHz (3′4×3′8:.) and 1.4 GHz (1′0×1′1), for spectral index studies.

see also: Tian & Leahy 2006, *A&A*, 451, 991. Erratum.

Kothes *et al.* 2006, *A&A*, 457, 1081. CGPS at 408 MHz (~3′:S=61±9 Jy) and 1420 MHz (~1′:S=29.4±2.7 Jy), including review of flux densities.

Lazendic & Slane 2006, *ApJ*, 647, 350. X-ray observations.

Green 2007, *BASI*, 35, 77. Review of radio spectrum.

Shi *et al.* 2008, *A&A*, 487, 601. Urumqi 25-m at 4.8 GHz (9′5), plus other survey observations for spectral studies.

Casandjian & Grenier 2008, *A&A*, 489, 849. γ -ray observations.

G152.4–2.1**RA:** 04^h07^m50^s**1-GHz flux/Jy:** 3.5?**Size/arcmin:** 100×95**Dec:** +49°11′**Spectral index:** 0.7?**Type:** S

Radio: Bilateral shell.

References:

Foster *et al.* 2013, *A&A*, 549, A107. Effelsberg 100-m at 2.7 GHz, including polarisation, plus various radio survey observations.

G156.2 + 5.7**RA:** 04^h58^m40^s**Dec:** +51°50′**1-GHz flux/Jy:** 5**Spectral index:** 0.5**Size/arcmin:** 110**Type:** S**Radio:** Faint shell, brighter in E and W.**Optical:** Filamentary ring and smaller patchy ring.**X-ray:** Faint shell.**References:**Pfeffermann *et al.* 1991, A&A, 246, L28. ROSAT detection.Reich *et al.* 1992, A&A, 256, 214. Effelsberg 100-m at 1.4 (9′ : S = 4.2 ± 1.0 Jy) and 2.7 GHz (4′.3 : S = 3.0 ± 1.0 Jy), plus H α and IRAS.Yamauchi *et al.* 1993, PASJ, 45, 795. Hard X-ray observations.Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.Yamauchi *et al.* 1999, PASJ, 51, 13. ASCA observations of some regions.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz, including polarisation.

Pannuti & Allen 2004, AdSpR, 33, 434. ASCA and RXTE observations.

Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.Kotthes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3′$: S = 8.1 ± 1.3 Jy), including review of flux densities.Xu *et al.* 2007, A&A, 470, 969. Urumqi 25-m at 4.8 GHz (9′.5 : S = 2.5 ± 0.5 Jy), including polarisation.

Gerardy & Fesen 2007, MNRAS, 376, 929. Optical observations.

Katsuda *et al.* 2009, PASJ, 61, S155. Suzaku observations.Uchida *et al.* 2012, PASJ, 64, 61. Suzaku observations.**G159.6 + 7.3****RA:** 05^h20^m00^s**Dec:** +50°00′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 240 × 180?**Type:** S**Optical:** Large, faint shell.**X-ray:** Possible emission.**References:**Fesen & Milisavljevic 2010, AJ, 140, 1163. H α and other observations.

G160.9+2.6

HB9

RA: 05^h01^m00^s**1-GHz flux/Jy:** 110**Size/arcmin:** 140×120**Dec:** +46°40′**Spectral index:** 0.64**Type:** S

Has been called G160.5+2.8 and G160.4+2.8.

Radio: Large, filamentary shell.

Optical: Incomplete shell.

X-ray: Centrally brightened.

Point sources: Pulsar within boundary of the remnant, plus several nearby compact radio sources.

Distance: Various observations suggests less than 4 kpc.

References:

d’Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.

Damashek *et al.* 1978, ApJ, 225, L31. Pulsar.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Dwarakanath *et al.* 1982, JApA, 3, 207. Radio observations at 34.5 MHz (26′×40′: $S=750\pm 150$ Jy), plus review of flux densities.

van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H α absorption to nearby point source.

Sequist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.

Leahy 1987, ApJ, 322, 917. Einstein observations.

Leahy & Roger 1991, AJ, 101, 1033. DRAO at 408 MHz (3′.5×4′.8) and 1.4 GHz (1′.0×1′.4), including H α and discussion of distance.

Yamauchi & Koyama 1993, PASJ, 45, 545. Hard X-ray observations.

Leahy & Aschenbach 1995, A&A, 293, 853. ROSAT observations.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Leahy *et al.* 1998, A&A, 339, 601. 232 MHz (3′.8×5′.2), 151 MHz (4′.2×5′.8) and Effelsberg 100-m at 4.7 GHz (2′.5) for spectral index studies.

Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density ($S=1130\pm 340$ Jy).

Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′.5: $S=91\pm 3$ Jy).

Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=102\pm 7$ Jy) and 1420 MHz ($\sim 1'$: $S=54.0\pm 2.9$ Jy), including polarisation and review of flux densities.

Leahy & Tian 2007, A&A, 461, 1013. CGPS at 408 MHz (2′.8×3′.9: $S=117.8\pm 5.3$ Jy) and 1.4 GHz (58″×80″: $S=65.9\pm 3.4$ Jy).

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5: $S=34\pm 3$ Jy), including polarisation and review of flux densities.

Gosachinskii 2013, AstL, 39, 179. H α observations of region.

G166.0+4.3

VRO 42.05.01

RA: 05^h26^m30^s**1-GHz flux/Jy:** 7**Size/arcmin:** 55×35**Dec:** +42°56′**Spectral index:** 0.37**Type:** S

Radio: Two arcs of strikingly different radii.

Optical: Nearly complete ring.

X-ray: Predominantly in SW.

Distance: H α indicates 4.5 kpc.

References:

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 5.2 \pm 1.0$ Jy).
 Lozinskaya 1979, AujPh, 32, 113. H α interferometry.
 Landecker *et al.* 1982, ApJ, 261, L41. DRAO at 1.4 GHz ($1'0 \times 1'4$), plus review of flux densities.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Pineault *et al.* 1985, A&A, 151, 52. VLA at 1.4 GHz ($16'' \times 20''$) of part of remnant, and optical observations.
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Braun & Strom 1986, A&AS, 63, 345. WSRT H α Observations.
 Pineault *et al.* 1987, ApJ, 315, 580. DRAO and VLA combined at 1.4 GHz ($20''$).
 Landecker *et al.* 1989, MNRAS, 237, 277. DRAO at 1.4 GHz ($1'0 \times 1'4$), including H α .
 Burrows & Guo 1994, ApJ, 421, L19. ROSAT images and spectra.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Guo & Burrows 1997, ApJ, 480, L51. ASCA observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Leahy & Tian 2005, A&A, 440, 929. CGPS at 408 MHz ($3'4 \times 5'0$) and 1.4 GHz ($1'0 \times 1'4$), for spectral index studies.
see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 8.1 \pm 0.9$ Jy) and 1420 MHz ($\sim 1' : S = 5.1 \pm 0.4$ Jy), including polarisation and review of flux densities.
 Bocchino *et al.* 2009, A&A, 498, 139. XMM observations.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'5 : S = 3.3 \pm 0.3$ Jy), including polarisation and review of flux densities.
 Araya *et al.* 2013, MNRAS, 434, 2202. Fermi observations.

G178.2–4.2

RA: 05^h35^m05^s

Dec: +28°11'

1-GHz flux/Jy: 2

Spectral index: 0.5

Size/arcmin: 72×62

Type: S

Radio: Faint shell, brighter in NE.

References:

Gao *et al.* 2011, A&A, 532, A144. Urumqi 25-m at 5 GHz ($9'5 : S = 1.0 \pm 0.1$ Jy), plus other observations.

G179.0+2.6

RA: 05^h53^m40^s

Dec: +31°05'

1-GHz flux/Jy: 7

Spectral index: 0.4

Size/arcmin: 70

Type: S?

Radio: Thick shell, with background extragalactic sources near centre.

References:

Fürst & Reich 1986, A&A, 154, 303. Effelsberg 100-m at 1.4 ($9'4$), 2.7 ($4'3$) and 4.75 GHz ($2'4$).
 Fürst *et al.* 1989, A&A, 223, 66. Observations of central, extragalactic source.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'5 : S = 3.2 \pm 0.3$ Jy), including polarisation and review of flux densities.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G180.0–1.7

S147

RA: 05^h39^m00^s**1-GHz flux/Jy:** 65**Size/arcmin:** 180**Dec:** +27°50′**Spectral index:** varies**Type:** S**Radio:** Large faint shell, with spectral break.**Optical:** Wispy ring.**X-ray:** Possible detection.**Point sources:** Pulsar within boundary, with faint wind nebula.**Distance:** Optical absorption towards stars indicates > 0.36 and < 0.88 kpc.**References:**van den Bergh *et al.* 1973, ApJ, 26, 19. Optical observations.Sofue *et al.* 1980, PASJ, 32, 1. Effelsberg 100-m at 5 GHz (2′.6) of parts.Kundu *et al.* 1980, A&A, 92, 225. Effelsberg 100-m at 2.7 GHz (5′.5 : S = 34.9±4 Jy) and 1.6 GHz (10′ : S = 60.2±6 Jy).

Angerhofer & Kundu 1981, AJ, 86, 1003. Arecibo at 430 MHz (9′ : S = 97±20 Jy).

Fürst *et al.* 1982, A&A, 115, 428. Observations of compact radio sources near the remnant.Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.

Fürst & Reich 1986, A&A, 163, 185. Effelsberg 100-m at 1.4, 2.7 and 4.7/5.0 GHz (9′.4, 4′.3 and 2′.4/2′.6).

Sauvageot *et al.* 1990, A&A, 227, 183. EXOSAT possible detection.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Anderson *et al.* 1996, ApJ, 468, L55. Pulsar detection.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.

Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′.5 : S = 77±10 Jy).

Romani & Ng 2003, ApJ, 585, L41. Chandra of pulsar.

Kramer *et al.* 2003, ApJ, 593, L31. Pulsar observations.

Sallmen & Welsh 2004, A&A, 426, 555. Optical absorption towards background stars.

Drew *et al.* 2005, MNRAS, 362, 753. H α imaging.Ng *et al.* 2007, ApJ, 654, 487. Pulsar and wind nebula observations.Xiao *et al.* 2008, A&A, 482, 783. Urumqi 25-m at 4.8 GHz (9′.5 : S = 15.4±3.0 Jy) and Effelsberg 100-m at 2.6 GHz (4′.4 : S = 34.6±4.0 Jy).Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.Katsuta *et al.* 2012, ApJ, 752, 135. Fermi observations.**G182.4 + 4.3****RA:** 06^h08^m10^s**1-GHz flux/Jy:** 0.5**Size/arcmin:** 50**Dec:** +29°00′**Spectral index:** 0.4**Type:** S**Radio:** Incomplete shell.**Optical:** Brighter in S and NW.**X-ray:** Diffuse emission.**References:**Kothes *et al.* 1998, A&A, 331, 661. Effelsberg 100-m at 1.4, 2.7, 4.9 and 10.5 GHz (9′.4 : S = 0.36±0.08 Jy, 4′.4 : S = 0.25±0.04 Jy, 2′.5 : S = 0.20±0.02 Jy and 1′.2 : S = 0.15±0.03 Jy), plus X-ray upper limit.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz and 4.9 GHz (3′).

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 0.26±0.5 Jy) including polarisation and review of flux densities.Sezer *et al.* 2012, MNRAS, 427, 1168. Optical and XMM observations.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G184.6—5.8

Crab Nebula, 3C144, SN1054

RA: 05^h34^m31^s

1-GHz flux/Jy: 1040

Size/arcmin: 7×5

Dec: +22°01′

Spectral index: 0.30

Type: F

This is the remnant of the SN of AD1054.

Radio: Filled-centre, central pulsar, with faint ‘jet’ (or tube) extending from the N edge.

Optical: Strongly polarised filaments, diffuse synchrotron emission, with ‘jet’ faintly visible.

X-ray: Central ‘torus’ around the pulsar.

Point sources: Pulsar powering the remnant.

Distance: Proper motions and radial velocities give 2 kpc.

References:

- Velusamy 1984, *Nature*, 308, 251. VLA at 1.4 GHz, radio detection of ‘jet’.
- Velusamy 1985, *MNRAS*, 212, 359. VLA at 1.4 GHz (15’’).
- Aller & Reynolds 1985, *ApJ*, 293, L73. Flux density decrease.
- Fesen & Gull 1986, *ApJ*, 306, 259. Deep [OIII] imagery of ‘jet’.
- Marcelin *et al.* 1990, *A&A*, 228, 471. Optical of ‘jet’.
- Hester *et al.* 1990, *ApJ*, 357, 539. Optical and IR images.
- Hickson & van den Bergh 1990, *ApJ*, 365, 224. Optical polarisation.
- Bietenholz & Kronberg 1990, *ApJ*, 357, L13. VLA at 1.4 GHz (1’’8).
- Bietenholz & Kronberg 1992, *ApJ*, 393, 206. VLA at 1.5 and 5 GHz (1’’8) and 1.5 and 14 GHz (6’’5) for spectral studies.
- Fesen & Staker 1993, *MNRAS*, 263, 69. [OIII] imaging of ‘jet’, and proper motion studies.
- Kassim *et al.* 1993, *AJ*, 106, 2218. VLA at 74 MHz (20’’).
- Véron-Cetty & Woltjer 1993, *A&A*, 270, 370. Continuum and [OIII] photometry.
- Wallace *et al.* 1994, *A&A*, 286, 565. H α of surroundings.
- Frail *et al.* 1995, *ApJ*, 454, L129. VLA at 333 MHz (20’’) for limits on shell.
- Bietenholz *et al.* 1997, *ApJ*, 490, 291. Comparison of VLA observations at 74 MHz, 327 MHz, 1.5 GHz and 5 GHz for spectral index studies.
- Blair *et al.* 1997, *ApJS*, 109, 473. HST imaging.
- Fesen *et al.* 1997, *AJ*, 113, 354. Limits on H α halo.
- van der Meulen *et al.* 1998, *A&A*, 330, 321. γ -ray observations.
- Tanimori *et al.* 1998, *ApJ*, 492, L33. γ -ray observations.
- Nugent 1998, *PASP*, 110, 831. Optical expansion.
- Sankrit *et al.* 1998, *ApJ*, 504, 344. HST images.
- Wallace *et al.* 1999, *ApJS*, 124, 181. DRAO at 1.4 GHz (1’0×2’8), plus Effelsberg 100-m, for H α studies.
- Greiveldinger & Aschenbach 1999, *ApJ*, 510, 305. X-ray variability of torus.
- Weisskopf *et al.* 2000, *ApJ*, 536, L81. Chandra observations.
- Sollerman *et al.* 2000, *ApJ*, 537, 861. HST observations.
- Aharonian *et al.* 2000, *ApJ*, 539, 317. γ -ray observations.
- Oser *et al.* 2001, *ApJ*, 547, 949. γ -ray observations.
- Willingale *et al.* 2001, *A&A*, 365, L212. XMM observations.
- Douvion *et al.* 2001, *A&A*, 373, 281. ISO observations.
- Bietenholz *et al.* 2001, *ApJ*, 560, 254. Multi-epoch VLA observations, showing variations near the pulsar.
- Bandiera *et al.* 2002, *A&A*, 386, 1044. 1.3 mm observations.
- Hester *et al.* 2002, *ApJ*, 577, L49. HST and Chandra multi-epoch observations.

Atkins *et al.* 2003, ApJ, 595, 803. γ -ray observations.
 Green *et al.* 2004, MNRAS, 355, 1315. Sub-mm and ISO observations.
 Mori *et al.* 2004, ApJ, 609, 186. Chandra observations.
 Čadež *et al.* 2004, ApJ, 609, 797. Optical observations.
 Aharonian *et al.* 2004, ApJ, 614, 897. γ -ray observations.
 Bietchholz *et al.* 2004, ApJ, 615, 794. VLA at 5 GHz (1''4) and HST multi-epoch observation for proper motion studies.
 Melatos *et al.* 2005, ApJ, 633, 931. Multi-epoch near-IR observations central region.
 Seward *et al.* 2006, ApJ, 636, 873. Chandra observations of scattering halo.
 Temim *et al.* 2006, AJ, 132, 1610. Spitzer observations.
see also: Temim *et al.* 2009, AJ, 137, 5155. Erratum.
 Aharonian *et al.* 2006, A&A, 457, 899. γ -ray observations
 Seward *et al.* 2006, ApJ, 652, 1277. Chandra observations.
 MacAlpine *et al.* 2007, AJ, 133, 81. Optical spectroscopy.
 Kaplan *et al.* 2008, ApJ, 677, 1201. HST proper motion of pulsar.
 Rudie *et al.* 2008, MNRAS, 384, 1200. [OIII] observations of 'jet', for proper motion.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Tziamtzis *et al.* 2009, A&A, 497, 167. Limits on H α halo.
 Carlebois *et al.* 2010, AJ, 139, 2083. Optical imaging spectroscopy.
 Aumount *et al.* 2010, A&A, 514, A70. IRAM 30-m at 150 GHz (16''7 : S = 244 \pm 24 Jy) and GBT at 90 GHz (9''3).
 Loh *et al.* 2012, MNRAS, 421, 789. IR observations of H₂.
 Lundqvist & Tziamtzis 2012, MNRAS, 423, 1571. Optical limits on outer shell.
 Temim *et al.* 2012, ApJ, 753, 72. Spitzer observations of dust.
 Gomez *et al.* 2012, ApJ, 760, 96. Herschel IR and sub-mm observations of dust.
 Loll *et al.* 2013, ApJ, 765, 152. HST optical line and continuum observations.
 Barlow *et al.* 2013, Science, 342, 1343. Herschel far-IR spectroscopy.

G189.1 + 3.0

IC443, 3C157

RA: 06^h17^m00^s**1-GHz flux/Jy:** 160**Size/arcmin:** 45**Dec:** +22°34'**Spectral index:** 0.36**Type:** C**Radio:** Limb-brightened to NE, with faint extension to the E.**Optical:** Brightest to the NE, with faint filaments outside the NE boundary.**X-ray:** Shell, brightest to the NE, plus compact source with nebula.**Point sources:** X-ray source and nebula in S.**Distance:** Mean optical velocity suggests 0.7--1.5 kpc, association with S249 gives 1.5--2 kpc.**References:**

Fesen 1984, ApJ, 281, 658. Optical of filament to far NE.
 Erickson & Mahoney 1985, ApJ, 290, 596. TPT at 4 frequencies between 31 and 74 MHz (13'x11' to 5'4x4'7), plus review of flux densities.
 Braun & Strom 1986, A&A, 164, 193. WSRT at 327 MHz (72''x185'') and 1.4 GHz (17''x43''), plus H_I and IRAS.
 Green 1986, MNRAS, 221, 473. 151 MHz observations (1'2x3'1) and Half-Mile Telescope at 1.4 GHz (2'1x5'4).
 Mufson *et al.* 1986, AJ, 92, 1349. Radio, IR, optical, UV and X-ray comparison, including VLA at 1.6 GHz (3''8x3''3 and 40'').
 Petre *et al.* 1988, ApJ, 335, 215. Einstein and other X-ray observations.
 Dickel *et al.* 1989, AJ, 98, 1363. VLA at 1.4 GHz (1''1x1''2) of NE.
 Burton *et al.* 1990, ApJ, 355, 197. IR observations of shocked O₂.
 Wood *et al.* 1991, AJ, 102, 224. VLA at 5 GHz (3''6x3''8) of northeast, including polarisation.

- Wang & Scoville 1992, ApJ, 386, 158. Observations of shocked molecular species.
- Turner *et al.* 1992, ApJ, 399, 114. Observations of shocked molecular species.
- Dickman *et al.* 1992, ApJ, 400, 203. Observations of shocked molecular species.
- van Dishoeck *et al.* 1993, A&A, 279, 541. Observations of shocked molecular species.
- Asaoka & Aschenbach 1994, A&A, 284, 573. X-ray, including possible overlapping remnant.
- Tauber *et al.* 1994, ApJ, 421, 570. Observations of shocked molecular species.
- Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
- Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.
- Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.
- Claussen *et al.* 1999, ApJ, 522, 349. High resolution observations of OH masers.
- Oliva *et al.* 1999, A&A, 341, L75. ISO observations.
- Cesarsky *et al.* 1999, A&A, 348, 945. ISO observations of shocked molecular H₂.
- Bocchino & Bykov 2000, A&A, 362, L29. BeppoSAX observations.
- Rho *et al.* 2001, ApJ, 547, 885. ISO and 2Mass IR spectroscopy and imaging.
- Olbert *et al.* 2001, ApJ, 554, L205. Chandra of compact X-ray source and surrounding nebula.
- Bocchino & Bykov 2001, A&A, 376, 248. XMM of compact X-ray source and surrounding nebula.
- Bocchino & Bykov 2003, A&A, 400, 203. XMM observations of compact sources.
- Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'.5 : S = 160±5 Jy).
- Welsh & Sallmen 2003, A&A, 408, 545. Optical absorption studies.
- Leahy 2004, AJ, 127, 2277. DRAO at 408 MHz (3'.3×8'.6) and 1.4 GHz (1'.0×2'.6), for spectral index studies.
see also: Leahy 2004, AJ, 128, 1478. Addendum.
- Snell *et al.* 2005, ApJ, 620, 758. Molecular line and IR observations of shocked material.
- Bykov *et al.* 2005, ApJ, 624, L41. Chandra observations of compact source.
- Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
- Gaensler *et al.* 2006, ApJ, 648, 1037. Chandra of X-ray source and nebula.
- Hewitt *et al.* 2006, ApJ, 652, 1288. GBT at 1.6 and 1.7 GHz (7'.2) for OH, and VLA at 330 MHz (74''×64'').
- Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
- Rosado *et al.* 2007, AJ, 133, 89. Observations of shocked H₂.
- Albert *et al.* 2007, ApJ, 664, L87. γ -ray observations.
- Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
- Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
- Lee *et al.* 2008, AJ, 135, 796. VLA at 1.4 GHz (39''×42'') including H_i.
- Troja *et al.* 2008, A&A, 485, 777. XMM observations.
- Bocchino *et al.* 2008, AdSpR, 41, 396. Integral observations.
- Bykov *et al.* 2008, ApJ, 676, 1050. XMM, Chandra and Spitzer observations.
- Bocchino *et al.* 2009, A&A, 498, 139. XMM observations.
- Hirschauer *et al.* 2009, ApJ, 696, 1533. Optical absorption spectroscopy.
- Acciari *et al.* 2009, ApJ, 698, L133. γ -ray observations.
- Yamaguchi *et al.* 2009, ApJ, 705, L6. Suzaku observations.
- Koo *et al.* 2010, AJ, 140, 262. H_i Zeeman splitting observations.
- Tavani *et al.* 2010, ApJ, 710, L151. γ -ray observations.
- Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
- Xu *et al.* 2011, ApJ, 727, 81. CO observations of region.
- Shinn *et al.* 2011, ApJ, 732, 124. AKARI observations of H₂.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9'.5 : S = 85±9 Jy), including polarisation and review of flux densities.
- Lee *et al.* 2012, ApJ, 749, 34. CO and HCO⁺ of region.
- Taylor *et al.* 2012, ApJ, 750, L15. Optical absorption of background stars.
- Yuan *et al.* 2012, ApJ, 753, 126. Spitzer spectroscopy.
- Shinn *et al.* 2012, ApJ, 759, 34. AKARI observations of H₂.
- Satterfield *et al.* 2012, AJ, 144, 27. Optical spectroscopy.
- Ackermann *et al.* 2013, Science, 339, 807. Fermi observations.
- Hezareth *et al.* 2013, A&A, 558, A45. CO observations, including polarisation, of region.
- Kokasho *et al.* 2013, ApJ, 768, L8. IR observations of [FeII], plus AKARI and Spitzer observations.

G190.9–2.2**RA:** 06^h01^m55^s**Dec:** +18°24′**1-GHz flux/Jy:** 1.3?**Spectral index:** 0.7?**Size/arcmin:** 70×60**Type:** S**Radio:** Incomplete shell.**References:**Foster *et al.* 2013, A&A, 549, A107. Various radio survey observations.**G192.8–1.1****RA:** 06^h09^m20^s**Dec:** +17°20′**1-GHz flux/Jy:** 20?**Spectral index:** 0.6?

PKS 0607+17

Size/arcmin: 78**Type:** S

Has been called G193.3–1.5. Has been regarded as part of the Origem Loop, a supposed larger remnant.

Radio: In complex region.**Optical:** Encompasses S261 and S254--258.**References:**

Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (9′:S=13±15% Jy).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 318 MHz (15′:S=74±32 Jy), and S_{610 MHz}=40 Jy.

Caswell 1985, AJ, 90, 1076. DRAO at 1.4 GHz (1′×3′3:S=18±3 Jy).

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Welsh *et al.* 2001, A&A, 372, 516. Far-UV spectroscopy.**G205.5+0.5****RA:** 06^h39^m00^s**Dec:** +06°30′**1-GHz flux/Jy:** 140**Spectral index:** 0.4

Monoceros Nebula

Size/arcmin: 220**Type:** S**Radio:** In complex region, parts may be H_{II} regions.**Optical:** Large ring, near Rosette nebula.**X-ray:** Possibly detected.

Distance: Mean optical velocity suggests 0.8 kpc, low frequency radio absorption suggests 1.6 kpc.

References:

Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (9′).

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′), part only.

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 111 MHz (1′:S=462±180 Jy) and S_{610 MHz}=245 Jy.Davies *et al.* 1978, A&AS, 31, 271. Deep optical plates.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Graham *et al.* 1982, A&A, 109, 145. Effelsberg 100-m at 2.7 GHz (4′4:S=97.6±12.5 Jy), plus review of flux densities.Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.Leahy *et al.* 1986, MNRAS, 220, 501. Einstein observations.

Odegard 1986, ApJ, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24′, 19′ and 16′).

Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ -ray emission.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Oliver *et al.* 1996, A&A, 315, 578. CO observations of some of surroundings.
 Jaffe *et al.* 1997, ApJ, 484, L129. γ -ray detection.
 Aharonian *et al.* 2004, A&A, 417, 973. γ -ray limits.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Xiao & Zhu 2012, A&A, 545, A86. Review of radio, H α and H α observations.

G206.9+2.3

PKS 0646+06

RA: 06^h48^m40^s**1-GHz flux/Jy:** 6**Size/arcmin:** 60×40**Dec:** +06°26′**Spectral index:** 0.5**Type:** S?**Radio:** Diffuse source near the Monoceros Nebula.**Optical:** Filaments detected.**X-ray:** Possibly detected.**References:**

Davies & Meaburn 1978, A&A, 69, 443. Optical observations.
 Nousek *et al.* 1981, ApJ, 248, 152. HEAO-1 X-ray limit.
 Graham *et al.* 1982, A&A, 109, 145. Effelsberg 100-m at 2.7 GHz (4′.4 : S = 4.1±0.6 Jy), plus review of flux densities.
 Rosado 1982, RMxAA, 5, 127. Optical observations.
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Leahy 1986, A&A, 156, 191. Einstein observations.
 Odegard 1986, ApJ, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24′, 19′ and 16′).
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 2.9±0.3 Jy), including polarisation and review of flux densities.

G213.0–0.6
RA: 06^h50^m50^s**1-GHz flux/Jy:** 21**Size/arcmin:** 160×140?**Dec:** –00°30′**Spectral index:** 0.4**Type:** S

Has also been called G213.3–0.4.

Radio: Large, faint shell.**Optical:** Filamentary shell.**Point sources:** Central X-ray source**References:**

Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′.5 : S = 22.0±3.7 Jy).
 Stupar & Parker 2012, MNRAS, 419, 1413. H α , radio survey and other observations.

G260.4–3.4

Puppis A, MSH 08–44

RA: 08^h22^m10^s**1-GHz flux/Jy:** 130**Size/arcmin:** 60×50**Dec:** –43°00′**Spectral index:** 0.5**Type:** S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Angular shell, brightest to the E, poorly defined to the W.

Optical: Nebulosity and wisps.

X-ray: Brightest to the E.

Point sources: Central possible pulsating X-ray source.

Distance: Association with H_I gives 2.2 kpc.

References:

- Green 1971, *AJPh*, 24, 773. Molonglo at 408 MHz (3′ : S = 198±20 Jy).
 Goudis & Meaburn 1978, *A&A*, 62, 283. H α + [NII] optical image.
 Petre *et al.* 1982, *ApJ*, 258, 22. Einstein observations.
 Milne *et al.* 1983, *MNRAS*, 204, 237. FIRST at 1415 MHz (50″), large scale emission missing.
 Teske & Petre 1987, *ApJ*, 318, 370. Coronal optical line emission.
 Dubner & Arnal 1988, *A&AS*, 75, 363. H_I and CO observations of surroundings.
 Arendt *et al.* 1990, *ApJ*, 350, 266. MOST at 843 MHz (44″×65″), with large scale emission added, plus IR, optical and X-ray observations.
 Dubner *et al.* 1991, *AJ*, 101, 1466. VLA at 327 MHz (43″×73″) and 1.5 GHz (43″×77″).
 Arendt *et al.* 1991, *ApJ*, 368, 474. IR observations.
 Milne *et al.* 1993, *MNRAS*, 261, 366. Parkes 64-m at 4.75 (4′.5 : S = 59±5 Jy) and 8.4 GHz (3′ : S = 38±4 Jy), plus polarisation.
 Berthiaume *et al.* 1994, *ApJ*, 425, 132. X-ray spectroscopy.
 Sutherland & Dopita 1995, *ApJ*, 439, 365. Spectrophotometry.
 Reynoso *et al.* 1995, *AJ*, 110, 318. VLA at 1.4 GHz (90″) including neutral hydrogen.
 Blair *et al.* 1995, *ApJ*, 454, L35. Far UV spectroscopy.
 Petre *et al.* 1996, *ApJ*, 465, L43. ROSAT of central source.
 Bock *et al.* 1998, *AJ*, 116, 1886. MOST at 843 MHz (43″×60″).
 Pavlov *et al.* 1999, *ApJ*, 511, L45. Possible pulsation detection from central X-ray source.
 Zavlin *et al.* 1999, *ApJ*, 525, 959. X-ray observations of central source.
 Bocchino *et al.* 2000, *A&A*, 359, 316. Optical studies of selected filaments in N.
 Woermann *et al.* 2000, *MNRAS*, 317, 421. OH observations.
 Gaensler *et al.* 2000, *ApJ*, 537, L35. Radio limit for nebula around possible pulsar.
 Reynoso *et al.* 2003, *MNRAS*, 345, 671. ATCA at 1.4 GHz (90″) for H_I near central X-ray source.
 Hwang *et al.* 2005, *ApJ*, 635, 355. Chandra observations of E edge.
 Hui & Becker 2006, *A&A*, 454, 543. XMM and Chandra observations.
 Hui & Becker 2006, *A&A*, 457, L33. Chandra proper motion study of central source.
 Castelletti *et al.* 2006, *A&A*, 459, 535. VLA at 1.4 GHz (16″×34″ : S = 114±8 Jy) and 327 MHz (45″×90″ : S = 263±20 Jy).
 Winkler & Petre 2007, *ApJ*, 670, 635. Chandra proper motion study of central source.
 Paron *et al.* 2008, *A&A*, 480, 439. CO observations of E.
 Hwang *et al.* 2008, *ApJ*, 676, 378. Suzaku observations.
 Katsuda *et al.* 2008, *ApJ*, 678, 297. XMM observations.
 Mignani *et al.* 2009, *A&A*, 500, 1211. Optical limits for compact X-ray source.
 Katsuda *et al.* 2010, *ApJ*, 714, 1725. Chandra and XMM observations.
 Arendt *et al.* 2010, *ApJ*, 725, 585. Spitzer observations.
 Becker *et al.* 2012, *ApJ*, 755, 141. Chandra proper motion study of central X-ray source.
 Katsuda *et al.* 2012, *ApJ*, 756, 49. XMM spectroscopy.
 Hewitt *et al.* 2012, *ApJ*, 759, 89. Fermi and WMAP 23 to 90 GHz observations.
 Dubner *et al.* 2013, *A&A*, 555, A9. XMM and Chandra observations.
 Katsuda *et al.* 2013, *ApJ*, 768, 182. XMM observations.

G261.9 + 5.5**RA:** 09^h04^m20^s**Dec:** -38°42'**1-GHz flux/Jy:** 10?**Spectral index:** 0.4?**Size/arcmin:** 40×30**Type:** S**Radio:** Faint shell with little limb brightening.**References:**

Hill 1967, *AJPh*, 20, 297. Parkes 64-m at 2650 MHz (7'5" : S = 7 Jy) also $S_{1410 \text{ MHz}} = 8 \text{ Jy}$, $S_{81.5 \text{ MHz}} = 25 \text{ Jy}$.

Colomb & Dubner 1980, *A&A*, 82, 244. Argentine 30-m dish at 1.4 GHz, for H_I possibly associated with remnant.

Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44"×71").

Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

G263.9—3.3**RA:** 08^h34^m00^s**Dec:** -45°50'**1-GHz flux/Jy:** 1750**Spectral index:** varies**Size/arcmin:** 255**Type:** C

Vela (XYZ)

This refers to the whole Vela XYZ complex, of which X has at times been classified as a separate (filled-centre) remnant. This remnant is overlapped by G260.4—3.4 and G266.2—1.2.

Radio: Large shell, with flatter spectrum component (Vela X), and pulsar nebula.

Optical: Filaments.

X-ray: Patchy shell, with extensions, central nebula and pulsar.

Point sources: Pulsar within Vela X, with one-sided 'jet'.

Distance: Vela pulsar parallax gives 0.3 kpc, optical spectra and H_I studies suggest 0.25 kpc.

References:

Milne 1968, *AJPh*, 21, 201. Parkes 64-m at various frequencies, including 408 MHz (48' : S = 2300±300 Jy), 635 MHz (31' : S = 2360±300 Jy), 1410 MHz (14' : S = 1640±300 Jy) and 2650 MHz (7'5" : S = 1400±250 Jy), plus discussion of the distance.

Milne 1980, *A&A*, 81, 293. Maps of Vela X with Parkes 64-m at 1.66, 2.7 and 5 GHz (12', 8'.4 and 4'.4).

Weiler & Panagia 1980, *A&A*, 90, 269. Clarification of notation of this region and review previous observations.

Bignami & Caraveo 1988, *ApJ*, 325, L5. Pulsar proper motion from optical observations.

Dwarakanath 1991, *JApA*, 12, 199. Gauribidanur 'T' array at 34.5 MHz (26'×84' : S = 1800 Jy for Vela X and S = 3900 Jy for Vela YZ), plus review of flux densities.

Bietenholz *et al.* 1991, *ApJ*, 376, L41. VLA at 5 GHz in vicinity of pulsar.

Willmore *et al.* 1992, *MNRAS*, 254, 139. Hard X-ray observations.

Dubner *et al.* 1992, *A&AS*, 96, 505. Argentine 30-m at 1.4 GHz (30') of surrounding H_I.

Strom *et al.* 1995, *Nature*, 373, 590. Radio of X-ray extensions.

Markwardt & Ögelman 1995, *Nature*, 375, 40. X-ray jet from pulsar.

Jenkins & Wallerstein 1995, *ApJ*, 440, 227. Optical absorption of associated neutral carbon cloud.

Milne 1995, *MNRAS*, 277, 1435. Parkes 64-m at 8.4 GHz (3'), including polarisation, of Vela X.

Danks & Sembach 1995, *AJ*, 109, 2627. Optical spectroscopy of background stars.

Blair *et al.* 1995, *AJ*, 110, 312. UV spectroscopy.

Duncan *et al.* 1996, MNRAS, 280, 252. Parkes 64-m at 2.4 GHz (8'9).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×62'') of part.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
 Frail *et al.* 1997, ApJ, 475, 224. VLA at 330 MHz (1'×1'8) and comparison with ROSAT observations of Vela X.
 Markwardt & Ögelman 1997, ApJ, 480, L13. ASCA observations of pulsar 'jet'.
 Raymond *et al.* 1997, ApJ, 482, 881. UV spectroscopy.
 Yoshikoshi *et al.* 1997, ApJ, 487, L65. γ -ray observations of pulsar.
 Jenkins *et al.* 1998, ApJ, 492, L147. UV absorption to background star.
 Dubner *et al.* 1998, AJ, 116, 813. Parkes 64-m at 1.4 GHz (15') for Hi studies.
 Bock *et al.* 1998, AJ, 116, 1886. MOST at 843 MHz (43''×60'').
 Cha *et al.* 1999, ApJ, 515, L25. Optical spectra, for distance.
 Lu & Aschenbach 2000, A&A, 362, 1083. ROSAT spatially resolved spectroscopy.
 Chadwick *et al.* 2000, ApJ, 537, 414. Limit on high energy γ -rays from pulsar.
 Sankrit *et al.* 2001, ApJ, 549, 416. Far-UV observations of selected region.
 Pavlov *et al.* 2001, ApJ, 554, L189. Chandra two epoch observations of pulsar nebula.
 Helfand *et al.* 2001, ApJ, 556, 380. Chandra observations of pulsar and its nebula.
 Miyata *et al.* 2001, ApJ, 559, L45. Chandra of X-ray extension.
 Alvarez *et al.* 2001, A&A, 372, 636. Radio spectral index studies.
see also: Alvarez et al. 2001, A&A, 379, 323. Erratum.
 Caraveo *et al.* 2001, ApJ, 561, 930. HST parallax observations of pulsar.
 Moriguchi *et al.* 2001, PASJ, 53, 1025. CO observations.
 Dodson *et al.* 2003, MNRAS, 343, 116. ATCA at 1.4, 2.4, 5.2 and 8.5 GHz (6''.2×8''.1, 26''.×36'', 12''.1×10''.5 and 11''.2×10''.6) of pulsar nebula.
 Sankrit *et al.* 2003, ApJ, 589, 242. Optical nebulosity to NE of G266.2–1.2.
 Nichols & Slavin 2004, ApJ, 610, 285. UV absorption toward background sources.
 Hales *et al.* 2004, ApJ, 613, 977. Vela X at 31 GHz (4'.1).
 Mongano *et al.* 2005, A&A, 436, 917. XMM and other X-ray observations of pulsar nebula.
 Miceli *et al.* 2005, A&A, 442, 513. XMM observations of N rim.
 Katsuda & Tsunemi 2005, PASJ, 57, 621. XMM observations of E.
 Aharonian *et al.* 2006, A&A, 448, L43. γ -ray observations.
 McConnell *et al.* 2006, AJ, 131, 648. ATCA at 4.9 GHz (12') including polarisation.
 Katsuda & Tsunemi 2006, ApJ, 642, 917. XMM observations of NE.
 Nishikida *et al.* 2006, ApJ, 644, L171. Far UV observations.
 Miceli *et al.* 2008, ApJ, 676, 1064. XMM observations.
 LaMassa *et al.* 2008, ApJ, 689, L121. XMM observations of Vela X.
 Grondin *et al.* 2013, ApJ, 774, 110. Fermi observations of Vela X.

G266.2–1.2

RA: 08^h52^m00^s
Dec: –46°20'

1-GHz flux/Jy: 50?
Spectral index: 0.3?

RX J0852.0–4622

Size/arcmin: 120
Type: S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Incomplete shell, confused by the Vela SNR.

Optical: Nebulosity offset to NE.

X-ray: Non-thermal shell, confused by the Vela SNR, with central source, and possible associated pulsar.

Point sources: Central X-ray source, with optical nebula, and possible associated pulsar.

Distance: X-ray data suggest an upper limit of 1 kpc.

References:

Aschenbach 1998, *Nature*, 396, 141. ROSAT identification.
 Iyudin *et al.* 1998, *Nature*, 396, 142. γ -ray observations.
 Combi *et al.* 1999, *ApJ*, 519, L177. Radio observations.
 Duncan & Green 2000, *A&A*, 364, 732. Parkes 64-m at 1.4 GHz (14'0), and comparison with other observations.
 Redman *et al.* 2000, *ApJ*, 543, L153. Optical of nearly nebulousity.
 Tsunemi *et al.* 2000, *PASJ*, 52, 887. ASCA spectral observations.
 Slane *et al.* 2001, *ApJ*, 548, 814. ASCA observations.
 Mereghetti *et al.* 2001, *ApJ*, 548, L213. BeppoSAX observations of central sources.
 Pavlov *et al.* 2001, *ApJ*, 559, L131. Chandra of central X-ray source.
 Moriguchi *et al.* 2001, *PASJ*, 53, 1025. CO observations.
 Pellizzoni *et al.* 2002, *A&A*, 393, L65. Optical observations of central source.
 Redman *et al.* 2002, *MNRAS*, 336, 1093. Optical nebulousity to NE.
 Kargaltsev *et al.* 2002, *ApJ*, 580, 1060. Chandra observations of central source.
 Sankrit *et al.* 2003, *ApJ*, 589, 242. Optical nebulousity to NE.
 Redman & Meaburn 2005, *MNRAS*, 356, 969. Possible pulsar association.
 Iyudin *et al.* 2005, *A&A*, 429, 225. XMM observations.
 Aharonian *et al.* 2005, *A&A*, 437, L7. γ -ray observations.
 Katagiri *et al.* 2005, *ApJ*, 619, L163. γ -ray observations.
 Bamba *et al.* 2005, *ApJ*, 632, 294. Chandra of NW rim.
 Reynoso *et al.* 2006, *A&A*, 449, 243. ATCA at 1.38 GHz (32'' \times 37'').
 Enomoto *et al.* 2006, *ApJ*, 652, 1268. γ -ray observations.
 Mignani *et al.* 2007, *A&A*, 473, 883. Deep optical observations of compact X-ray source.
 Aharonian *et al.* 2007, *ApJ*, 661, 236. γ -ray observations.
 Katsuda *et al.* 2008, *ApJ*, 678, L35. XMM proper motion study.

G272.2–3.2

RA: 09^h06^m50^s
Dec: –52°07'

1-GHz flux/Jy: 0.4
Spectral index: 0.6

Size/arcmin: 15?
Type: S?

Radio: Diffuse shell.

Optical: Detected.

X-ray: Centrally brightened.

References:

Greiner *et al.* 1994, *A&A*, 286, L35. ROSAT observations, plus optical observations.
 Duncan *et al.* 1997, *MNRAS*, 289, 97. Parkes 64-m at 1.4 GHz (18' : $S = 0.38 \pm 0.09$ Jy) 2.4 GHz (10'6 : $S = 0.25 \pm 0.04$ Jy) and 4.8 GHz (5'7 : $S = 0.17 \pm 0.02$ Jy), MOST at 843 MHz (45'' \times 70'' : $S = 0.45 \pm 0.10$ Jy), and ATCA at 2.4 GHz (37'' \times 52''), plus ROSAT observations.
 Harrus *et al.* 2001, *ApJ*, 552, 614. ASCA and ROSAT observations, plus review of earlier observations.

G279.0+1.1

RA: 09^h57^m40^s
Dec: –53°15'

1-GHz flux/Jy: 30?
Spectral index: 0.6?

Size/arcmin: 95
Type: S

Radio: Faint, incomplete shell.

Optical: Detected.

Point sources: Pulsar nearby.

References:

Woermann & Jonas 1988, MNRAS, 234, 971. Hartesbeesthoek 26-m at 1.6 ($30' : S = 25.2 \pm 4$ Jy) and 2.3 GHz ($20' : S = 20.7 \pm 3$ Jy).
 Duncan *et al.* 1995, MNRAS, 277, 319. Parkes 64-m at 1.4 ($18' : S = 28 \pm 3$ Jy) and 2.4 GHz ($11' : S = 20 \pm 2$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 53''$) of part.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G284.3–1.8

MSH 10–53

RA: 10^h18^m15^s**1-GHz flux/Jy:** 11?**Size/arcmin:** 24?**Dec:** –59°00′**Spectral index:** 0.3?**Type:** S

Has been called G284.2–1.8.

Radio: Incomplete, poorly defined shell.

Point sources: Pulsar with wind nebula nearby.

References:

Ruiz & May 1986, ApJ, 309, 667. CO and optical observations.
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz ($43'' \times 50''$) and Parkes 64-m at 8.4 GHz ($3' : S = 5.4 \pm 0.8$ Jy) including polarisation, plus earlier flux densities.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
 Camilo *et al.* 2001, ApJ, 557, L51. Observations of nearby pulsar.
 Camilo *et al.* 2004, ApJ, 616, 1118. Chandra observations of pulsar and nebula.
 Abramowski *et al.* 2012, A&A, 541, A5. γ -ray observations.

G286.5–1.2**RA:** 10^h35^m40^s**1-GHz flux/Jy:** 1.4?**Size/arcmin:** 26×6**Dec:** –59°42′**Spectral index:** ?**Type:** S?

Radio: Double, elongated arc.

Optical: Detected.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 1.6$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G289.7–0.3**RA:** 11^h01^m15^s**1-GHz flux/Jy:** 6.2**Size/arcmin:** 18×14**Dec:** –60°18′**Spectral index:** 0.2?**Type:** S

Radio: Incomplete shell.

Point sources: Compact radio source near centre.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 6.4 \pm 0.5$ Jy), plus Parkes 64-m at 4.5 GHz ($S = 7.5 \pm 2.5$ Jy) and 8.55 GHz ($S = 3.6 \pm 0.9$ Jy).

G290.1–0.8

MSH 11–61A

RA: 11^h03^m05^s**1-GHz flux/Jy:** 42**Size/arcmin:** 19×14**Dec:** –60°56′**Spectral index:** 0.4**Type:** S**Radio:** Elongated, clumpy shell.**Optical:** Filaments detected.**X-ray:** Centrally brightened.**Point sources:** Pulsar nearby.**Distance:** H α absorption indicates 7±1 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4.4′:S=20.2 Jy).
 Elliott & Malin 1979, MNRAS, 186, 45P. Optical image and spectra.
 Kirshner & Winkler 1979, ApJ, 227, 853. Optical observations.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×50″).
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43″×49″:S=45±11 Jy), and Parkes 64-m at 8.4 GHz (3′:S=19.5±1.0 Jy), including polarisation.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×49″:S=43 Jy).
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Kaspi *et al.* 1997, ApJ, 485, 820. Pulsar detection.
 Gotthelf & Kaspi 1998, ApJ, 497, L29. ASCA observations of pulsar.
 Slane *et al.* 2002, ApJ, 564, 284. ASCA observations.
 Filipović *et al.* 2005, SerAJ, 170, 47. ATCA at 1.4 GHz (21″), plus other observations, including CO of surroundings.
 Reynoso *et al.* 2006, MNRAS, 369, 416. ATCA at 1.4 GHz (22″.5×25″.0) including H α .

G291.0–0.1

(MSH 11–62)

RA: 11^h11^m54^s**1-GHz flux/Jy:** 16**Size/arcmin:** 15×13**Dec:** –60°38′**Spectral index:** 0.29**Type:** C**Radio:** Centrally brightened core, with surrounding arcs.**Optical:** Detected.**X-ray:** Centrally brightened.**Point sources:** Central compact X-ray source.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Roger *et al.* 1986, MNRAS, 219, 815. MOST at 843 MHz (43″×50″:S=17.2±1.0 Jy), and Parkes 64-m at 5 and 8.4 GHz (4.6 and 3′:S=10.4±0.4 Jy and 9′.1±0.2), with polarisation.
 Wilson 1986, ApJ, 302, 718. Einstein observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×49″:S=12.7 Jy).
 Harrus *et al.* 1998, ApJ, 499, 273. ASCA observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Slane *et al.* 2012, ApJ, 749, 131. Chandra, XMM and Fermi observations.

G292.0+1.8

MSH 11–54

RA: 11^h24^m36^s**1-GHz flux/Jy:** 15**Size/arcmin:** 12×8**Dec:** −59°16′**Spectral index:** 0.4**Type:** C**Radio:** Centrally brightened source surrounded by a plateau of faint emission.**Optical:** Oxygen rich.**X-ray:** Ring of emission, with diffuse central nebula and pulsar.**Point sources:** Central pulsar.**Distance:** H α absorption implies 6.0 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Lockhart *et al.* 1977, MNRAS, 179, 147. Fleurs at 1415 MHz (50″: $S=13.0$ Jy).
 Goss *et al.* 1979, MNRAS, 188, 357. Optical spectra.
 Clark *et al.* 1980, MNRAS, 193, 129. X-ray spectrum.
 Tuohy *et al.* 1982, ApJ, 260, L65. Einstein observations.
 Dopita & Tuohy 1984, ApJ, 282, 135. Optical spectra.
 Braun *et al.* 1986, A&A, 162, 259. MOST at 843 MHz (0′.8) and IRAS.
 Hughes & Singh 1994, ApJ, 422, 126. EXOSAT spectrum.
 Sutherland & Dopita 1995, ApJ, 439, 365. Spectrophotometry.
 Hughes *et al.* 2001, ApJ, 559, L153. Chandra observations, including central nebula.
 Camilo *et al.* 2002, ApJ, 567, L71. Pulsar detection.
 Park *et al.* 2002, ApJ, 564, L39. Chandra observations.
 Gonzalez & Safi-Harb 2003, ApJ, 583, L91. Chandra observations.
 Hughes *et al.* 2003, ApJ, 591, L139. Chandra observations of pulsar.
 Gaensler & Wallace 2003, ApJ, 594, 326. ATCA at 1.4, 2.3 and 5.2 GHz (9″.6×8″.0, 7″.2×6″.2 and 5″.5×4″.8: $S=11.9\pm 0.1, 11.4\pm 0.1$ and 8.8 ± 0.1), plus H α observations.
 Park *et al.* 2004, ApJ, 602, L33. Chandra observations.
 Vink *et al.* 2004, NuPhS, 132, 62. XMM observations.
 Ghavamian *et al.* 2005, ApJ, 635, 365. Optical imaging spectroscopy.
 Winkler & Long 2006, AJ, 132, 360. Optical observations.
 Park *et al.* 2007, ApJ, 670, L121. Chandra observations.
 Zhariikov *et al.* 2008, A&A, 492, 805. Possible optical counterpart to pulsar and nebula.
 Winkler *et al.* 2009, ApJ, 692, 1489. Optical proper motion studies.
 Ghavamian *et al.* 2009, ApJ, 696, 1307. Spitzer spectroscopy.
 Lee *et al.* 2009, ApJ, 706, 441. IR observations.
 Lee *et al.* 2010, ApJ, 711, 861. Chandra observations.
 Ghavamian *et al.* 2012, ApJ, 750, 39. Spitzer observations
 Zhariikov *et al.* 2013, A&A, 554, A120. IR of pulsar wind nebula.

G292.2–0.5**RA:** 11^h19^m20^s**1-GHz flux/Jy:** 7**Size/arcmin:** 20×15**Dec:** −61°28′**Spectral index:** 0.5**Type:** S**Radio:** Shell.**X-ray:** Shell, brighter to W, with central nebula.**Point sources:** Central, young pulsar.**Distance:** H α absorption indicates 8.4 kpc.**References:**

Camilo *et al.* 2000, ApJ, 541, 367. Pulsar detection.
 Crawford *et al.* 2001, ApJ, 554, 152. ATCA at 1.4 GHz ($29'' \times 25'' : S = 5.6 \times 0.3$ Jy) and 2.5 GHz ($21'' \times 20''$).
 Pivovarov *et al.* 2001, ApJ, 554, 161. ROSAT and ASCA observations.
 Gonzalez & Safi-Harb 2003, ApJ, 591, L143. Chandra observations of pulsar.
 Caswell *et al.* 2004, MNRAS, 352, 1405. ATCA at 5 GHz ($1' : S = 2.8$ Jy), including polarisation, and 1.4 GHz for H α absorption.
 Gonzalez & Safi-Harb 2005, ApJ, 619, 856. Chandra observations.
 Gonzalez *et al.* 2005, ApJ, 630, 489. XMM observations of pulsar.
 Safi-Harb *et al.* 2008, ApJ, 684, 532. Chandra observations of pulsar and nebula.
 Kumar *et al.* 2012, ApJ, 754, 96. Chandra and XMM observations.
 Ng *et al.* 2012, ApJ, 761, 65. XMM observations.

G293.8+0.6

RA: 11^h35^m00^s

Dec: -60°54'

1-GHz flux/Jy: 5?

Spectral index: 0.6?

Size/arcmin: 20

Type: C

Radio: Central source, with faint extended plateau.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3' : S = 9.0$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 2.1$ Jy).
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 51''$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 49'' : S = 2.6$ Jy).

G294.1-0.0

RA: 11^h36^m10^s

Dec: -61°38'

1-GHz flux/Jy: >2?

Spectral index: ?

Size/arcmin: 40

Type: S

Radio: Faint shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 49'' : S > 2$).

G296.1-0.5

RA: 11^h51^m10^s

Dec: -62°34'

1-GHz flux/Jy: 8?

Spectral index: 0.6?

Size/arcmin: 37×25

Type: S

Incorporates the previously catalogued remnant G296.1-0.7. Has been called G296.05-0.50.

Radio: Irregular shell, with nearby H α regions.

Optical: Detected.

X-ray: Irregular, incomplete shell.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3' : S = 6.9$ Jy) and Parkes 64-m at 5 GHz ($4' : S > 0.74$).
 Longmore *et al.* 1977, MNRAS, 181, 541. Optical spectra.
 van den Bergh 1978, ApJS, 38, 119. Optical observations.

Markert *et al.* 1981, ApJ, 248, L17. Einstein observations.
 Caswell & Barnes 1983, ApJ, 271, L55. Molonglo at 408 MHz ($3':S=12.4$ Jy).
 Bignami *et al.* 1986, ApJ, 302, 606. EXOSAT and Einstein observations.
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43''\times 48'' : S > 2.4$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Castro *et al.* 2011, ApJ, 734, 86. XMM observations.
 Gök & Sezer 2012, MNRAS, 419, 1603. Suzaku observations.

G296.5 + 10.0

PKS 1209–51/52

RA: 12^h09^m40^s**1-GHz flux/Jy:** 48**Size/arcmin:** 90×65**Dec:** −52°25′**Spectral index:** 0.5**Type:** S

Has been called G296.5+9.7.

Radio: Shell with two bright limbs.

Optical: Detected.

X-ray: Incomplete shell, with central pulsar.

Point sources: Central pulsar.

References:

Irvine & Irvine 1974, ApJ, 192, L111. Optical observations.
 Danziger & Dennefeld 1976, PASP, 88, 44. Optical spectra.
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
 Tuohy *et al.* 1979, ApJ, 230, L27. X-ray detection with HEAO-1 A2 experiment.
 Ruiz 1983, AJ, 88, 1210. Optical spectra.
 Dubner *et al.* 1986, AJ, 91, 343. Argentine 30-m dish at 1.4 GHz (34'), plus H_i.
 Kellett *et al.* 1987, MNRAS, 225, 199. EXOSAT of the west of the remnant, including the compact source.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44''\times 56''$).
 Matsui *et al.* 1988, ApJ, 329, 838. Einstein observations, including compact source.
 Roger *et al.* 1988, ApJ, 332, 940. MOST at 843 MHz ($44''\times 56''$).
 Bignami *et al.* 1992, ApJ, 389, L67. Optical in vicinity of X-ray source.
 Milne & Haynes 1994, MNRAS, 270, 106. Parkes 64-m at 2.4 GHz ($8'3 : S = 33\pm 3$ Jy), 4.8 GHz ($4'5 : S = 23.3\pm 3$ Jy) and 8.4 GHz ($3'0 : 18.8\pm 3$), including polarisation and review of flux densities.
 Mereghetti *et al.* 1996, ApJ, 464, 842. Radio, optical and X-ray observations of central source.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
 Vasisht *et al.* 1997, ApJ, 476, L43. ASCA observations of neutron star.
 Zavlin *et al.* 1998, A&A, 331, 821. ROSAT and ASCA observations of neutron star.
 Giacani *et al.* 2000, AJ, 119, 281. ATCA at 1.4 GHz ($2'7\times 4'0$) for H_i studies.
 Zavlin *et al.* 2000, ApJ, 540, L25. Chandra observations of central pulsar.
 Gotthelf & Halpern 2007, ApJ, 664, L35. X-ray timing observations of pulsar.
 Harvey-Smith *et al.* 2010, ApJ, 712, 1157. ATCA at 1.4 GHz ($1'8\times 3'3$), including polarisation.
 Araya *et al.* 2013, MNRAS, 434, 2202. Fermi observations.

G296.7–0.9**RA:** 11^h55^m30^s**Dec:** –63°08′**1-GHz flux/Jy:** 3**Spectral index:** 0.5**Size/arcmin:** 15×8**Type:** S**Radio:** Bilateral shell.**X-ray:** Brighter to SE.**References:**Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT observations.Robbins *et al.* 2012, MNRAS, 419, 2623. ATCA at 1.4 GHz (33″×45″: $S=2.5\pm0.2$ Jy), plus MOST at 843 MHz (43″×49″), plus other observations.

Prinz & Becker 2013, A&A, 550, A33. XMM observations.

G296.8–0.3**RA:** 11^h58^m30^s**Dec:** –62°35′**1-GHz flux/Jy:** 9**Spectral index:** 0.6**Size/arcmin:** 20×14**Type:** S

1156–62

Radio: Shell, brighter to the NW.**X-ray:** Detected.**Distance:** H_i absorption gives 9.6 kpc.**References:**Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′: $S=15.0$ Jy) and Parkes 64-m at 5 GHz (4′: $S=3.2$ Jy).

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′4) and 5 GHz (4′4).

Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×48″: $S=9.2$ Jy).Gaensler *et al.* 1998, MNRAS, 296, 813. ATCA at 1.3 GHz (22″×24″: $S=7.0\pm0.3$ Jy), including polarisation and H_i observations, plus review of flux densities.Sánchez-Ayaso *et al.* 2012, Ap&SS, 337, 573. XMM and IR observations.**G298.5–0.3****RA:** 12^h12^m40^s**Dec:** –62°52′**1-GHz flux/Jy:** 5?**Spectral index:** 0.4?**Size/arcmin:** 5?**Type:** ?**Radio:** Not well resolved, may be part of a larger ring?**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).

Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×48″: $S=1.8$ Jy).Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G298.6—0.0**RA:** 12^h13^m41^s**Dec:** -62°37'**1-GHz flux/Jy:** 5?**Spectral index:** 0.3**Size/arcmin:** 12×9**Type:** S

Has been called G298.6—0.1.

Radio: Incomplete shell, in complex region.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×50'').
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'' : S=7.4 Jy).
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G299.2—2.9**RA:** 12^h15^m13^s**Dec:** -65°30'**1-GHz flux/Jy:** 0.5?**Spectral index:** ?**Size/arcmin:** 18×11**Type:** S**Radio:** Faint source.**Optical:** Filaments in W.**X-ray:** Centrally brightened with shell at higher energies.**References:**

Busser *et al.* 1996, A&A, 310, L1. ROSAT detection, plus optical studies.
 Slane *et al.* 1996, ApJ, 465, 840. Einstein, IRAS and radio observations.
 Bai & Wang 2000, ApJ, 539, 760. ASCA observations.
 Park *et al.* 2007, ApJ, 665, 1173. Chandra observations.

G299.6—0.5**RA:** 12^h21^m45^s**Dec:** -63°09'**1-GHz flux/Jy:** 1.0?**Spectral index:** ?**Size/arcmin:** 13**Type:** S**Radio:** Faint shell, brightest to E.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'' : S=1.1 Jy).

G301.4—1.0**RA:** 12^h37^m55^s**Dec:** -63°49'**1-GHz flux/Jy:** 2.1?**Spectral index:** ?**Size/arcmin:** 37×23**Type:** S**Radio:** Faint, incomplete shell, with possible extension to southwest.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'' : S=2.3 Jy).

G302.3+0.7**RA:** 12^h45^m55^s**Dec:** -62°08′**1-GHz flux/Jy:** 5?**Spectral index:** 0.4?**Size/arcmin:** 17**Type:** S**Radio:** Distorted shell, in complex region, with possibly associated filament.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′:S=7.5 Jy) and Parkes 64-m at 5 GHz (4′:S=3.0 Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×49″:S=3.2 Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.**G304.6+0.1****RA:** 13^h05^m59^s**Dec:** -62°42′**1-GHz flux/Jy:** 14**Spectral index:** 0.5**Size/arcmin:** 8**Type:** S

Kes 17

Radio: Incomplete shell.**X-ray:** Detected.**Distance:** Possible limit of > 9.7 kpc from H_I absorption.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4′4″:S=6.9 Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×48″:S=18 Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.Combi *et al.* 2010, A&A, 523, A76. XMM observationsLee *et al.* 2011, ApJ, 740, 31. AKARI and Spitzer observations.Wu *et al.* 2011, ApJ, 740, L12. Fermi observations.Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.

Gök & Sezer 2012, MNRAS, 423, 1215. Suzaku observations.

Gelfand *et al.* 2013, ApJ, 777, 148. ATCA at 1.4 GHz (8″×23″:S=10.9±0.4 Jy), plus other observations.**G306.3-0.9****RA:** 13^h21^m50^s**Dec:** -63°34′**1-GHz flux/Jy:** 0.16?**Spectral index:** 0.5?**Size/arcmin:** 4**Type:** S?**Radio:** Diffuse emission.**X-ray:** Partial shell.**References:**Reynolds *et al.* 2013, ApJ, 766, 112. Chandra observations, and ATCA at 5.5 GHz (23″×26″), and 9 GHz.

G308.1–0.7**RA:** 13^h37^m37^s**Dec:** –63°04′**1-GHz flux/Jy:** 1.2?**Spectral index:** ?**Size/arcmin:** 13**Type:** S**Radio:** Faint shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'':S=1.3 Jy).

G308.4–1.4**RA:** 13^h41^m30^s**Dec:** –63°44′**1-GHz flux/Jy:** 0.4?**Spectral index:** ?**Size/arcmin:** 12×6?**Type:** S?

W part has been called G308.3–1.4.

Radio: Complex structure, with multiple arcs.**X-ray:** Limb brightened partial shell in W.**References:**Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT observations.

Prinz & Becker 2012, A&A, 544, A7. ATCA at 1.4 GHz (53''×64'':S=0.33 Jy) and 2.5 GHz (29''×35'':S=0.24 Jy), plus Chandra and other observations.

Hui *et al.* 2012, ApJ, 750, 7. XMM and other observations.De Horta *et al.* 2013, MNRAS, 428, 1980. ATCA at 1.4 GHz (54''×65'') and 2.5 GHz (29''×35''), plus other observations.**G308.8–0.1****RA:** 13^h42^m30^s**Dec:** –62°23′**1-GHz flux/Jy:** 15?**Spectral index:** 0.4?**Size/arcmin:** 30×20?**Type:** C?

Incorporates previous catalogued remnant G308.7+0.0.

Radio: Bright ridge in north, and arc to south.**Point sources:** Pulsar near centre of remnant.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3':S=16.7 Jy) and Parkes 64-m at 5 GHz (4':S=7.0 Jy).

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4'.4:S=6.5 Jy).

Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'').

Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.

Caswell *et al.* 1992, ApJ, 399, L151. MOST at 843 MHz (43''×49'').Kaspi *et al.* 1992, ApJ, 399, L155. Pulsar observations.Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G309.2–0.6**RA:** 13^h46^m31^s**1-GHz flux/Jy:** ??**Size/arcmin:** 15×12**Dec:** –62°54′**Spectral index:** 0.4?**Type:** S

Has been called G309.2–0.7.

Radio: Distorted shell.

X-ray: Extended emission, with unrelated central source.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : S = 10.0 Jy) and Parkes 64-m at 5 GHz (4′ : S = 3.9 Jy).

Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1′5).

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×50″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×48″ : S = 6 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Gaensler *et al.* 1998, MNRAS, 299, 812. ATCA at 1.3 GHz (24″×23″ : S = 5.2±0.2 Jy).

Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT observations.

Rakowski *et al.* 2006, ApJ, 649, L111. Observations of central source.

Safi-Harbi *et al.* 2007, ApJ, 659, 407. Observations of central source.

G309.8+0.0**RA:** 13^h50^m30^s**1-GHz flux/Jy:** 17**Size/arcmin:** 25×19**Dec:** –62°05′**Spectral index:** 0.5**Type:** S

Radio: Distorted shell.

Point sources: Steep radio spectrum source near the centre of the remnant.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : S = 26.4 Jy) and Parkes 64-m at 5 GHz (4′ : S = 7.4 Jy).

Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (1′).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×49″ : S > 8.8).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G310.6–1.6**RA:** 14^h00^m45^s**1-GHz flux/Jy:** ?**Size/arcmin:** 2.5**Dec:** –63°26′**Spectral index:** ?**Type:** C?

X-ray: Bright central nebula, with faint shell.

Point sources: X-ray pulsar.

References:

Renaud *et al.* 2010, ApJ, 716, 663. Chandra and other observations.

G310.6–0.3

Kes 20B

RA: 13^h58^m00^s**1-GHz flux/Jy:** 5?**Size/arcmin:** 8**Dec:** –62°09′**Spectral index:** ?**Type:** S**Radio:** Asymmetric shell.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'':*S*=5.4 Jy).Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.**G310.8–0.4**

Kes 20A

RA: 14^h00^m00^s**1-GHz flux/Jy:** 6?**Size/arcmin:** 12**Dec:** –62°17′**Spectral index:** ?**Type:** S**Radio:** Arc in E, in complex region.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'':*S*=6.9 Jy).Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.**G311.5–0.3****RA:** 14^h05^m38^s**1-GHz flux/Jy:** 3?**Size/arcmin:** 5**Dec:** –61°58′**Spectral index:** 0.5**Type:** S**Radio:** Shell, not well resolved.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3').

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'':*S*=2.9 Jy).Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.**G312.4–0.4****RA:** 14^h13^m00^s**1-GHz flux/Jy:** 45**Size/arcmin:** 38**Dec:** –61°44′**Spectral index:** 0.36**Type:** S**Radio:** Irregular, incomplete shell.**X-ray:** Weak emission in W.**Point sources:** Nearby γ -ray sources and pulsars.**Distance:** H α absorption suggests > 6 kpc and possibly > 14 kpc.**References:**Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3':*S*=56 Jy).Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'':*S*>19), plus Parkes 64-m at 4.5 GHz (*S*=30±2 Jy) and 8.55 GHz (*S*=17±4 Jy).Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Case & Bhattacharya 1999, ApJ, 521, 246. Nearby γ -ray sources.

Roberts *et al.* 1999, ApJ, 515, 712. MOST at 843 MHz ($43'' \times 49''$).

Doherty *et al.* 2003, MNRAS, 339, 1048. ATCA at 1.4 GHz ($25''$) plus H α absorption, and Chandra observations.

G312.5–3.0

RA: $14^{\text{h}}21^{\text{m}}00^{\text{s}}$

Dec: $-64^{\circ}12'$

1-GHz flux/Jy: 3.5?

Spectral index: ?

Size/arcmin: 20×18

Type: S

Radio: Distorted shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz ($10'.4$)

Kane & Vaughan 2003, MNRAS, 344, 625. ATCA at 1.4 GHz ($129'' \times 116''$) and 2.4 GHz ($75'' \times 67''$).

G315.1+2.7

RA: $14^{\text{h}}24^{\text{m}}30^{\text{s}}$

Dec: $-57^{\circ}50'$

1-GHz flux/Jy: ?

Spectral index: ?

Size/arcmin: 190×150

Type: S

Radio: Poorly defined shell.

Optical: Filaments, brighter in NE.

References:

Duncan *et al.* 1997, MNRAS, 287, 722. Parkes 64-m at 2.4 GHz ($11': S=19 \pm 3$ Jy).

Combi *et al.* 1998, A&A, 333, 298. Radio survey observations.

Stupar *et al.* 2007, MNRAS, 374, 1441. Optical and various radio observations.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G315.4–2.3

RA: $14^{\text{h}}43^{\text{m}}00^{\text{s}}$

Dec: $-62^{\circ}30'$

1-GHz flux/Jy: 49

Spectral index: 0.6

RCW 86, MSH 14–63

Size/arcmin: 42

Type: S

Possibly the remnant of the SN of AD185?

Radio: Shell, brightest to the SW.

Optical: Bright, radiative filaments, with some faint Balmer dominated filaments.

X-ray: Partial shell, with thermal and non-thermal emission.

Point sources: Several X-ray sources.

Distance: Optical observations imply 2.3 kpc.

References:

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz ($3': S=86$ Jy) and Parkes 64-m at 5 GHz ($4': S=18.2$ Jy).

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'.4$) and 5 GHz ($4'.4$).

Winkler 1978, ApJ, 221, 220. X-ray detection.

Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
 Piskunov *et al.* 1984, ApJ, 277, 710. Einstein observations
 Nugent *et al.* 1984, ApJ, 284, 612. X-ray spectrum.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 50'').
 Long & Blair 1990, ApJ, 358, L13. Balmer dominated optical filaments.
 Greidanus & Strom 1990, A&A, 240, 385. IRAS observations.
 Kaastra *et al.* 1992, A&A, 264, 654. Ginga X-ray spectra.
 Strom 1994, MNRAS, 268, L5. Historical association
 Chin & Huang 1994, Nature, 371, 398. Questioning of historical association.
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'' : $S > 22$).
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Smith *et al.* 1997, AJ, 114, 2664. Observations of Balmer dominated filaments.
 Vink *et al.* 1997, A&A, 328, 628. ASCA spectroscopy.
 Bocchino *et al.* 2000, A&A, 360, 671. BeppoSAX observations of N and SW.
 Bamba *et al.* 2000, PASJ, 52, 1157. ASCA observations.
 Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
 Dickel *et al.* 2001, ApJ, 546, 447. ATCA at 1.34 GHz (8''), including polarisation.
 Borkowski *et al.* 2001, ApJ, 550, 334. ASCA observations.
 Rho *et al.* 2002, ApJ, 581, 1116. Chandra observations.
 Gvaramadze & Vikhlinin 2003, A&A, 401, 625. Chandra point source search.
 Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
 Vink *et al.* 2006, ApJ, 648, L33. Chandra and XMM observations.
 Ueno *et al.* 2007, PASJ, 59, S171. Suzaku observations.
 Aharonian *et al.* 2009, ApJ, 692, 1500. γ -ray observations.
 Yamaguchi *et al.* 2011, PASJ, 63, S837. Suzaku observations.
 Helder *et al.* 2011, ApJ, 737, 85. XMM and optical observations.
 Williams *et al.* 2011, ApJ, 741, 96. Spitzer observations.
 Mignani *et al.* 2012, MNRAS, 425, 2309. X-ray and optical observations of compact X-ray sources.
 Lemoine-Goumard *et al.* 2012, A&A, 545, A28. γ -ray observations.
 Helder *et al.* 2013, MNRAS, 435, 910. Optical proper motion studies.
 Castro *et al.* 2013, ApJ, 779, 49. Chandra of NW.

G315.4–0.3

RA: 14^h35^m55^s

Dec: –60°36'

1-GHz flux/Jy: 8

Spectral index: 0.4

Size/arcmin: 24 \times 13

Type: ?

Radio: Irregular non-thermal emission, with H_{II} region superposed in E.

Optical: Detected.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : $S = 15.9$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 4.9$ Jy).
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'' : $S = 6.25$ Jy), re-assessment of earlier flux densities.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 49'' : $S = 3.1$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G315.9–0.0**RA:** 14^h38^m25^s**Dec:** –60°11′**1-GHz flux/Jy:** 0.8?**Spectral index:** ?**Size/arcmin:** 25×14**Type:** S

Has been called G315.8–0.0.

Radio: Faint, distorted shell, with elongated trail to pulsar.

Point sources: Pulsar at end of radio trail.

References:

Kesteven *et al.* 1987, AujPh, 40, 855. MOST at 843 MHz (44″×50″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×50″:S=0.9 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.

Camilo *et al.* 2009, ApJ, 703, L55. Pulsar detection.

Ng *et al.* 2012, ApJ, 746, 105. ATCA at 1.4, 2.5, 5 and 9 GHz, including polarisation, of pulsar trail.

G316.3–0.0

(MSH 14–57)

RA: 14^h41^m30^s**Dec:** –60°00′**1-GHz flux/Jy:** 20?**Spectral index:** 0.4**Size/arcmin:** 29×14**Type:** S

Radio: Distorted shell, with possible ‘blowout’.

X-ray: Detected.

Distance: H_i absorption data suggests > 7.2 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H_i absorption.

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4′4″:S=16.7 Jy).

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×51″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×50″:S=20 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

G317.3–0.2**RA:** 14^h49^m40^s**Dec:** –59°46′**1-GHz flux/Jy:** 4.7?**Spectral index:** ?**Size/arcmin:** 11**Type:** S

Radio: Incomplete shell.

Optical: Detected.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×50″:S=5.2 Jy).

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G318.2+0.1**RA:** 14^h54^m50^s**Dec:** -59°04'**1-GHz flux/Jy:** >3.9?**Spectral index:** ?**Size/arcmin:** 40×35**Type:** S**Radio:** Faint shell, with central H_{II} region.**X-ray:** Sources within remnant.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'' : S > 4.3).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Bocchino *et al.* 2001, A&A, 367, 629. BeppoSAX and ROSAT observations.**G318.9+0.4****RA:** 14^h58^m30^s**Dec:** -58°29'**1-GHz flux/Jy:** 4?**Spectral index:** 0.2?**Size/arcmin:** 30×14**Type:** C

May not be a SNR?

Radio: Complex arcs, with off-centre core.**References:**

Whiteoak 1990, Nature, 347, 157. MOST at 843 MHz (43''×51'').

Whiteoak 1993, ApJ, 415, 701. MOST at 843 MHz (43''×51'' : S = 4.8±0.6 Jy), Parkes 64-m at 4.5 GHz (4'.7 : S = 3.7±0.2 Jy) and 8.4 GHz (2'.8 : S = 3.0±0.4 Jy) including polarisation, and ATCA at 1.4 GHz and 4.8 GHz (11''×13'') of core.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'' : S = 4.8 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G320.4-1.2****RA:** 15^h14^m30^s**Dec:** -59°08'**1-GHz flux/Jy:** 60?**Spectral index:** 0.4

MSH 15-52, RCW 89

Size/arcmin: 35**Type:** C

Has been suggested as the remnant of the SN of AD185?

Radio: Ragged shell.**Optical:** RCW 89 is the H_α emitting region to the NW.**X-ray:** Partial shell, central nebula and pulsar and 'jet'.**Point sources:** Radio and X-ray pulsar, with wind nebula.**Distance:** H_I absorption indicates 5.2 kpc.**References:**Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).

Dopita *et al.* 1977, ApJ, 214, 179. Optical spectra.Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'').

Seward & Harnden 1982, ApJ, 256, L45. X-ray observations of pulsar.

Manchester *et al.* 1982, ApJ, 262, L31. Radio observations of pulsar.Seward *et al.* 1983, ApJ, 267, 698. X-ray, Optical and IR.

van den Bergh & Kamper 1984, ApJ, 280, L51. Optical expansion.

Seward *et al.* 1984, *Apj*, 281, 650. X-ray observations of pulsar and nebulosity.
 Lortet *et al.* 1987, *A&A*, 180, 65. Optical observations.
 Trussoni *et al.* 1990, *A&A*, 234, 403. EXOSAT observations.
 Asaoka & Koyama 1990, *PASJ*, 42, 625. Ginga X-ray spectrum.
 Arendt 1991, *AJ*, 101, 2160. IRAS observations, including compact source.
 Milne *et al.* 1993, *MNRAS*, 264, 853. Parkes 64-m at 4.8 GHz (4'5 : S = 37±7 Jy) and 8.4 GHz (3'0 : S = 24±4 Jy), including polarisation and review of flux densities.
 Strom 1994, *MNRAS*, 268, L5. Historical association.
 Chin & Huang 1994, *Nature*, 371, 398. Questioning of historical association.
 Matz *et al.* 1994, *Apj*, 434, 288. X-ray observations of pulsar.
 Schaefer 1995, *AJ*, 110, 1793. Questioning of historical association.
 Du Plessis *et al.* 1995, *Apj*, 453, 746. Hartesbeesthoek 26-m at 2.3, 5 and 8.5 GHz (S = 42, 35.6 and 14.5 Jy).
 Greiveldinger *et al.* 1995, *Apj*, 454, 855. ROSAT observations.
 Trussoni *et al.* 1996, *A&A*, 306, 581. ROSAT observations.
 Tamura *et al.* 1996, *PASJ*, 48, L33. ASCA observations.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×50'' : S = 62 Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Marsden *et al.* 1997, *Apj*, 491, L39. X-ray spectroscopy.
 Brazier & Becker 1997, *MNRAS*, 284, 335. ROSAT observations.
 Gaensler *et al.* 1999, *MNRAS*, 305, 724. ATCA at 1.4 GHz (24''×21''), plus H α observations, and 5.3 GHz (15''×10'').
 Sako *et al.* 2000, *Apj*, 537, 422. Possible high energy γ -ray detection of pulsar.
 Mineo *et al.* 2001, *A&A*, 380, 695. BeppoSAX observations.
 Gaensler *et al.* 2002, *Apj*, 569, 878. Chandra observations of pulsar and nebula.
 Dubner *et al.* 2002, *AJ*, 123, 337. ATCA at 1.4 GHz (4'0×2'7), plus H α observations.
 Aharonian *et al.* 2005, *A&A*, 435, L17. γ -ray detection.
 Yatsu *et al.* 2005, *Apj*, 631, 312. Chandra observations of pulsar and jet.
 DeLaney *et al.* 2006, *Apj*, 640, 929. Chandra and ROSAT multi-epoch observations of pulsar wind nebula.
 Kaplan & Moon 2006, *Apj*, 644, 1056. Possible IR detection of pulsar.
 Forot *et al.* 2006, *Apj*, 651, L45. X-ray observations of pulsar and nebula.
 Nakamori *et al.* 2008, *Apj*, 677, 297. γ -ray observations.
 Yatsu *et al.* 2009, *PASJ*, 61, 129. Chandra observations of pulsar wind nebula.
 Koo *et al.* 2011, *Apj*, 732, 6. AKARI and Spitzer observations.

G320.6–1.6

RA: 15^h17^m50^s

Dec: –59°16'

1-GHz flux/Jy: ?

Spectral index: ?

Size/arcmin: 60×30

Type: S

Radio: Faint shell, overlapping G320.4–1.2 in W.

Optical: Detected.

References:

Milne *et al.* 1993, *MNRAS*, 264, 853. Parkes 64-m at 4.8 GHz (4'5) and 8.4 GHz (3'0), including polarisation.
 Duncan *et al.* 1995, *MNRAS*, 277, 36. Parkes 64-m at 2.4-GHz (10'4).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×50'' : S > 9.3).
 Stupar & Parker 2011, *MNRAS*, 414, 2282. H α observations.

G321.9–1.1**RA:** 15^h23^m45^s**Dec:** –58°13′**1-GHz flux/Jy:** >3.4?**Spectral index:** ?**Size/arcmin:** 28**Type:** S**Radio:** Faint shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×51″:S>3.8).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G321.9–0.3****RA:** 15^h20^m40^s**Dec:** –57°34′**1-GHz flux/Jy:** 13**Spectral index:** 0.3**Size/arcmin:** 31×23**Type:** S**Radio:** Shell brighter to the W, with Cir X-1 to N.**Point sources:** Compact, probably thermal source at S edge.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′:S=18.3 Jy) and Parkes 64-m at 5 GHz (4′:S=7.8 Jy).

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×52″).

Stewart *et al.* 1993, MNRAS, 261, 593. ATCA at 1.5 GHz (21″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×51″:S>8.3).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Mignani *et al.* 2002, A&A, 386, 487. HST proper motion study of Cir X-1.Tudose *et al.* 2006, MNRAS, 372, 417. MOST at 843 MHz.**G322.1+0.0****RA:** 15^h20^m49^s**Dec:** –57°10′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 8×4.5?**Type:** S?**Radio:** Circular shell, with extension to S.**X-ray:** Diffuse emission.**Point sources:** Cir X-1 HMXB at centre.**References:**Heinz *et al.* 2013, ApJ, 779, 171. Chandra observations and ATCA at 1.1--3.1 GHz (4′.0×4′.9).**G322.5–0.1****RA:** 15^h23^m23^s**Dec:** –57°06′**1-GHz flux/Jy:** 1.5**Spectral index:** 0.4**Size/arcmin:** 15**Type:** C**Radio:** Shell with central extended source.**Point sources:** PN Pe 2-8 within boundary.**References:**

Whiteoak 1992, MNRAS, 256, 121. MOST at 843 MHz (43″×51″:S=2.0±0.3 Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 51''$: $S = 1.7$ Jy), plus Parkes 64-m at 4.5 GHz ($= 0.89 \pm 0.13$).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G323.5 + 0.1

RA: $15^{\text{h}}28^{\text{m}}42^{\text{s}}$

Dec: $-56^{\circ}21'$

1-GHz flux/Jy: 3?

Spectral index: 0.4?

Size/arcmin: 13

Type: S

Radio: Distorted shell, confused with thermal emission.

Point sources: Compact, probably thermal source near centre.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3'$: $S = 4.2$ Jy) and Parkes 64-m at 5 GHz ($4'$: $S = 1.5$ Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 52''$: $S = 4.2$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G326.3 – 1.8

RA: $15^{\text{h}}53^{\text{m}}00^{\text{s}}$

Dec: $-56^{\circ}10'$

1-GHz flux/Jy: 145

Spectral index: varies

Size/arcmin: 38

Type: C

MSH 15–56

Has been called G326.2–1.7.

Radio: Shell, with elongated, flat-spectrum core.

Optical: Emission around the shell.

X-ray: Shell, with central extended emission.

Point sources: Compact X-ray source.

References:

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz ($3'$: $S = 180$ Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.

Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz ($0'.8$: $S > 95$) and Parkes 64-m at 14.7 GHz ($2'.2$: $S = 69 \pm 8$ Jy).

van den Bergh 1979, ApJ, 227, 497. Optical observations.

Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.

Dennefeld 1980, PASP, 92, 603. Optical spectra.

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 53''$).

Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz ($43'' \times 52''$: $S = 153 \pm 40$ Jy), and Parkes 64-m at 8.4 GHz ($3'$: $S = 68 \pm 5$ Jy), including polarisation.

Seward 1990, ApJS, 73, 781. Einstein observations.

Kassim *et al.* 1993, ApJ, 419, 733. ROSAT image.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 52''$: $S > 130$).

Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Dickel *et al.* 2000, ApJ, 543, 840. ATCA at 1.34 GHz ($8''.5 \times 6''.4$: $S > 60$ Jy), 4.80 GHz ($3''.8$: $S = 25$ Jy for core only) and 8.64 GHz ($3''.2$: $S = 15$ Jy for core only).

Temim *et al.* 2013, ApJ, 768, 61. XMM and Chandra observations.

Yatsu *et al.* 2013, ApJ, 773, 25. XMM and Chandra observations.

G327.1–1.1**RA:** 15^h54^m25^s**Dec:** –55°09′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 18**Type:** C**Radio:** Shell, with off-centre core.**X-ray:** Diffuse, with core.**References:**

Milne & Dickel 1974, *AJPh*, 27, 549. Parkes 64-m at 2.7 GHz (8′.4 : S = 10 ± 15% Jy).
 Clark *et al.* 1975, *AJPA*, 37, 1. Molonglo at 408 MHz (3′ : S = 10.6 Jy) and Parkes 64-m at 5 GHz (4′ : S = 4.3 Jy).
 Lamb & Markert 1981, *Apj*, 244, 94. Einstein observations.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43″ × 52″ : S = 7.6 Jy).
 Seward *et al.* 1996, *Apj*, 471, 887. ROSAT observations.
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Sun *et al.* 1999, *Apj*, 511, 274. ASCA and ROSAT observations.
 Bocchino & Bandiera 2003, *A&A*, 398, 195. BeppoSAX observations.
 Temim *et al.* 2009, *Apj*, 691, 895. Chandra and XMM observations.

G327.2–0.1**RA:** 15^h50^m55^s**Dec:** –54°18′**1-GHz flux/Jy:** 0.4**Spectral index:** ?**Size/arcmin:** 5**Type:** S

Has been called G327.24–0.13.

Radio: Shell, possibly with central emission.**Point sources:** Central pulsar (magnetar).**References:**

Camilo *et al.* 2007, *Apj*, 666, L93. Pulsar observations.
 Gelfand & Gaensler 2007, *Apj*, 667, 1111. Chandra and XMM observations, and 843 GHz (43″ × 53″ : S = 0.5 ± 0.1 Jy) and SGPS at 1.4 GHz (S = 0.3 ± 0.1 Jy).

G327.4+0.4**RA:** 15^h48^m20^s**Dec:** –53°49′**1-GHz flux/Jy:** 30?**Spectral index:** 0.6**Size/arcmin:** 21**Type:** S

Kes 27

Has been called G327.3+0.4 and G327.3+0.5.

Radio: Incomplete, multi-arc shell, brightest to the SE.**X-ray:** Diffuse, best defined to E.**Distance:** H_I absorption indicates 4.3 to 5.4 kpc.**References:**

Caswell *et al.* 1975, *AJPA*, 37, 39. Molonglo at 408 MHz (3′ : S = 58 Jy) and Parkes 64-m at 5 GHz (4′ : S = 12.4 Jy).
 Dickel & Milne 1976, *AJPh*, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Lamb & Markert 1981, *Apj*, 244, 94. Einstein observations.
 Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44″ × 55″).

Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz ($43'' \times 53''$: $S = 32.2 \pm 6$ Jy), and Parkes 64-m at 8.4 GHz ($3'$: $S = 9.4 \pm 0.8$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 53''$: $S = 25$ Jy).
 Seward *et al.* 1996, ApJ, 471, 887. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ($2'.0 \times 1'.8$), plus HI.
 Enoguchi *et al.* 2002, PASJ, 54, 229. ASCA observations.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Chen *et al.* 2008, ApJ, 676, 1040. Chandra observations.

G327.4 + 1.0

RA: $15^{\text{h}}46^{\text{m}}48^{\text{s}}$

Dec: $-53^{\circ}20'$

1-GHz flux/Jy: 1.9?

Spectral index: ?

Size/arcmin: 14

Type: S

Radio: Asymmetric shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 54''$: $S = 2.1$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ($2'.0 \times 1'.8$), plus HI.

G327.6 + 14.6

RA: $15^{\text{h}}02^{\text{m}}50^{\text{s}}$

Dec: $-41^{\circ}56'$

1-GHz flux/Jy: 19

Spectral index: 0.6

SN1006, PKS 1459–41

Size/arcmin: 30

Type: S

This is the remnant of the SN of AD1006.

Radio: Shell, with two bright arcs.

Optical: Filaments to the NW, with broad H α component.

X-ray: Thermal shell, with non-thermal limb-brightened arcs.

Point sources: The background Schweizer--Middleditch star is near the middle of the remnant.

Distance: Optical spectra and proper motion indicate 2.2 kpc.

References:

van den Bergh 1976, ApJ, 208, L17. Optical observations.
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'.4$) and 5 GHz ($4'.4$).
 Becker *et al.* 1980, ApJ, 240, L33. X-ray spectroscopy.
 Schweizer & Middleditch 1980, ApJ, 241, 1039. Possible stellar remnant.
 Pye *et al.* 1981, MNRAS, 194, 569. Einstein observations.
 Caswell *et al.* 1983, MNRAS, 204, 921. FIRST at 1415 MHz ($77''$).
 Reynolds & Gilmore 1986, AJ, 92, 1138. VLA at 1.37 and 1.67 GHz ($16'' \times 20''$).
 Kirshner *et al.* 1987, ApJ, 315, L135. Broad H α optical component.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 66''$).
 Fesen *et al.* 1988, ApJ, 327, 164. UV absorption spectra of the Schweizer--Middleditch star.
 Roger *et al.* 1988, ApJ, 332, 940. MOST at 843 MHz ($44'' \times 66''$: $S = 17.5 \pm 1.5$ Jy).
 Long *et al.* 1988, ApJ, 333, 749. Optical proper motion for distance.
 Jones & Pye 1989, MNRAS, 238, 567. EXOSAT observations.
 Leahy *et al.* 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.

- Wu *et al.* 1993, ApJ, 416, 247. UV spectra.
- Reynolds & Gilmore 1993, AJ, 106, 272. VLA at 1.37 and 1.67 GHz (24''), including polarisation.
- Moffett *et al.* 1993, AJ, 106, 1566. VLA at 1.37 and 1.67 GHz (10'' \times 14''), for proper motion studies.
- Ozaki *et al.* 1994, PASJ, 46, 367. X-ray observations.
- Raymond *et al.* 1995, ApJ, 454, L31. Far UV spectroscopy.
- Koyama *et al.* 1995, Nature, 378, 255. ASCA observations.
- Willingale *et al.* 1996, MNRAS, 278, 749. ROSAT observations.
- Blair *et al.* 1996, ApJ, 468, 871. Optical absorption studies.
- Laming *et al.* 1996, ApJ, 472, 267. Modelling of optical spectra, including distance.
- Wu *et al.* 1997, ApJ, 477, L53. Far UV absorption spectra of the Schweizer--Middleditch star.
- Winkler & Long 1997, ApJ, 486, L137. UV absorption spectra of background quasar.
- Winkler & Long 1997, ApJ, 491, 829. ROSAT and optical images.
- Tanimori *et al.* 1998, ApJ, 497, L25. γ -ray detection.
- Vink *et al.* 2000, A&A, 354, 931. X-ray spectroscopy.
- Burleigh *et al.* 2000, A&A, 356, 585. Optical spectroscopy of the Schweizer--Middleditch star.
- Allen *et al.* 2001, ApJ, 558, 739. ASCA, ROSAT and other X-ray observations.
- Dubner *et al.* 2002, A&A, 387, 1047. ATCA at 1.4 GHz (4'.7 \times 3'.0) for H α , plus CO observations.
- Ghavamian *et al.* 2002, ApJ, 572, 888. Optical of filaments in NW.
- Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
- Winkler *et al.* 2003, ApJ, 585, 324. Optical proper motion studies.
- Vink *et al.* 2003, ApJ, 587, L31. XMM observations.
- Bamba *et al.* 2003, ApJ, 589, 827. Chandra observations of NE.
- Korreck *et al.* 2004, ApJ, 615, 280. Far-uv observations.
- Aharonian *et al.* 2005, A&A, 437, 135. γ -ray limit.
- Winkler *et al.* 2005, ApJ, 624, 189. HST absorption towards background sources.
- Kalemci *et al.* 2006, ApJ, 640, L55. γ -ray limit upper limit.
- Kalemci *et al.* 2006, ApJ, 644, 274. Integral observations.
- Acero *et al.* 2007, A&A, 475, 883. XMM observations.
- Raymond *et al.* 2007, ApJ, 659, 1257. HST H α observations.
- Hamilton *et al.* 2007, MNRAS, 381, 771. UV spectroscopy of the Schweizer--Middleditch star.
- Bamba *et al.* 2008, AdSpR, 41, 411. Suzaku observations.
- Cassam-Chenaï *et al.* 2008, ApJ, 680, 1180. ATCA and VLA at 1.5 GHz (6'' \times 9'') plus Chandra observations.
- Yamaguchi *et al.* 2008, PASJ, 60, S141. Suzaku observations.
- Bamba 2008, PASJ, 60, S153. Suzaku observations.
- Dyer *et al.* 2009, AJ, 137, 2956. GBT and VLA observations at 1.4 GHz.
- Winkler *et al.* 2011, ApJ, 742, 80. Multi-epoch UV spectroscopy of the Schweizer--Middleditch star.
- Miceli *et al.* 2012, A&A, 546, A66. XMM observations.
- Broersen *et al.* 2013, A&A, 552, A9. XMM observations of NW knot.
- Miceli *et al.* 2013, A&A, 550, A80. XMM observations.
- Katsuda *et al.* 2013, ApJ, 763, 85. Chandra proper motion study of NW.
- Winkler *et al.* 2013, ApJ, 764, 156. Spitzer observations.
- Uchida *et al.* 2013, ApJ, 771, 56. Suzaku observations.
- Reynoso *et al.* 2013, AJ, 145, 104. VLA and ATCA at 1.4 GHz (10''), including polarisation.
- Nikolić *et al.* 2013, Science, 340, 45. Optical spectroscopy.

G328.4+0.2

(MSH 15–57)

RA: 15^h55^m30^s**1-GHz flux/Jy:** 15**Size/arcmin:** 5**Dec:** –53°17′**Spectral index:** 0.0**Type:** F**Radio:** Amorphous emission, with central bar.**X-ray:** Detected at high energies.**Distance:** H_I absorption indicates > 17.4 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50′′).
 Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43′′×54′′: S=15 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Gaensler *et al.* 2000, ApJ, 542, 380. ATCA at 1.4 GHz (16′′5×19′′5: S=14.3±0.1 Jy) and 4.5 GHz (2′′0×1′′5: S=12.5±0.2 Jy).
 Hughes *et al.* 2000, ApJ, 542, 386. ASCA observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1′8×2′0), plus H_I.
 Johnston *et al.* 2004, MNRAS, 348, L19. ATCA at 19 GHz (6′′1×7′′7), including polarisation.
 Gelfand *et al.* 2007, ApJ, 663, 468. ATCA at 1.4 GHz (5′′8×7′′0: S=13.8±0.4 Jy), plus XMM observations.

G329.7+0.4**RA:** 16^h01^m20^s**1-GHz flux/Jy:** >34?**Size/arcmin:** 40×33**Dec:** –52°18′**Spectral index:** ?**Type:** S**Radio:** Diffuse shell, in complex region.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43′′×53′′: S>38).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2′0×1′8), plus H_I.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G330.0+15.0

Lupus Loop

RA: 15^h10^m00^s**1-GHz flux/Jy:** 350?**Size/arcmin:** 180?**Dec:** –40°00′**Spectral index:** 0.5?**Type:** S**Radio:** Low surface brightness loop with H_I shell.**X-ray:** Detected, with central source.**Point sources:** Central, possibly pulsating, X-ray source.**References:**

Milne 1971, AujPh, 24, 757. Parkes 64-m at 408 MHz (48′), 635 MHz (31′) and 1410 MHz (15′).
 Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (8′4: S=120±30% Jy).
 Toor 1980, A&A, 85, 184. X-ray image and spectrum.
 Colomb & Dubner 1982, A&A, 112, 141. Argentine 30-m dish at 1.42 GHz (30′), H_I observations.
 Leahy *et al.* 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.
 Ozaki *et al.* 1994, PASJ, 46, 367. X-ray observations.

Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
Shinn *et al.* 2006, ApJ, 644, L189. Far UV observations.

G330.2 + 1.0

RA: 16^h01^m06^s
Dec: -51°34′

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 11
Type: S?

Radio: Clumpy non-thermal emission, possibly a distorted shell.

X-ray: Shell.

Point sources: Central compact X-ray source.

Distance: H_I absorption indicates > 4.9 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: S=8.6 Jy) and Parkes 64-m at 5 GHz (4′: S=4.0 Jy).
Caswell *et al.* 1983, MNRAS, 204, 915. FIRST at 1415 MHz (52″×47″), and MOST at 843 MHz (43″×55″).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×55″: S=4.7 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2′.0×1′.8), plus H_I.
Torii *et al.* 2006, PASJ, 58, L11. ASCA detection.
Park *et al.* 2009, ApJ, 695, 431. XMM and Chandra observations.

G332.0 + 0.2

RA: 16^h13^m17^s
Dec: -50°53′

1-GHz flux/Jy: 8?
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Incomplete shell.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×57″).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×55″: S=8.9 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2′.0×1′.8), plus H_I.

G332.4 - 0.4

RCW 103

RA: 16^h17^m33^s
Dec: -51°02′

1-GHz flux/Jy: 28
Spectral index: 0.5

Size/arcmin: 10
Type: S

Radio: Shell, brightest to the S.

Optical: Filaments correspond well to the radio shell, brightest in SE.

X-ray: Brightest to NW, with point source near centre.

Point sources: Central, variable X-ray source, and nearby pulsar.

Distance: H_I absorption indicates 3.1 kpc.

References:

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1975, *A&A*, 45, 239. Parkes H α absorption.
 Tuohy & Garmire 1980, *ApJ*, 239, L107. Einstein detection of X-ray point source.
 Caswell *et al.* 1980, *MNRAS*, 190, 881. FIRST at 1415 MHz (50'').
 Lamb & Markert 1981, *ApJ*, 244, 94. Einstein observations.
 Ruiz 1983, *AJ*, 88, 1210. Optical spectra.
 Tuohy *et al.* 1983, *ApJ*, 268, 778. X-ray observations.
 Leibowitz & Danziger 1983, *MNRAS*, 204, 273. Optical spectra.
 Nugent *et al.* 1984, *ApJ*, 284, 612. X-ray spectrum (and Einstein image from Tuohy, private communication).
 Meaburn & Allen 1986, *MNRAS*, 222, 593. Optical spectra.
 Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44'' \times 57'').
 Oliva *et al.* 1990, *A&A*, 240, 453. IR spectroscopy.
 Dickel *et al.* 1996, *AJ*, 111, 340. ATCA at 1.36 (8'') and 2.37 GHz (4''5''), including polarisation.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' \times 55'': $S=34$ Jy).
 Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant.
 Gotthelf *et al.* 1997, *ApJ*, 487, L175. ASCA observations of compact X-ray source.
 Carter *et al.* 1997, *PASP*, 109, 990. Optical expansion.
 Torii *et al.* 1998, *ApJ*, 494, L207. ASCA detection of nearby pulsar.
 Kaspi *et al.* 1998, *ApJ*, 503, L161. Pulsar observations.
 Gotthelf *et al.* 1999, *ApJ*, 514, L107. X-ray variability of central source.
 Oliva *et al.* 1999, *A&A*, 343, 943. ISO spectroscopy.
 Torii *et al.* 2000, *ApJ*, 534, L71. X-ray timing on pulsar, including glitch.
 Reynoso *et al.* 2004, *PASA*, 21, 82. ATCA at 1.4 GHz (50''), including H α absorption to central source.
 Russeil *et al.* 2005, *A&A*, 429, 497. H α observations.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Paron 2006, *PASA*, 23, 69. CO and HCO $^+$ observations of surroundings.
 De Luca 2006, *Science*, 313, 814. XMM observations of central source.
 De Luca 2007, *Ap&SS*, 308, 231. XMM observation of periodicity of central source.
 Matsumoto *et al.* 2007, *PASJ*, 59, S199. Suzaku observations.
 De Luca 2008, *ApJ*, 682, 1185. IR observations of central source.
 Anderson *et al.* 2011, *ApJ*, 742, 7. Spitzer observations.

G332.4 + 0.1

MSH 16–51, Kes 32

RA: 16^h15^m20^s**1-GHz flux/Jy:** 26**Size/arcmin:** 15**Dec:** –50°42'**Spectral index:** 0.5**Type:** S

Has been called G332.4+0.2.

Radio: Distorted shell, with thermal jet and plume adjacent.

Optical: Detected.

X-ray: Shell, brightest to NW.

Point sources: Pulsar nearby.

References:

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Roger *et al.* 1985, *Nature*, 316, 44. MOST at 843 MHz (44'' \times 57'').
 Caraveo 1993, *ApJ*, 415, L111. Nearby pulsar.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' \times 56'': $S=29$ Jy).
 Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant.
 Brinkmann *et al.* 1999, *A&A*, 346, 599. ROSAT image of remnant and nearby pulsar.
 Vink *et al.* 2004, *ApJ*, 604, 693. Chandra observations.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer possible detection.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G332.5–5.6

RA: 16^h43^m20^s

Dec: –54°30′

1-GHz flux/Jy: 2?

Spectral index: 0.7?

Size/arcmin: 35

Type: S

Radio: Bipolar shell, with central emission also.

Optical: Patchy filaments.

X-ray: Emission from centre.

References:

Reynoso & Green 2007, MNRAS, 375, 92. ATCA at 1.4 GHz (40″ : $S = 1.90 \pm 0.15$ Jy) and 2.4 GHz (90″ : $S = 1.3 \pm 0.2$ Jy) including polarisation, and 843 MHz (43″ × 53″) plus ROSAT observations.

Stupar *et al.* 2007, MNRAS, 381, 377. Optical observations, plus ATCA at 1.4 GHz (58″) and 2.4 GHz (95″) including polarisation, and other observations.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G335.2+0.1

RA: 16^h27^m45^s

Dec: –48°47′

1-GHz flux/Jy: 16

Spectral index: 0.5

Size/arcmin: 21

Type: S

Radio: Well defined shell.

Point sources: Old pulsar within remnant boundary.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : $S = 27.1$ Jy) and Parkes 64-m at 5 GHz (4′ : $S = 8.6$ Jy).

Kaspi *et al.* 1996, AJ, 111, 2028. Pulsar observations.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ × 57″ : $S = 16$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G336.7+0.5

RA: 16^h32^m11^s

Dec: –47°19′

1-GHz flux/Jy: 6

Spectral index: 0.5

Size/arcmin: 14 × 10

Type: S

Radio: Irregular shell.

Optical: Detected.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″ × 60″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ × 58″ : $S = 6.1$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G337.0–0.1

(CTB 33)

RA: 16^h35^m57^s**1-GHz flux/Jy:** 1.5**Size/arcmin:** 1.5**Dec:** –47°36′**Spectral index:** 0.6?**Type:** S

This entry refers to a small (1'5) SNR, not the larger previously catalogued G337.0–0.1. Has mistakenly been called G337.7–0.1.

Radio: Shell, in a complex region.

Point sources: Associated with a soft gamma repeater.

Distance: Association with CTB 33 gives 11 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×58'' : S=21 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Sarma *et al.* 1997, ApJ, 483, 335. ATCA at 1.4 GHz (12'') including H_I, and 8.9 GHz (13''×15'') for recombination lines, clarifying extent of the remnant.
 Woods *et al.* 1999, ApJ, 519, L139. Soft gamma repeater observations.
 Hurley *et al.* 2000, ApJ, 528, L21. ASCA observations of soft gamma repeater.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Castro *et al.* 2013, ApJ, 774, 36. Fermi observations.

G337.2–0.7**RA:** 16^h39^m28^s**1-GHz flux/Jy:** 1.5**Size/arcmin:** 6**Dec:** –47°51′**Spectral index:** 0.4**Type:** S

Radio: Shell, brighter in S.

X-ray: Extended emission.

Distance: H_I absorption suggests 2.0 to 9.3 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : S=3.8 Jy) and Parkes 64-m at 5 GHz (4' : S=0.70 Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×58'' : S=2.0 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT observations.
 Rakowski *et al.* 2006, ApJ, 646, 982. ATCA observations at 1.3 GHz (15'' : S=1.55±0.05 Jy) and 5 GHz (15'' : S=0.93±0.02 Jy), plus H_I and Chandra observations.

G337.2+0.1**RA:** 16^h35^m55^s**1-GHz flux/Jy:** 1.5?**Size/arcmin:** 3×2**Dec:** –47°20′**Spectral index:** ?**Type:** ?

Radio: Not well defined.

X-ray: Detected.

Distance: Association with H_I hole gives 14 kpc.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×59'' : S=1.6 Jy)

Combi *et al.* 2005, A&A, 431, L9. ASCA and other observations.
 Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations of nearby source.
 Combi *et al.* 2006, ApJ, 653, L41. XMM observations.

G337.3+1.0

Kes 40

RA: 16^h32^m39^s**1-GHz flux/Jy:** 16**Size/arcmin:** 15×12**Dec:** −46°36′**Spectral index:** 0.55**Type:** S**Radio:** Nearly complete shell.**References:**

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′:S=24.6 Jy) and Parkes 64-m at 5 GHz (4′:S=7.2 Jy).
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′4) and 5 GHz (4′4).
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×61″).
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43″×59″:S=14.8±3.0 Jy), and Parkes 64-m at 8.4 GHz (3′:S=5.1±0.6 Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×59″:S=20 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G337.8−0.1

Kes 41

RA: 16^h39^m01^s**1-GHz flux/Jy:** 18**Size/arcmin:** 9×6**Dec:** −46°59′**Spectral index:** 0.5**Type:** S**Radio:** Distorted shell.**X-ray:** Centrally brightened.**Distance:** H_i absorption suggests 11 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes H_i absorption.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×59″:S=18 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
 Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz, for associated OH masers.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Köthes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including H_i.
 Combi *et al.* 2008, A&A, 488, L25. XMM observations.

G338.1+0.4
RA: 16^h37^m59^s**1-GHz flux/Jy:** 4?**Size/arcmin:** 15?**Dec:** −46°24′**Spectral index:** 0.4**Type:** S**Radio:** Arc in NE, merging with thermal emission in S.**Optical:** Detected.**X-ray:** Detected.**References:**

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Zealey *et al.* 1979, *A&AS*, 38, 39. Optical detection.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' \times 59'' : $S=3.8$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
 Caswell 2004, *MNRAS*, 349, 99. ATCA at 1.7 GHz of associated OH masers.

G338.3–0.0

RA: 16^h41^m00^s
Dec: –46°34'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 8
Type: C?

Radio: Irregular shell, in complex region.

X-ray: Central X-ray source and nebula.

Point sources: Central X-ray source.

Distance: H_i observations suggest 8 to 13 kpc.

References:

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' \times 59'' : $S=7.4$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Aharonian *et al.* 2005, *Science*, 307, 1938. γ -ray detection.
 Aharonian *et al.* 2006, *ApJ*, 636, 777. γ -ray observations.
 Landi *et al.* 2006, *ApJ*, 651, 190. X-ray observations.
 Funk *et al.* 2007, *ApJ*, 662, 517. XMM observations.
 Lemiere *et al.* 2009, *ApJ*, 706, 1269. Chandra observations of central source and nebula.
 Slane *et al.* 2010, *ApJ*, 720, 266. Fermi observations.
 Castelletti *et al.* 2011, *A&A*, 536, A98. GMRT at 235 MHz (10'' \times 26''), 610 MHz (5''.0 \times 12''.6) and 1280 MHz (5''.0 \times 6''.4), plus ATCA at 2.3 GHz (4''.3 \times 4''.8).

G338.5+0.1

RA: 16^h41^m09^s
Dec: –46°19'

1-GHz flux/Jy: 12?
Spectral index: ?

Size/arcmin: 9
Type: ?

Radio: Circle of non-thermal emission in complex region, not well defined.

Distance: H_i absorption suggests 11 kpc.

References:

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' \times 59'' : $S=13$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Kothes & Dougherty 2007, *A&A*, 468, 993. SGPS at 1.4 GHz including H_i.

G340.4 + 0.4**RA:** 16^h46^m31^s**Dec:** -44°39'**1-GHz flux/Jy:** 5**Spectral index:** 0.4**Size/arcmin:** 10×7**Type:** S**Radio:** Distorted shell, elongated east--west.**Optical:** Detected.**References:**

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3': S=8.2 Jy) and Parkes 64-m at 5 GHz (4': S=2.9 Jy).

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50").

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (75"×48" : S = 9.8±0.9 Jy) and 1.4 GHz (27"×9" : S = 3.6±0.1 Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43"×61" : S = 5.9 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G340.6 + 0.3**RA:** 16^h47^m41^s**Dec:** -44°34'**1-GHz flux/Jy:** 5?**Spectral index:** 0.4?**Size/arcmin:** 6**Type:** S**Radio:** Incomplete shell.**Optical:** Possible associated filaments.**Distance:** H α absorption suggests 15 kpc.**References:**

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3': S=7.0 Jy) and Parkes 64-m at 5 GHz (4': S=2.8 Jy).

Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50").

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (75"×48" : S = 9.2±0.9 Jy) and 1.4 GHz (27"×9" : S = 5.8±0.1 Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43"×61" : S = 4.5 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including H α .

G341.2 + 0.9**RA:** 16^h47^m35^s**Dec:** -43°47'**1-GHz flux/Jy:** 1.5?**Spectral index:** 0.6?**Size/arcmin:** 22×16**Type:** C**Radio:** Incomplete shell, with extension to SW.**Point sources:** Pulsar in W, with wind nebula.**References:**

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (54"×116" : S = 3.0±0.1 Jy) and 1.4 GHz (21"×25" : S = 12.5±0.05 Jy).

Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4 and 4.9 GHz (25") of pulsar wind nebula.

G341.9–0.3**RA:** 16^h55^m01^s
Dec: –44°01′**1-GHz flux/Jy:** 2.5
Spectral index: 0.5**Size/arcmin:** 7
Type: S**Radio:** Incomplete shell, brightest to NE.**References:**

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′: $S=7.4$ Jy) and Parkes 64-m at 5 GHz (4′: $S=1.7$ Jy).

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50″), revision of previous flux densities.

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (35″×11″: $S=2.2\pm 0.1$ Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×62″: $S=2.7$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.0–0.2**RA:** 16^h54^m50^s
Dec: –43°53′**1-GHz flux/Jy:** 3.5?
Spectral index: 0.4?**Size/arcmin:** 12×9
Type: S**Radio:** Distorted shell.**References:**

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50″), estimate $S_{408\text{ MHz}}=5$ Jy, $S_{5\text{ GHz}}=2$ Jy from previous maps.

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (35″×11″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×62″: $S=3.5$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.1+0.9**RA:** 16^h50^m43^s
Dec: –43°04′**1-GHz flux/Jy:** 0.5?
Spectral index: ?**Size/arcmin:** 10×9
Type: S**Radio:** Incomplete shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×63″: $S=0.6$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G343.0–6.0

RCW 114

RA: 17^h25^m00^s**1-GHz flux/Jy:** ?**Size/arcmin:** 250**Dec:** –46°30′**Spectral index:** ?**Type:** S**Radio:** Faint, poorly defined.**Optical:** Filamentary shell.**References:**

Walker & Zealey 2001, MNRAS, 325, 287. Optical observations, and review of earlier observations.

Welsh *et al.* 2003, A&A, 403, 605. Optical spectroscopy.Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.Kim *et al.* 2010, ApJ, 709, 823. UV observations.**G343.1–2.3****RA:** 17^h08^m00^s**1-GHz flux/Jy:** 8?**Size/arcmin:** 32?**Dec:** –44°16′**Spectral index:** 0.5?**Type:** C?**Radio:** Incomplete shell?**X-ray:** Pulsar wind nebula.**Point sources:** Pulsar near edge, with wind nebula.**References:**McAdam *et al.* 1993, Nature, 361, 516. MOST at 843 MHz (smoothed to 2′).Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (56″×111″: $S=10.6$ Jy) and 1.4 GHz (22″×27″) near pulsar.Becker *et al.* 1995, A&A, 298, 528. ROSAT of pulsar, and limit for remnant.Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4, 4.9 and 8.5 GHz (25″) of pulsar wind nebula.Gotthelf *et al.* 2002, ApJ, 567, L125. Pulsar x-ray detection.

Dodson & Golap 2002, MNRAS, 334, L1. ATCA at 1.4 GHz (70″×47″) including polarisation, and Chandra observations of pulsar wind nebula.

Aharonian *et al.* 2005, A&A, 432, L9. γ -ray limit.Abramowski *et al.* 2011, A&A, 528, A143. γ -ray observations.**G343.1–0.7****RA:** 17^h00^m25^s**1-GHz flux/Jy:** 7.8**Size/arcmin:** 27×21**Dec:** –43°14′**Spectral index:** 0.55**Type:** S**Radio:** Shell, with smaller thermal shell adjacent.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×63″: $S=58.5\pm 0.6$), plus Parkes 64-m at 4.5 GHz ($S=3.9\pm 0.6$ Jy) and 8.55 GHz ($S=2.4\pm 0.5$ Jy)Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G344.7–0.1**RA:** 17^h03^m51^s**Dec:** –41°42′**1-GHz flux/Jy:** 2.5?**Spectral index:** 0.3?**Size/arcmin:** 8**Type:** C?**Radio:** Asymmetric shell, with possible core.**X-ray:** Detected.**Distance:** H_i absorption and association with features suggests 6.3 kpc.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=4.7$ Jy) and Parkes 64-m at 5 GHz (4′: $S=1.3$ Jy).Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (30″×43″: $S=1.7\pm0.1$ Jy).Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×65″: $S=2.5$ Jy).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.Yamauchi *et al.* 2005, PASJ, 57, 459. ASCA observations.Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.Giacani *et al.* 2011, A&A, 531, A138. VLA and ATCA at 1.4 GHz (5″6×8″), ATCA at 5 GHz (10″×13″) and 8.4 GHz (5″×10″), plus XMM observations and H_i from SGPS.Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.Yamaguchi *et al.* 2012, ApJ, 749, 137. Suzaku observations.Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.**G345.7–0.2****RA:** 17^h07^m20^s**Dec:** –40°53′**1-GHz flux/Jy:** 0.6?**Spectral index:** ?**Size/arcmin:** 6**Type:** S**Radio:** Poorly defined diffuse shell.**Point sources:** Old pulsar nearby.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×66″: $S=0.7$ Jy).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G346.6–0.2****RA:** 17^h10^m19^s**Dec:** –40°11′**1-GHz flux/Jy:** 8?**Spectral index:** 0.5?**Size/arcmin:** 8**Type:** S**Radio:** Irregular shell.**X-ray:** Centrally brightened.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=14.9$ Jy) and Parkes 64-m at 5 GHz (4′: $S=4.3$ Jy).Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (31″×43″: $S=8.1\pm0.9$ Jy).Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×67″: $S=8.7$ Jy).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Sezer *et al.* 2011, MNRAS, 415, 301. Suzaku observations.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Yamauchi *et al.* 2013, PASJ, 65, 6. Suzaku observations.

G347.3–0.5
RX J1713.7–3946
RA: 17^h13^m50^s
1-GHz flux/Jy: 30?

Size/arcmin: 65×55

Dec: –39°45′

Spectral index: ?

Type: S?

Radio: Faint emission.

X-ray: Non-thermal, limb-brightened to W, with central source.

Point sources: Central X-ray source.

Distance: Association with molecular clouds and X-ray observations imply 1.3 kpc.

References:

- Koyama *et al.* 1997, PASJ, 49, L7. ASCA of NW.
 Slane *et al.* 1999, ApJ, 525, 357. ASCA and other observations.
 Muraishi *et al.* 2000, A&A, 354, L57. High energy γ -ray detection.
 Butt *et al.* 2001, ApJ, 562, L167. Associated γ -ray emission.
 Ellison *et al.* 2001, ApJ, 563, 191. ATCA at 1.4 GHz (46''×36''), and ASCA observations.
 Enomoto *et al.* 2002, Nature, 416, 823. γ -ray observations.
 Uchiyama *et al.* 2002, PASJ, 54, L73. ASCA observations.
 Uchiyama *et al.* 2003, A&A, 400, 567. Chandra spectroscopy.
 Pannuti *et al.* 2003, ApJ, 593, 377. ROSAT and ASCA observations.
 Lazendic *et al.* 2003, ApJ, 593, L27. Chandra, XMM and other X-ray observations of central source.
 Fukui *et al.* 2003, PASJ, 55, L61. CO observations of surroundings.
 Cassam-Chenaï *et al.* 2004, A&A, 427, 199. XMM and other observations.
 Aharonian *et al.* 2004, Nature, 432, 75. γ -ray detection.
 Lazendic *et al.* 2004, ApJ, 602, 271. Chandra observations of parts, and ATCA at 1.4 GHz (36''×46'').
- Hiraga *et al.* 2005, A&A, 431, 953. XMM observations,
 Moriguchi *et al.* 2005, ApJ, 631, 947. CO observations of surroundings.
 Aharonian *et al.* 2007, A&A, 464, 235. γ -ray observations.
see also: Aharonian *et al.* 2011, A&A, 531, C1. Erratum.
 Uchiyama *et al.* 2007, Nature, 449, 576. Chandra multi-epoch observations for study of small-scale variability.
 Tanaka *et al.* 2008, ApJ, 685, 988. Suzaku observations.
 Takahashi *et al.* 2008, PASJ, 60, S131. Suzaku observations of SW.
 Mignani *et al.* 2008, A&A, 484, 457. Optical and IR observations of central source.
 Acero *et al.* 2009, A&A, 505, 157. XMM observations, plus other radio, IR and γ -ray data.
 Sano *et al.* 2010, ApJ, 724, 59. CO observations of region.
 Maxted *et al.* 2012, MNRAS, 422, 2230. Molecular line observations.
see also: Maxted *et al.* 2013, MNRAS, 430, 2511. Erratum.
 Sano *et al.* 2013, ApJ, 778, 59. CO, SGPS H_I and Suzaku observations.

G348.5—0.0**RA:** 17^h15^m26^s**1-GHz flux/Jy:** 10?**Size/arcmin:** 10?**Dec:** -38°28'**Spectral index:** 0.4?**Type:** S?**Radio:** Arc, overlapping G348.5+0.1.**References:**

Kassim *et al.* 1991, Apj, 374, 212. VLA at 333 MHz (46''×53''), 1.4 GHz (18''×33'') and part at 5 GHz (2''×3''9).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×69'':S=10.2 Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Hewitt *et al.* 2008, Apj, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Hewitt *et al.* 2009, Apj, 694, 1266. Spitzer spectroscopy.

Anderson *et al.* 2011, Apj, 742, 7. Spitzer observations.

G348.5+0.1

CTB 37A

RA: 17^h14^m06^s**1-GHz flux/Jy:** 72**Size/arcmin:** 15**Dec:** -38°32'**Spectral index:** 0.3**Type:** S**Radio:** Shell, poorly define to S and W, overlapping G348.5—0.0 in E.**X-ray:** Brighter to W.**Distance:** H_I absorption indicates 8.0 kpc.**References:**

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3':S=97 Jy).

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4'.4:S=43 Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.

Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz (0'.8:S>50) and Parkes 64-m at 14.7 GHz (2'.2:S=18±5 Jy).

Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20''×45'').

Kassim *et al.* 1991, Apj, 374, 212. VLA at 333 MHz (46''×53''), 1.4 GHz (18''×33'') and part at 5 GHz (2''×3''9).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×69'':S=71 Jy).

Brogan *et al.* 2000, Apj, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

Reynoso & Mangum 2000, Apj, 545, 874. CO observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Aharonian *et al.* 2008, A&A, 490, 685. γ -ray observations.

Castro & Slane 2010, Apj, 717, 372. Fermi observations.

Anderson *et al.* 2011, Apj, 742, 7. Spitzer observations.

Sezer *et al.* 2011, MNRAS, 417, 1387. Suzaku observations.

Tian & Leahy 2012, MNRAS, 421, 2593. SGPS H_I absorption observations.

Maxted *et al.* 2013, MNRAS, 434, 2188. Molecular line observations of region.

G348.7 + 0.3

CTB 37B

RA: 17^h13^m55^s**1-GHz flux/Jy:** 26**Size/arcmin:** 17?**Dec:** -38°11'**Spectral index:** 0.3**Type:** S**Radio:** Incomplete shell with faint eastern extensions.**X-ray:** Diffuse emission.**Point sources:** X-ray pulsar.**Distance:** H_I absorption suggests 13 kpc.**References:**Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3': S=34 Jy).

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4'.4: S=32 Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz (0'.8: S>20) and Parkes 64-m at 14.7 GHz (2'.2: S=8±3 Jy).

Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20''×45'').

Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz (46''×53'').

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×69'': S=33 Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.Aharonian *et al.* 2008, A&A, 486, 829. γ -ray and Chandra observations.Nakamura *et al.* 2009, PASJ, 61, S197. Suzaku and Chandra observations.

Halpern & Gotthelf 2010, ApJ, 710, 941. Chandra detection of pulsar.

Halpern & Gotthelf 2010, ApJ, 725, 1384. Chandra observations of pulsar.

Tian & Leahy 2012, MNRAS, 421, 2593. SGPS H_I observations.**G349.2 - 0.1****RA:** 17^h17^m15^s**1-GHz flux/Jy:** 1.4?**Size/arcmin:** 9×6**Dec:** -38°04'**Spectral index:** ?**Type:** S**Radio:** Elongated shell, adjacent to bright H_{II} region.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×70'': S=1.6 Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.**G349.7 + 0.2****RA:** 17^h17^m59^s**1-GHz flux/Jy:** 20**Size/arcmin:** 2.5×2**Dec:** -37°26'**Spectral index:** 0.5**Type:** S**Radio:** Incomplete clumpy shell, with enhancement to the S.**X-ray:** Irregular shell, brighter to S and E.**Distance:** H_I absorption indicates 14.8 kpc, association with OH features gives 22 kpc.**References:**

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3': S=31.0 Jy) and Parkes 64-m at 5 GHz (4': S=9.1 Jy), no maps.

Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.
 Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz ($3''4 \times 14''5$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 71''$; $S=22$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Reynoso & Mangum 2001, AJ, 121, 347. CO observations of the vicinity.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Slane *et al.* 2002, ApJ, 580, 904. ASCA observations.
 Dubner *et al.* 2004, A&A, 426, 201. CO observations of surroundings.
 Lazendic *et al.* 2005, ApJ, 618, 733. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189.GBT at 1.6 and 1.7 GHz for OH masers.
 Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Castro & Slane 2010, ApJ, 717, 372. Fermi observations.
 Lazendic *et al.* 2010, MNRAS, 409, 371. OH, CO and other molecular line observations of region.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.

G350.0–2.0

RA: 17^h27^m50^s
Dec: –38°32'

1-GHz flux/Jy: 26
Spectral index: 0.4

Size/arcmin: 45
Type: S

Incorporates the previously catalogued G350.0–1.8 in the NW.

Radio: Shell, brightest in NW.

Optical: Detected.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3'$: $S=49.5$ Jy) and Parkes 64-m at 5 GHz ($4'$: $S=13.6$ Jy).
 Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz ($4'4$).
 Gaensler 1998, ApJ, 493, 781. VLA and Parkes 64-m at 1.4 GHz ($18'' \times 21''$: $S=22.3 \pm 0.3$ Jy), clarifying extent of remnant.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G350.1–0.3

RA: 17^h17^m40^s
Dec: –37°24'

1-GHz flux/Jy: 6?
Spectral index: 0.8?

Size/arcmin: 4?
Type: ?

Radio: Several clumps of emission.

X-ray: Diffuse emission, with compact source.

Point sources: X-ray source.

Distance: H α absorption indicates 4.5 to 10.7 kpc, possible interaction with molecular cloud indicates 4.5 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3'$: $S=10.7$) and Parkes 64-m at 5 GHz ($4'$: $S=1.7$).
 Salter *et al.* 1986, A&A, 162, 217. VLA at 1.5 ($4'' \times 10''4$) and 4.8 GHz ($15'' \times 35''$) and 15 GHz ($5'' \times 2''$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 71''$).
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.

Gaensler *et al.* 2008, ApJ, 680, L37. VLA at 4.8 GHz ($11''.4 \times 5''.6$) plus XMM observations.
 Lovchinsky *et al.* 2011, ApJ, 731, 70. Chandra and Spitzer observations.

G351.2+0.1

RA: $17^{\text{h}}22^{\text{m}}27^{\text{s}}$
Dec: $-36^{\circ}11'$

1-GHz flux/Jy: 5?
Spectral index: 0.4

Size/arcmin: 7
Type: C?

Has been called G351.3+0.2.

Radio: Distorted shell, with possible flat-spectrum core.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz ($3': S=8.1$ Jy) and Parkes 64-m at 5 GHz ($4': S=3.1$ Jy).
 Becker & Helfand 1988, AJ, 95, 883. VLA at 5 GHz ($15''$), and at 15 GHz of core.
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($32'' \times 36'' : S=4.8 \pm 0.2$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 73'' : S=5.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G351.7+0.8

RA: $17^{\text{h}}21^{\text{m}}00^{\text{s}}$
Dec: $-35^{\circ}27'$

1-GHz flux/Jy: 10
Spectral index: 0.5?

Size/arcmin: 18×14
Type: S

Radio: Elongated shell, adjacent to bright H_{II} region.

Point sources: Pulsar nearby.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 74'' : S=11$ Jy).
 Tian *et al.* 2007, MNRAS, 378, 1283. SGPS at 1.4 GHz ($100'' : S=8.4 \pm 0.7$ Jy) including H_I.

G351.9-0.9

RA: $17^{\text{h}}28^{\text{m}}52^{\text{s}}$
Dec: $-36^{\circ}16'$

1-GHz flux/Jy: 1.8?
Spectral index: ?

Size/arcmin: 12×9
Type: S

Radio: Asymmetric shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 73'' : S=2.0$ Jy).

G352.7-0.1

RA: $17^{\text{h}}27^{\text{m}}40^{\text{s}}$
Dec: $-35^{\circ}07'$

1-GHz flux/Jy: 4
Spectral index: 0.6

Size/arcmin: 8×6
Type: S

Radio: Distorted shell.

X-ray: Detected.

Distance: H_I absorption indicates 6.8 to 8.4 kpc.

References:

Clark *et al.* 1975, *AujPA*, 37, 1. Molonglo at 408 MHz ($3': S=9.6$ Jy) and Parkes 64-m at 5 GHz ($4': S=2.3$ Jy).
 Caswell *et al.* 1983, *MNRAS*, 203, 595. FIRST at 1415 MHz ($1'.1$).
 Dubner *et al.* 1993, *AJ*, 105, 2251. VLA at 1.47 GHz ($34'': S=3.4\pm 0.4$ Jy).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz ($43''\times 75'': S=4.4$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Kinugasa *et al.* 1998, *PASJ*, 50, 249. ASCA observations.
 Koralesky *et al.* 1998, *AJ*, 116, 1323. VLA search for OH emission.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
 Giacani *et al.* 2009, *A&A*, 507, 841. VLA at 4.8 GHz ($9''\times 12''$), plus H α and XMM observations.

G353.6–0.7

RA: $17^{\text{h}}32^{\text{m}}00^{\text{s}}$

Dec: $-34^{\circ}44'$

1-GHz flux/Jy: 2.5?

Spectral index: ?

Size/arcmin: 30

Type: S

Radio: Shell, brighter to S.

X-ray: Detected.

Point sources: Central X-ray source.

References:

Tian *et al.* 2008, *ApJ*, 679, L85. SGPS at 1.4 GHz ($100''$), plus 843 MHz ($43''$) and X-ray observations.
 Tian *et al.* 2010, *ApJ*, 712, 790. XMM, Suzaku and CO observations.
 Halpern & Gotthelf 2010, *ApJ*, 710, 941. XMM observations.
 Abramowski *et al.* 2011, *A&A*, 531, A81. γ -ray observations.
 Bamba *et al.* 2012, *ApJ*, 756, 149. Suzaku observations.
see also: Bamba *et al.* 2012, *ApJ*, 761, 80. Erratum.
 Klochkov *et al.* 2013, *A&A*, 556, A41. XMM and other X-ray observations.

G353.9–2.0

RA: $17^{\text{h}}38^{\text{m}}55^{\text{s}}$

Dec: $-35^{\circ}11'$

1-GHz flux/Jy: 1?

Spectral index: 0.5?

Size/arcmin: 13

Type: S

Radio: Shell, with central double source.

References:

Green 2001, *MNRAS*, 326, 283. VLA at 327 MHz ($3'.0\times 2'.7$) and 1.4 GHz ($42''\times 36''$), plus 8.4 GHz ($8''4\times 6''.1$) of central source only.

G354.1+0.1

RA: $17^{\text{h}}30^{\text{m}}28^{\text{s}}$

Dec: $-33^{\circ}46'$

1-GHz flux/Jy: ?

Spectral index: varies

Size/arcmin: $15\times 3?$

Type: C?

Is this a SNR?

Radio: Elongated N--S.

Point sources: Pulsar at S tip.

References:

Frail *et al.* 1994, *ApJ*, 437, 781. VLA at 330 MHz ($47'' \times 99''$) and 1.4 GHz ($8''.8 \times 21''$).

G354.8–0.8

RA: $17^{\text{h}}36^{\text{m}}00^{\text{s}}$

Dec: $-33^{\circ}42'$

1-GHz flux/Jy: 2.8?

Spectral index: ?

Size/arcmin: 19

Type: S

Radio: Distorted shell.

References:

Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz ($43'' \times 78''$: $S=3.1$ Jy).

Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, *AJ*, 116, 1323. VLA search for OH emission.

G355.4+0.7

RA: $17^{\text{h}}31^{\text{m}}20^{\text{s}}$

Dec: $-32^{\circ}26'$

1-GHz flux/Jy: 5?

Spectral index: ?

Size/arcmin: 25

Type: S

Radio: Faint, incomplete shell.

References:

Gray 1994, *MNRAS*, 270, 835. MOST at 843 MHz ($43'' \times 77''$).

Roy & Bhatnagar 2006, *JPhCS*, 54, 152. GMRT at 330 MHz ($1'$: $S=8.9 \pm 1.3$ Jy).

G355.6–0.0

RA: $17^{\text{h}}35^{\text{m}}16^{\text{s}}$

Dec: $-32^{\circ}38'$

1-GHz flux/Jy: 3?

Spectral index: ?

Size/arcmin: 8×6

Type: S

Radio: Well defined shell.

X-ray: Centrally brightened.

References:

Gray 1994, *MNRAS*, 270, 847. MOST at 843 MHz ($43'' \times 80''$: $S=2.6$ Jy).

Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.

Roy & Bhatnagar 2006, *JPhCS*, 54, 152. GMRT at 330 MHz ($1'$: $S=3.3 \pm 0.5$ Jy).

Marquez-Lopez & Phillips 2010, *MNRAS*, 407, 94. Mid-IR observations.

Minami *et al.* 2013, *PASJ*, 65, 99. Suzaku observations.

G355.9–2.5

RA: $17^{\text{h}}45^{\text{m}}53^{\text{s}}$

Dec: $-33^{\circ}43'$

1-GHz flux/Jy: 8

Spectral index: 0.5

Size/arcmin: 13

Type: S

Radio: Distorted shell, brightest to SE.

References:

Clark *et al.* 1975, *AujPA*, 37, 1. Molonglo at 408 MHz ($3'$: $S=12.3$ Jy) and Parkes 64-m at 5 GHz ($4'$: $S=3.4$ Jy).

Dubner *et al.* 1993, *AJ*, 105, 2251. VLA at 1.47 GHz ($32'' \times 34''$: $S=5.0 \pm 0.3$ Jy).

Gray 1994, *MNRAS*, 270, 835. MOST at 843 MHz ($43'' \times 77''$).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.

G356.2 + 4.5

RA: 17^h19^m00^s
Dec: −29°40′

1-GHz flux/Jy: 4
Spectral index: 0.7

Size/arcmin: 25
Type: S

Has been called G356.2+4.4.

Radio: Faint shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10′.4).
 Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (3′×1′.5 : S = 8.1±1.7 Jy), and NVSS at 1.4 GHz.

G356.3 – 1.5

RA: 17^h42^m35^s
Dec: −32°52′

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 20×15
Type: S

Radio: Double arc.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43″×79″ : S = 2.8 Jy).
 Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (0′.8×1′.7 : S = 5.7±0.2).

G356.3 – 0.3

RA: 17^h37^m56^s
Dec: −32°16′

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 11×7
Type: S

Has been suggested this part of a larger SNR.

Radio: Diffuse emission.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43″×81″ : S = 2.6 Jy).
 Roy & Pramesh Rao 2002, MNRAS, 329, 775. GMRT at 330 MHz (2′.7×4′.8).

G357.7 – 0.1

RA: 17^h40^m29^s
Dec: −30°58′

1-GHz flux/Jy: 37
Spectral index: 0.4

MSH 17–39
Size/arcmin: 8×3?
Type: ?

Has been suggested that this is not a SNR.

Radio: Multiple arcs and filaments, brighter to NW ‘head’.

X-ray: Detected from NW ‘head’, and SW ‘tail’.

Distance: H_i absorption suggests beyond Galactic Centre.

References:

Caswell *et al.* 1975, *AujPA*, 37, 39. Molonglo at 408 MHz ($3': S=54.2$ Jy) and Parkes 64-m at 5 GHz ($4': S=18.5$ Jy).

Milne & Dickel 1975, *AujPh*, 28, 209. Parkes 64-m at 5 GHz ($4'.4: S=14.6$ Jy).

Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'.6$).

Caswell *et al.* 1980, *MNRAS*, 190, 881. FIRST at 1415 MHz ($50''$).

Weiler & Panagia 1980, *A&A*, 90, 269. Effelsberg 100-m at 9 GHz ($1'.5$) (private communication from Baker).

Shaver *et al.* 1985, *Nature*, 313, 113. VLA at 1.4 GHz ($3''.8\times 10''.9$) and 5 GHz ($12''\times 26''$).

Becker & Helfand 1985, *Nature*, 313, 115. VLA at 1.4 GHz and 5 GHz.

Helfand & Becker 1985, *Nature*, 313, 118. Suggest it is not a SNR.

Shaver *et al.* 1985, *A&A*, 147, L23. Observations of peripheral compact source.

Caswell *et al.* 1989, *PASA*, 8, 184. MOST at 843 MHz ($43''\times 83''$).

Gray 1994, *MNRAS*, 270, 835. MOST at 843 MHz ($43''\times 84''$).

Stewart *et al.* 1994, *Apj*, 432, L39. ATCA at 4.79 and 5.84 GHz ($12''\times 22''$) and Effelsberg 100-m at 10.6 GHz ($1'$), including polarisation.

Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant, including masers.

Yusef-Zadeh *et al.* 1999, *Apj*, 527, 172. VLA of nearby OH masers.

LaRosa *et al.* 2000, *AJ*, 119, 207. VLA at 333 MHz ($43''\times 24''$).

Brogan *et al.* 2000, *Apj*, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

Brogan & Goss 2003, *AJ*, 125, 272. VLA at 1.4 GHz ($13''.6\times 11''.4$), including H_I, and 8.3 GHz ($14''.3\times 6''.8$) recombination line observation of H_{II} region.

Gaensler *et al.* 2003, *Apj*, 594, L35. Chandra detection.

Lazendic *et al.* 2003, *AN*, 324 (No 51), 157. Molecular line observations.

Burton *et al.* 2004, *MNRAS*, 348, 638. IR and radio observations of H_{II} region.

Lazendic *et al.* 2004, *MNRAS*, 354, 393. IR and molecular line observations.

Hewitt *et al.* 2008, *Apj*, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Phillips *et al.* 2009, *MNRAS*, 397, 1215. Observations of interactions with surroundings.

Castro *et al.* 2013, *Apj*, 774, 36. Fermi observations.

G357.7 + 0.3

RA: 17^h38^m35^s
Dec: -30°44'

1-GHz flux/Jy: 10
Spectral index: 0.4?

Size/arcmin: 24
Type: S

Radio: Non-thermal shell in complex region.

References:

Reich & Fürst 1984, *A&AS*, 57, 165. Effelsberg 100-m at 2.7 GHz ($4'.3: S=7\pm 1.5$ Jy), $S_{5\text{ GHz}}=5.5\pm 1.5$ Jy from surveys.

Gray 1994, *MNRAS*, 270, 835. MOST at 843 MHz ($43''\times 84''$).

Yusef-Zadeh *et al.* 1999, *Apj*, 527, 172. VLA of nearby OH masers.

Hewitt *et al.* 2008, *Apj*, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Phillips & Marquez-Lugo 2010, *MNRAS*, 409, 701. Spitzer observations of region.

G358.0 + 3.8

RA: 17^h26^m00^s
Dec: -28°36'

1-GHz flux/Jy: 1.5?
Spectral index: ?

Size/arcmin: 38
Type: S

Radio: Faint shell.

References:

Duncan *et al.* 1995, *MNRAS*, 277, 36. Parkes 64-m at 2.4-GHz ($10'.4$).

Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz ($2'2 \times 1'3$: $S = 2.5 \pm 1.3$ Jy), and NVSS at 1.4 GHz.

G358.1 + 0.1

RA: $17^{\text{h}}37^{\text{m}}00^{\text{s}}$
Dec: $-29^{\circ}59'$

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 20
Type: S

Radio: Faint shell.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 77''$).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1'$: $S = 6.0 \pm 2.5$ Jy).

G358.5 – 0.9

RA: $17^{\text{h}}46^{\text{m}}10^{\text{s}}$
Dec: $-30^{\circ}40'$

1-GHz flux/Jy: 4?
Spectral index: ?

Size/arcmin: 17
Type: S

Radio: Shell, brighter to NE.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 77''$).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1'$: $S = 8.0 \pm 2.5$ Jy).

G359.0 – 0.9

RA: $17^{\text{h}}46^{\text{m}}50^{\text{s}}$
Dec: $-30^{\circ}16'$

1-GHz flux/Jy: 23
Spectral index: 0.5

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Optical: Detected.

X-ray: Partial shell.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4'3$).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 86''$).
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz ($43'' \times 24''$).
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz ($8''4 \times 12''8$) of part.
 Bamba *et al.* 2009, ApJ, 691, 1854. Suzaku observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G359.1–0.5**RA:** 17^h45^m30^s**Dec:** –29°57′**1-GHz flux/Jy:** 14**Spectral index:** 0.4?**Size/arcmin:** 24**Type:** S**Radio:** Non-thermal shell in complex region, crossed by the ‘snake’.**Optical:** Detected.**X-ray:** Centrally brightened.**Point sources:** Several compact radio sources near centre, OH masers around edge.**References:**Downes *et al.* 1979, A&AS, 35, 1. From observations by Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6″:S=13 Jy).

Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz (4′3″:S=10±1.5 Jy) and 4.8 GHz (2′4″:S=8.1±0.5 Jy).

Uchida *et al.* 1992, ApJ, 398, 128. VLA at 1.5 GHz (11″×10″), and observations of nearby molecular material.Uchida *et al.* 1992, AJ, 104, 1533. VLA at 1.4 GHz.

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43″×85″).

Yusef-Zadeh *et al.* 1995, Science, 270, 1801. VLA at 1.4 GHz (33″×31″), and 1.7 GHz for OH survey.LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43″×24″).Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.Lazendic *et al.* 2002, MNRAS, 331, 537. Observations of shocked molecular gas where the ‘snake’ crosses the remnant.Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8″4×12″8).Aharonian *et al.* 2008, A&A, 483, 509. XMM and γ -ray observations.Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.Bamba *et al.* 2009, ApJ, 691, 1854. Suzaku observations.Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.Ohnishi *et al.* 2011, PASJ, 63, 527. Suzaku observations.**G359.1+0.9****RA:** 17^h39^m36^s**Dec:** –29°11′**1-GHz flux/Jy:** 2?**Spectral index:** ?**Size/arcmin:** 12×11**Type:** S**Radio:** Shell, brightest in E.**References:**

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43″×88″:S=4.3 Jy).

Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1′:S=4.3±1.0 Jy).

Law *et al.* 2008, ApJS, 177, 515. VLA at 1.4 GHz (10′9″×15′9″:S=1.3±0.5 Jy).